

## RESEARCH COMMUNICATION

# Suppressive Effects of Okinawan Food Items on Free Radical Generation from Stimulated Leukocytes and Identification of Some Active Constituents: Implications for the Prevention of Inflammation-associated Carcinogenesis

Akira Murakami\*, Hisashi Ishida, Kimie Kubo, Ikuyo Furukawa, Yasutaka Ikeda, Megumi Yonaha<sup>1</sup>, Yohko Aniya<sup>2</sup>, Hajime Ohigashi

## Abstract

Okinawa prefecture in Japan is a distinct area characterized by unique traditional food habits and longevity. Prolonged exposure to activated leukocytes, playing pivotal roles in chronic inflammation-associated carcinogenesis, is known to lead to oxidative and nitrosative damage to macromolecules in the body since they are primary sources of free radicals, such as superoxide anion ( $O_2^-$ ) and nitric oxide (NO). In this study, we estimated anti-oxidative and anti-nitrosative activities of Okinawan food items by employing two cellular experimental systems: (1) phorbol ester-induced  $O_2^-$  generation from differentiated HL-60 human promyelocytic leukemia cells; and (2) lipopolysaccharide (LPS)-induced NO generation in RAW264.7 murine macrophages. A total of 138 food items, consisting of 42 samples unique to Okinawa and 96 common in the Japanese main island, were purchased at local markets in Okinawa and extracted with chloroform. When tested at a concentration of 100  $\mu\text{g/ml}$ , 38% (16/42) of the former showed 70% or more inhibition of  $O_2^-$  generation while 21% (20/96) of the latter did so. In parallel, 64% (27/42) of the former showed significant NO generation suppression in contrast to 48% (46/96) of the latter. Twenty-one active species were further tested at a concentration of 20  $\mu\text{g/ml}$ , and eleven species, including sugar cane, wild turmeric, and zedoary, were indicated to be most promising items with anti-oxidative and anti-nitrosative properties. In addition, some of the active constituents (chebulagic acid, a resveratrol derivative, and sesquiterpenoids) were identified. Our results suggest that food items typical in the Okinawa area have higher cancer preventive potential than those common in Japan.

**Key Words:** Free radicals - Okinawa foodstuffs - inflammatory cells - screening

*Asian Pacific J Cancer Prev*, 6, 437-448

## Introduction

It is known that plants have potentials to biosynthesize certain chemicals in response to environmental stress stimuli such as ultraviolet (UV) exposure, invading insects, bacteria, and viruses. For instance, they are forced to biosynthesize flavonoids and carotenoids to quench or scavenge free radicals derived from UV light exposure. Based on this principle, it can be hypothesized that the plants exposed to harsher stress must produce more biologically active phytochemicals in terms with quality and quantity. In fact, we have previously shown that extracts from edible plants

in subtropical Southeast Asian countries, exposed to intense sun light and thriving microorganisms, show much higher anti-tumor promoting activity as detected by phorbol ester-induced EB virus activation as compared with those common in Japan (Koshimizu, 1988; Murakami, 1995, 1998, 2000a). Also it is important to note that activity-guiding separation of some of the active extracts resulted in isolation and identification of the active compounds with notable cancer preventive potentials and reasonable molecular mechanisms (Nakamura, 1998, 1999, 2000; Murakami, 1996, 2002, 2003, 2004; Tanaka, 1997ab, 2001; Ohata, 1998; Kobayashi, 1998; Miyauchi, 2000; Mofatt, 2000, 2002; Ito, 2004, 2005;

*Division of Food Science and Biotechnology, Graduate School of Agriculture, Kyoto University, Kyoto 606-8502; <sup>1</sup>Department of Research and Development, Ryukyu Bio-resource Development, Co. Ltd., Uruma city; <sup>2</sup>Laboratory of Molecular Pharmacology, Graduate School of Medicine, University of the Ryukyus, Nishihara, Okinawa, Japan \*To whom correspondence should be addressed. FAX: +81-75-753-6281; email: cancer@kais.kyoto-u.ac.jp*

Ichikawa, 2005; Takada, 2005).

Both endogenous (such as the arachidonate cascade, cytochrome P450, xanthine oxidase, and NADPH oxidase) and exogenous (such as UV light, heavy metals, and food types) oxidative stress are believed to be highly associated with carcinogenesis processes. In immune systems, neutrophils and macrophages are activated in response to exogenous stimuli, including microorganism invasion, to generate free radicals for host protection. It is important, however, to understand that imbalances of redox regulation and prolonged radical generation have been observed in some chronic inflammatory diseases of the digestive tract (Babbs, 1992; Nieto, 2000; Yamada, 1991). In fact, the activated leukocyte-derived oxidative damage may be involved in carcinogenesis by activating procarcinogens (Trush, 1985), participating in malignant transformation (Weitzman, 1985,1990), interacting with human epithelial cells (Kamp, 1989), and modulating DNA bases (Shacter, 1998; Shen, 2000). The NADPH oxidase system is present in inflammatory cells and consists of multiple components to produce superoxide anion ( $O_2^-$ ) (Kobayashi, 2001). Beyond its action as a precursor of various reactive oxygen species,  $O_2^-$  itself is involved in several signal transduction pathways (Barrett, 1999; Lee, 1998; Purohit, 1994; Sakai, 1994), while  $O_2^-$  generation plays some important roles in colonic inflammation (Tardieu, 2000), cellular transformation (Nakamura, 1998), and carcinogenesis (Schwarz, 1984). On the other hand, there is now ample evidence that excess and/or prolonged generation of nitric oxide (NO) causes carcinogenesis in digestive organs (Goto, 1999; Ambs, 1998). NO is drastically released by the inducible enzyme, inducible NO synthase (iNOS). Moreover, it is of paramount importance to note that peroxynitrite, formed by a non-enzymatic reaction of  $O_2^-$  with NO, is markedly mutagenic and responsible for tumor development (Maeda, 1998). Collectively, suppression of leukocytic free radical generation may reduce the risks of carcinogenesis.

The above mentioned background information led us to examine the suppressive effects of Okinawan food items on  $O_2^-$  and NO generation from stimulated leukocytes, i.e., respectively differentiated HL-60 and RAW264.7 macrophages. As we had expected, their anti-oxidative and anti-nitrosative activities were remarkably higher than those common in Japan, supporting our hypothesis that food items from subtropical areas are promising and interesting sources of cancer preventive agents based on the differences in environmental conditions. Additionally, the isolation and identification of some active constituents are described.

## Materials and Methods

### Chemicals

12-*O*-Tetradecanoylphorbol-13-acetate (TPA) was obtained from Research Biochemicals International (Natick, MA). Dulbecco's modified eagle medium (DMEM), RPMI 1640 medium, and fetal bovine serum (FBS) were purchased from Gibco RBL (Rockville, NY). Lipopolysaccharide (LPS,

*E. coli* serotype 0127, B8) was purchased from Difco Labs (Detroit, MI) and interferon (IFN)- $\gamma$  from Genzyme (Cambridge, MA). Cytochrome C was obtained from Sigma (St. Louis, MO). Chebulagic acid was donated from Dr. Toshio Ichiba, Okinawa Industrial Technology, Center, Okinawa, Japan. All other chemicals were purchased from Wako Pure Chemical Industries (Osaka, Japan), unless specified otherwise.

### Cells

HL-60 cells (human promyelocytes) and RAW 264.7 cells (mouse macrophages) were purchased from American Type Culture Collection (Rockville, MA) and cultured respectively in RPMI and DMEM media. Ten percent FBS, 200 U/ml penicillin, and 250 ng/mL streptomycin were added to each, and the cell lines were maintained at 37°C in a 5% CO<sub>2</sub> atmosphere.

### Food Samples

Each sample, purchased at local markets in Okinawa in 2002, was cut into small pieces and soaked in chloroform followed by one-week extraction.

### $O_2^-$ Generation in Differentiated HL-60 Cells

An inhibitory test of TPA-induced  $O_2^-$  generation was performed as previously reported (Murakami, 2002). Briefly, HL-60 cells ( $5 \times 10^5$  cells/ml) were incubated in 1.3% dimethylsulfoxide (DMSO) in RPMI 1640 medium for 6 days. Differentiated HL-60 cells ( $1 \times 10^6$ ) were incubated with the test compound, then dissolved in DMSO (0.5%, v/v) and 1 mL of Hank's buffer for 15 minutes. After washing, the cells were further incubated with 100 nM TPA and 1 mg/mL cytochrome C at 37°C for 60 minutes. Cell viability was measured by trypan blue-dye exclusion test. The level of extracellular  $O_2^-$  was then measured by a cytochrome C reduction method. Cells treated with only the vehicle plus cytochrome C and with TPA plus cytochrome C were used as negative and positive controls, respectively.

### NO<sub>2</sub><sup>-</sup> Production

A suppressive test of NO generation was done as previously reported (Murakami, 2002). RAW 264.7 cells ( $8 \times 10^5$  cells/2 mL) in a 60-mm dish were treated with LPS (100 ng/mL), tetrahydrobiopterin (10 mg/mL), IFN- $\gamma$  (100 U/mL), L-arginine (2 mM), and specified concentrations of the test compound dissolved in DMSO (0.5%, v/v). After 12 hours, the levels of nitrite and cytotoxicity were measured by Griess and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assays, respectively. Cells treated with only the vehicle and with LPS/IFN- $\gamma$  were used as negative and positive controls, respectively.

### Inhibitor Rate (IR) and Cell Viability (CV)

Each experiment was done in duplicate, and the data are expressed by the mean value. The standard errors did not exceed 10% of the means (data not shown). The IR in each assay was calculated using the following equation: IR (%)

=  $\{1 - [(test\ sample\ data) - (negative\ control\ data)] / [(positive\ control\ data) - (negative\ control\ data)]\}^{-1} \times 100$ . Similarly, CV was calculated using the following equation:  $CV (\%) = \{1 - [(test\ sample\ data) - (negative\ control\ data)] / [(positive\ control\ data) - (negative\ control\ data)]\}^{-1} \times 100$ . The statistical significance of differences between groups in each assay was assessed by a Student's t-test (two-sided) that assumed unequal variance.

#### Isolation and Identification of Active Constituents

##### 3-methoxy-5-hydroxystilbene

Dried leaves of *Alpinia speciosa* K. schum (1 kg) was extracted with chloroform at room temperature for a week. The extract was filtered and concentrated in vacuo using an evaporator, and then the concentrate (28 g) was subjected to a column chromatography on Wakogel C-200 (increasing the volume of ethyl acetate in *n*-hexane as elute). Eighty-percent ethyl acetate fraction (3.3 g) was then subjected to a reverse phase (C<sub>18</sub>), medium pressure column chromatography on ODS gel (Yamamura Chemical, Kyoto, Japan), increasing the volume of methanol in H<sub>2</sub>O as elute. Fifty-percent methanol fraction (106 mg) was purified by preparative thin layer chromatography (25% ethyl acetate in *n*-hexane) and preparative high performance liquid chromatography (HPLC) (column, J'sphere, Yamamura Chemical; mobile phase, 60% acetonitrile in H<sub>2</sub>O; flow rate, 7.0 ml/min; detection, 254 nm; retention time, 11.0 min) to afford 3-methoxy-5-hydroxystilbene (purity > 95%). The chemical identity was confirmed by comparing its spectral data (nuclear magnetic resonance, NMR, and mass spectrometry, MS) with those previously reported (Koon-Shin, 1998).

##### Sesquiterpenes

*Curcuma longa* L. (1 kg) was extracted with chloroform at room temperature. The extract was concentrated in vacuo followed by partition between chloroform and water. The chloroform layer thus obtained (3.4 g) was subjected silica gel column chromatography which was eluted with increasing amounts of ethyl acetate in *n*-hexane. The 2.5% ethyl acetate fraction (800 mg) was further separated on silica gel (0-5% ethyl acetate in *n*-hexane), which had been pretreated with 5% AgNO<sub>3</sub> in water and dried. This step allowed us to purify β-atlantone (29 mg). Each fraction was subjected to HPLC (column, YMC ODS AQ-302, Yamamura Chemical; mobile phase, 85% acetonitrile in H<sub>2</sub>O; flow rate, 1.0 ml/min; detection, 254 nm) to purify sesquiterpenes (retention time, min; and yield, mg): ar-turmerone (4.0, 130 mg), germacrone (4.8, 61 mg), β-turmerone (5.2, 157 mg), and α-turmerone (5.3, 100 mg). The retention time of β-atlantone in these HPLC conditions was 5.3 min, and could hardly be separated from those for α- and β-turmerones. The chemical identity of isolated sesquiterpenes was confirmed by comparing their spectral data (NMR and MS) with those previously reported (Yang, 2005; Li, 2003; Itokawa, 1984; Golding, 1992). The purity of each compound was > 95%.

## Results and Discussion

### Screening of Okinawan Food Items

A total of 138 food samples, collected in Okinawa prefecture, consists of 47 families of vegetables, fruits, seaweed, and others. Eight major families are Leguminosae, Solanaceae (10 samples each), Compositae, Rutaceae (9 samples each), Umbelliferae (8 samples), Cucurbitaceae, Gramineae, and Zingiberaceae (7 samples each), and Cruciferae (6 samples) (Table 1). These samples were extracted with chloroform and anti-oxidative and anti-nitrosative activities were evaluated at a concentration of 100 μg/mL for suppression of phorbol ester-induced O<sub>2</sub><sup>-</sup> generation in DMSO-differentiated HL-60 cells and that of LPS-induced NO generation in RAW264.7 macrophages, respectively. It is notable that both Gramineae and Zingiberaceae samples showed relatively high frequency of marked suppressive activity (70% > IR) toward both O<sub>2</sub><sup>-</sup> and NO generation (5/7, 71% for O<sub>2</sub><sup>-</sup> and 6/7, 86% for NO, with respect to each family). These two promising families are followed by Cruciferae (3/6, 50% for O<sub>2</sub><sup>-</sup> and 6/6, 100%) and Cucurbitaceae (2/7, 29% for O<sub>2</sub><sup>-</sup> and 5/7, 71% for NO). Interestingly, the samples of Umbelliferae has a quite contrastive profile for free radical generation suppression since one out of 8 samples (13%) markedly attenuated O<sub>2</sub><sup>-</sup> generation, whereas a high rate (8/8, 100%) was seen for NO generation reduction, suggesting that NO suppressive constituents widely occur in this family, but they have little ability to attenuate O<sub>2</sub><sup>-</sup> generation. In this context, it is intriguing to point out that the Umbelliferae plants abundantly biosynthesize the coumarin-related compounds which have been to demonstrated by us to be able to efficiently suppress NO generation in macrophages (Murakami, 1999, 2000b).

Then we classified all samples tested into those commonly seen in Japanese main island and those typical in Okinawan area. Test samples showing 70% or more suppression are designated as 'effective sample'. Figure 1 shows the results of the classification with respect to the suppression of O<sub>2</sub><sup>-</sup> (panel A), NO (panel B) and both (panel C), in which the portion of 'effective samples' are hatched in black. It should be noted that Okinawan samples exhibit markedly higher rates of 'effective samples' in each evaluation system. Those results are consistent with our previous work reporting that the extracts from subtropical foods exhibited higher in vitro anti-tumor promoting activity than those from Japanese common ones (Koshimizu, 1988; Murakami, 1995, 1998, 2000a), implying again that subtropical plants produce biologically active chemicals in terms of quantity and quality to protect themselves from invading microorganisms and insects as well as photo-damages by sunlight.

Since the numbers of effective samples in O<sub>2</sub><sup>-</sup> and NO assay systems are 31 and 79, respectively, second screening tests, in which sample concentration was decreased to 20 and 4 μg/mL, were done to focus on promising samples which deserve the isolation and identification of active

**Table 1. Suppressive Effects of Okinawan and Common Food Items on TPA-induced O<sub>2</sub>- generation in Differentiated HL-60 Cells and LPS/IFN- $\gamma$ -induced NO Generation in RAW265.7 Cells**

Family/species	Common name	Part tested	O <sub>2</sub> - IR (%)	CV (%)	NO IR (%)	CV (%)
Actinidiaceae						
<i>Actinidia chinensis</i> Planch.	Kiwi Fruit	Fruit	16	87	100	100
Anacardiaceae						
<i>Mangifera indica</i> L.	Mango	Fruit	6.0	8.1	31	100
Auriculariaceae						
<i>Auricularia auricula</i> Underwood.	Cloud ear mushroom#	Whole Part	92	0	76	97
Araceae						
<i>Colocasia esculenta</i> Schott.	Taro (fresh)	Stalks	2.2	83	81	79
	Taro (powdered)	Stalks	0	91	54	94
Bromeliaceae						
<i>Ananas comosus</i> Merrill.	Pineapple*	Fruit	3.4	80	30	86
Caricaceae						
<i>Carica papaya</i> L.	Papaya*	Fruit	0	69	13	100
	Papaya*		4.2	76	86	99
Cheneopodeaceae						
<i>Chenopodium</i> L.	<i>Akino-wasuregusa</i> *	Roots	24	81	85	65
<i>var.centrorubrum</i> Mkino.	<i>Akino-wasuregusa</i> *	Stem	2.7	91	43	96
<i>Spinacia oleracea</i> L.	Spinach	Leaves, Stem	7.0	83	100	92
Coccoloba						
<i>Coccoloba umifera</i>	<i>Umi-budo</i> *	Whole Part	2.8	96	10	100
Combretaceae						
<i>Terminalia catappa</i>	Indian Almond*#	Leaves	91	0	100	34
	Indian Almond* (fermented)	Leaves	4.5	74	100	36
Compositae						
<i>Artemisia campestris</i> L.	<i>Ryukyu-yomogi</i> *	Leaves	55	81	100	34
<i>A. princeps</i> Pamp.	Worm wood	Leaves, Stem	3.4	81	87	51
<i>Asparagus officinalis</i> L.	Common Asparagus	Stem	4.6	89	47	100
<i>Carthamus tinctorius</i> L.	Safflower	Flower	27	67	100	34
<i>Chrysanthemum coronarium</i> L.	Garland	Leaves, Stem	0.9	51	100	41
<i>Cirsium japonicum</i>	Shima-gobo*	Fruit	3.9	81	17	55
<i>Glycyrrhiza bicolor</i> DC.	Two-Colored Gynura*	Leaves	1.2	81	0	84
<i>Ixeris dentate</i> Nakai.	Nigana*	Leaves, Stem	12	97	81	94
<i>Lactuca sativa</i> L.	Lettuce	Leaves	28	94	75	100
Convolvulaceae						
<i>Ipomoea aquatica</i> Forsk.	Water Convolvulus*	Leaves, Stem	2.6	57	77	64
<i>I. batatas</i>	<i>Beni-imo</i> (white)*	Stalks	36	96	100	100
	<i>Beni-imo</i> (brown)*	Stalks	42	76	78	77
<i>I batatas</i> Lam.	Sweet Potato	Root	6.0	86	63	100
Cruciferae						
<i>Brassica chinensis</i> L.	Qing gin cai	Leaves, Stem	6.7	73	100	51
<i>B. compestris</i> L.var.pervilids.	Komatsuna	Leaves, Stem	14	82	86	100
<i>B. jnucea</i>	Leaf Mustard	Leaves, Stem	0.4	89	60	45
<i>B. oleracea var.capitata</i> L.	Cabbage	Leaves	2.7	85	0	64
<i>Nasturtium officinale</i> RBr.	Creson#	Leaves, Stem	96	11	97	47
<i>Raphanus sativus</i>	Radish	Leaves	0.8	70	100	69
	Radish#	Sprout	93	93	100	28
	Radish#	Stem	92	24	100	31
Cucurbitaceae						
<i>Berincasa hispida</i> Cogn.	White Gourd	Fruit	1.4	89	66	100
<i>Cucubita moschata</i> Poir.	Pumpkin	Fruit	18	76	100	100
<i>var.meronaeformis</i>						
<i>Cucumis sabivus</i> L.	Cucumber	Fruit	31	80	80	41
<i>Lagenaria hispida</i> Hera.	Bottle Gourd#	Fruit	97	0	100	71
<i>Luffa cylindrical</i> Roem.	Sponge Gourd*#	Fruit	94	83	79	35
<i>Momordica charantia</i> L.	Balsam Pear*	Fruit	3.1	100	76	100
<i>M. grosvenori</i> S.		Stalks	0	74	7.4	69
Dioscoreaceae						
<i>Dioscorea japonica</i> Thunberg.	Yam	Stalks	29	30	89	62

Table 1. Continued

Family/species	Common name	Part tested	O <sub>2</sub> - IR (%)	CV (%)	NO IR (%)	CV (%)
<b>Elaeagnaceae</b>						
<i>Elaeagnus glabra</i> Thurb. ex Murray.	<i>Tsuru-gumi*</i>	Fruit	31	93	100	86
<b>Gracilariaceae</b>						
<i>Gracilaria blodgettii</i>	<i>Kubire*#</i>	Whole Part	99	0	100	36
<b>Gramineae</b>						
<i>Coix lachryma-jobi</i> L. <i>var.lachryma-jobi</i>	Job's Tear*#	Grain	99	48	100	18
<i>Oryza sativa</i> L.	Rice	Grain	100	42	39	100
	<i>Mochigome#</i>	Grain	91	20	94	95
	<i>Kurokome#</i>	Grain	98	66	98	100
<i>Cymbopogon citratus</i>	Lemongrass	Leaves	0	98	100	47
<i>Saccharum officinarum</i> L.	Sugar Cane*#	Stem	90	100	100	32
<i>Zizania Latifolia</i> Stapf	Manchurian Wild Rice*	Grain	43	70	100	44
<b>Hypneaceae</b>						
<i>Hypnea charoides</i> Lamour.	Carrageenan*	Whole Part	9.4	100	100	100
<b>Labiatae</b>						
<i>Mentha arvensis</i>	Cool Mint	Leaves	25	74	100	77
<i>Orthosiphon stamineus</i> Benth.	Cats Whiskers*	Leaves	63	68	100	71
<i>Perilla frutescens</i> Britton <i>var.crispa</i> Hassk.	Perilla #	Leaves	89	36	97	42
<b>Lauraceae</b>						
<i>Persea americana</i> Mill.	Avocado	Fruit	26	84	100	100
<b>Leguminosae</b>						
<i>Archis hypogena</i> L.	Peanut	Nuts	0.6	75	13	100
<i>Cassia obtusifolia</i> L.	Oriental Senna	Leaves	2.0	96	100	34
<i>Glycine max</i> Merr.	Soybean	Beans	6.0	87	24	51
		Sprout	4.9	93	91	100
<i>Phasedus vulgaris</i> L.	Common Bean	Pod	0	88	29	100
<i>P. vulgaris</i> L.	<i>Uzura-mame</i>	Pod	13	92	55	100
<i>Pisum sativum</i> L.	Pea	Beans	2.5	70	25	100
<i>Vaccinium uliginosum</i>	<i>Kuromame</i> bean	Beans	58	89	69	87
<i>Vigna angularis</i> Ohwi et.Chashi.	Adzuki bean	Beans	8.2	89	48	100
<i>V. radiata</i> Rwlczek.	Mung bean	Beans	7.0	99	15	70
<b>Liliaceae</b>						
<i>Allium bakeri</i>	<i>Shima-rakkyo</i>	Bulb	21	97	56	82
<i>A sativum</i> L.	Garlic	Bulb	20	75	83	100
		Leaves	0	81	100	59
<i>Aloe arborescens</i> Mill.	Aloe	Leaves	3.3	73	18	100
<b>Malvaceae</b>						
<i>Hibiscus mutabilis</i> L.	Hibiscus#	Flower	99	0	100	63
<b>Musaceae</b>						
<i>Musa acuminata</i> Colla L.	Banana	Fruit	1.9	92	3.8	100
<b>Myrtaceae</b>						
<i>Eucalyptus globules</i> Labill.	Blue Gum#	Leaves	89	0	100	75
<i>Psidium guajava</i> L.	Guava (white)*	Fruit	1.8	80	27	100
	Guava (pink)*	Fruit	13	100	23	100
	Guava*#	Leaves	91	0	100	79
<b>Oxalidaceae</b>						
<i>Arerrhoa carambola</i> L.	Star Fruit	Fruit	14	100	27	14
<b>Passiflora</b>						
<i>Passiflora edulis</i> Sims.	Passion Fruit	Fruit	0.6	93	12	100
<b>Pedaliaceae</b>						
<i>Sesamum indicum</i> L.	Sesame (white)	Seeds	0	84	22	71
	Sesame (black)	Seeds	0	96	15	84
<b>Pinaceae</b>						
<i>Annina atemoya</i> hort.	Custard apple	Fruit	98	0	65	52
<b>Piperaceae</b>						
<i>Piper retrofractum</i> Vahl.	Hihatsu-modoki	Seeds	3.6	82	76	72

(Continued next page)

**Table 1. Continued**


Family/species	Common name	Part tested	O <sub>2</sub> - IR (%)	CV (%)	NO IR (%)	CV (%)
Plumbaginaceae						
<i>Limonium wrightii</i> O.K.	Ukon-isomatsu*#	Spray	88	100	96	14
Rhodomelaceae						
<i>Digenea simplex</i> C.Ag.	Digenea*#	Whole Part	77	0	100	29
Rosaceae						
<i>Eriobotrya japonica</i> Lindl.	Loquat	Fruit	33	60	54	100
<i>Prunus campanulata</i> Maxim	<i>Kanhi-zakura</i> *	Fruit	10	81	66	100
Rutaceae						
<i>Citrus deressa</i>	Shekwasha*	Fruit	0	91	17	100
		Juice	1.4	88	18	78
<i>C. limon</i> Burm.	Lemon#	Fruit	79	0	100	62
<i>C. natuaidai</i> Hayata.	Natsudaikai Orange	Fruit	0	74	12	68
<i>Citrus nobilis</i>	Marcott	Fruit	2.6	93	4.1	68
<i>C. tamurana</i> hort.ex Tanaka.	Hyuganatsu	Fruit	1.2	87	0	85
<i>C. tankan</i> Hayata.	Tankan	Fruit	4.4	93	0	90
<i>Fortunella crassifolia</i> Swingle.	Kumquats	Fruit	6.7	94	36	100
<i>Toddalia asiatica</i> (L.) Lam.	<i>Sarukake-mikan</i> *	Bark	21	98	100	48
Sapotaceae						
<i>Pouteria campechiana</i> (HBK) Baehni.	Canistel	Fruit	2.7	100	16	85
Saururaceae						
<i>Houttuynia cordata</i> Thunb.	Dokudami	Leaves	27	93	100	79
Sargassum						
<i>Sarugassum fusiforma</i> (Harvey) Setchell.	Yunabaru-hijiki*	Whole Part	11	80	100	41
Solanaceae						
<i>Capsicum annuum</i> CV.	Paprica (red)	Fruit	3.7	97	0	100
	Paprica (yellow)	Fruit	1.8	91	71	34
<i>C. annuum</i> var.grossum.Sendt.	Bell Pepper	Fruit	3.4	92	10	99
<i>C. annuum</i> L.var.angulosum Mill.	Sweet pepper	Fruit	9.0	93	39	100
<i>Lycium barbarum</i> L.	Chinese Wolfberry	Seed	8.5	84	75	100
<i>Lycopersicon esculentum</i> Mill.	Tomato	Fruit	2.3	92	92	84
	Tomato Cherry	Fruit	0.6	87	100	22
<i>Physalis alkekengi</i> L.var.dulce Pers.	Japanese Lantern Plant	Fruit	4.0	82	5.4	100
<i>Solanum melongena</i> L.	Eggplant	Fruit	0	86	53	100
<i>Solanum tuberosum</i> L.	Potato	Stem	0	71	42	70
Spermatochneae						
<i>Nemacystis decipiens</i> Kuckuck.	<i>Mozuku</i> #	Whole Part	96	0	100	59
Theaceae						
<i>Camelia sinensis</i> O.Kuntze.	Tea	Leaves	15	71	65	93
Tiliaceae						
<i>Corchorus olitorius</i>	Jew's mallow	Leaves	48	70	79	53
Tricholomataceae						
<i>Lentinus edodes</i> Singer.	Shiitake mushroom#	Whole Part	85	0	100	60
<i>Lyophyllum decastes</i> .	Shimeji mashroom#	Whole Part	91	100	100	13
Ulmaceae						
<i>Celtis sinensis</i> var japonica	Chinese Hackberry	Whole Part	98	0	59	45
Umbelliferae						
<i>Apium Graveolens</i> L.var.dulce Pers.	Celery	Leaves, Stem	1.8	95	86	42
<i>Chryptotaenia japonica</i> Hassk.	Japanese Hornwort	Leaves, Stem	68	49	87	31
<i>Daucus carota</i> L.	Carrot	Root	7.2	91	98	37
	<i>Shima-ninjin</i> *	Root	23	85	100	91
	<i>Kyo-ninjin</i>	Root	7.0	85	100	38
<i>Foericulum vulgare</i> Mill.	Fennel	Leaves, Stem	10	67	88	51
<i>Peucedanum japonicum</i> Thurb.	Botan-boufu*#	Root	100	50	91	49
	Botan-boufu*	Stem	67	89	100	34
Vitaceae						
<i>Vitis vinifera</i> L.	<i>Kyoho</i>	Fuit	0	78	8.4	77
Zingiberaceae						
<i>Alpinia speciosa</i> K. Schum.	Shell Flower	Leaves	21	81	100	35

(Continued next page)


Table 1. Continued

Family/species	Common name	Part tested	O <sub>2</sub> <sup>-</sup> IR (%)	CV (%)	NO IR (%)	CV (%)
	(fermented)*#	Leaves	90	45	100	38
	(powdered)*	Leaves	44	70	100	33
<i>Curcuma aromatica</i> Salisb.	Wild Turmeric*#	Root	94	19	93	61
<i>C. longa</i> L.	Turmeric*#	Root	94	0	76	52
<i>C. zedoaria</i> Rosc.	Zedoary*#	Root	89	0	100	35
<i>Zingiber officinale</i> Rosc.	Ginger	Root	90	14	51	41
Miscellaneous						
	<i>To-fuyo</i> *		54	93	95	90
	<i>To-fuyo</i> (liquid)*		9.0	93	32	100
<i>Laticauda semifasciata</i>	Sea snake (dried) *	Whole	5.0	93	41	93

Differentiated HL-60 cells ( $1 \times 10^6$ ) were incubated with the test compound, then dissolved in DMSO (0.5%, v/v) and 1 mL of Hank's buffer for 15 minutes. After washing, the cells were further incubated with 100 nM of TPA and 1 mg/mL of cytochrome C at 37°C for 60 minutes. Cell viability (CV) was measured using a trypan blue-dye exclusion test. The level of extracellular O<sub>2</sub><sup>-</sup> was then measured with a cytochrome C reduction method. Cells treated with only the vehicle plus cytochrome C and those with TPA plus cytochrome C were used as negative and positive controls, respectively. RAW 264.7 cells ( $8 \times 10^5$  cells/2 mL) in a 60-mm dish were treated with LPS (100 ng/mL), tetrahydrobiopterin (10 mg/mL), IFN- $\gamma$  (100 U/mL), L-arginine (2 mM), and specified concentrations of the test compound dissolved in DMSO (0.5%, v/v). After 12 hours, the levels of nitrite and CV were measured using Griess and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assays, respectively. Cells treated with only the vehicle and those with LPS/IFN- $\gamma$  were used as negative and positive controls, respectively. #Samples that showed 70% suppression or more in both assay systems. \*Food items that specifically occur in Okinawa prefecture. Data are shown as averages from the results of duplicate experiments.



**Figure 1. Comparison of Suppressive Potencies of Okinawan and Common Food Items on Free Radical Generation from Stimulated Leukocytes.** The proportions of food item extracts which showed 70% or more suppression of O<sub>2</sub><sup>-</sup> (Panel A), NO (Panel B), and both (Panel C) are shown as the hatched area in each graph.



**Figure 2. Chemical Structures of Phytochemicals Examined in an LPS-induced NO Generation Test.** Origins as follows: chebulagic acid (*Terminalia catappa*), 3-methoxy-5-hydroxystilbene (*Alpinia speciosa* K. schum), resveratrol (grape, etc.), ar-turmerone,  $\alpha$ - and  $\beta$ -turmerone,  $\beta$ -atlantone, germacrone, and curcumin (*Curcuma longa* L.).

principles. Twenty-three samples which showed 70% suppression or more in both assays were selected for secondary tests. As the results, 13 samples showed 70% suppression or more toward  $O_2^-$  generation at a concentration of 20  $\mu\text{g/ml}$  and only sugar cane and wild turmeric (4  $\mu\text{g/ml}$ ) attenuated  $O_2^-$  generation by 50% or more (Table 2). On the other hand, most of the samples (21/23, 91%) suppressed NO generation by 70% or more at a concentration of 20  $\mu\text{g/ml}$ , and 39% (9/23) of the total were still suppressive at a concentration of 4  $\mu\text{g/ml}$ . The present results, together with our previous extensive screening for the anti-tumor promoting activities (Koshimizu, 1988; Murakami, 1995, 1998, 2000a) and anti-oxidation activities (Kim, 1998, 2002) of Asian food items, may provide a useful database to search for food materials and its constituents which possess notable cancer preventive potentials derived from their anti-inflammatory properties. To support this notion, it can be indicated that we have so far reported several phytochemicals which are functionally novel and exhibited significant cancer preventive activity in a series of rodent models (Nakamura, 1998, 1999, 2000; Murakami, 1996, 2003, 2004; Tanaka, 1997ab, 2001; Kobayashi, 1998; Miyauchi, 2000). Of great

importance, those agents have been proven to show biochemical properties of  $O_2^-$  and NO generation suppression in vitro and in vivo (Nakamura, 1998, 1999, 2000; Murakami, 1996, 2002, 2003, 2004; Ohata, 1998). Thus, it is worth making an attempt to isolate and identify the active principles occurring in highly promising food samples listed in Table 2.

#### The Active Principles Which Suppress NO Generation

Based on the present results, literature information, and sample availability, we attempted to identify the active components of three food items, i.e., *getto* (*Alpinia speciosa* K. Schum, Zingiberaceae), turmeric (*Curcuma longa* L., Zingiberaceae), and momotamana (*Terminalia catappa*, Combretaceae). The Spectroscopic data, including those from  $^1\text{H-NMR}$  and MS analyses, they were revealed to be 3-methoxy-5-hydroxystilbene, five known sesquiterpenes, and chebulagic acid (an ellagitannin), respectively, by comparing them with previously published data (Koon-Shin, 1998; Yang, 2005; Li, 2003; Itokawa, 1984; Golding, 1992). 3-Methoxy-5-hydroxystilbene has been isolated another species (*Alpinia katsumadai*, Koon-Shin, 1998) but isolated

**Table 2. Suppressive Activities Toward Free Radical Generation by Okinawan and Common Food Items in the Second Set of Screening Tests**

Common name	Superoxide				Nitric oxide			
	% suppression		% cell viability		% suppression		% cell viability	
	20	4	20	4	20	4	20	4 ( $\mu\text{g/mL}$ )
Blue gum	85	45	28	100	100	100	35	94
<i>Botan-boufu</i> (stem)*	33	4.6	63	100	100	89	62	100
<i>Botan-boufu</i> (root)*	95	29	60	100	100	100	37	16
Bottle gourd	95	0	100	100	93	21	92	100
Cresson	100	40	100	100	100	100	36	87
Digenea*	100	4.5	100	99	100	3.6	86	97
<i>Getto</i> *	56	4.8	100	100	100	74	55	60
<i>Getto</i> * (fermented)	39	2.7	100	100	100	76	27	93
Hibiscus*	82	12	80	100	75	0	100	0
Indian almond*	51	14	100	100	100	83	43	77
Indian almond* (fermented)	44	26	73	100	100	76	44	100
Job's tear*	44	0.6	100	100	100	45	53	100
<i>Kubire</i> *	89	13	100	100	92	23	85	84
Mitsuba	68	0.8	92	93	100	100	33	38
<i>Moasa</i> *	95	24	79	100	100	19	67	86
Radish (sprout)	49	12	100	100	100	56	71	98
Radish (stem)	77	0	63	100	100	69	50	80
Rice (black)	52	0	54	100	52	0	90	0
Rice ( <i>mochigome</i> )	41	6.5	100	100	21	10	96	100
<i>Shiitakemashroom</i>	100	1.1	0	100	86	19	92	53
Sugar cane*	100	55	100	100	100	89	92	76
Wild turmeric*	100	68	8.2	100	100	40	39	91
Zedoary*	81	18	86	100	100	20	62	92

HL-60 cells ( $5 \times 10^5$  cells/ml) were incubated in 1.3% dimethylsulfoxide (DMSO) in RPMI 1640 medium for 6 days. Differentiated HL-60 cells ( $1 \times 10^6$ ) were incubated with the test compound, then dissolved in DMSO (0.5%, v/v) and 1 mL of Hank's buffer for 15 minutes. After washing, the cells were further incubated with 100 nM of TPA and 1 mg/mL of cytochrome C at 37°C for 60 minutes. Cell viability was measured using a trypan blue-dye exclusion test. The level of extracellular  $O_2^-$  was then measured with a cytochrome C reduction method. Cells treated with only the vehicle plus cytochrome C and those with TPA plus cytochrome C were used as negative and positive controls, respectively. RAW 264.7 cells ( $8 \times 10^5$  cells/2 mL) in a 60-mm dish were treated with LPS (100 ng/mL), tetrahydrobiopterin (10 mg/mL), IFN- $\gamma$  (100 U/mL), L-arginine (2 mM), and specified concentrations of the test compound dissolved in DMSO (0.5%, v/v). After 12 hours, the levels of nitrite and cytotoxicity were measured using Griess and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assays, respectively. Cells treated with only the vehicle and those with LPS/IFN- $\gamma$  were used as negative and positive controls, respectively. \*Food items specific to Okinawa prefecture. Data are shown as averages from the results of duplicate experiments.




from *A. speciosa* for the first time. A grape stilbene, resveratrol, and the yellow pigment in turmeric, curcumin, were also subjected to NO generation test because their ability to suppress NO generation has already been reported (Cho, 2002; Brouet, 1995) and because 3-methoxy-5-hydroxystilbene is structurally analogous to resveratrol.

The test agents were added to a RAW264.7 cell culture at the concentrations of 4, 20 and 100  $\mu\text{M}$ . As shown in Figure. 3A-C, chubulagic acid showed a concentration-dependent NO generation suppression without noticeable cytotoxicity. In accordance with the previous report (Cho, 2002), resveratrol markedly suppressed NO generation, and its analog, 3-methoxy-5-hydroxystilbene, was found to be cytotoxic at a higher concentration (Figurs. 3B and 3C). Curcumin abolished NO generation at a concentration of 20  $\mu\text{M}$  whereas it was highly cytotoxic at 100  $\mu\text{M}$  as detected by the value of  $\text{NO}_2^-$  concentration/cell viability (Figure. 3C). Germacrone, a sesquiterpenoid in turmeric, did not show substantial suppression even at a concentration of 100  $\mu\text{M}$ . NO generation suppressive potencies of  $\alpha,\beta$ -turmerone, ar-turmerone, and  $\beta$ -atlantone were nearly identical and comparable to that of curcumin. Yet, these sesquiterpene

are not cytotoxic even at a concentration of 100  $\mu\text{M}$ , in contrast to curcumin, suggesting that they are promising anti-inflammatory and anti-carcinogenic agents in turmeric. In addition, it can be pointed out that their total content in turmeric is comparable to that of curcumin (data not shown).

To our best knowledge, there has been no known reports on the physiological functions of 3-methoxy-5-hydroxystilbene. On the other hand, the NO generation suppressive effects of ar-turmerone, germacrone, and  $\beta$ -turmerone (Hong, 2002; Lee, 2002; Masuda, 1998) have recently and independently been described, but those of other sesquiterpenoids ( $\alpha$ -turmerone and  $\beta$ -atlantone) as well as simultaneous evaluation of those five sesquiterpene was done by us for the first time. Yoshikawa et al. demonstrated that the oral activity of germacrone for suppressing D-galactosamine- or tumor necrosis factor- $\alpha$ -induced liver injury in rats (Morikawa, 2002). In addition, an ethanol extract from turmeric, containing curcumin and sesquiterpenoids, significantly ameliorated symptoms of type 2 diabetes genetically diabetic KK-Ay mice (Kuroda, 2005; Nishiyama, 2005). Additionally, chebulagic acid has been shown to inhibit topoisomerase I (Hecht, 1992) and



**Figure 3. Effects of Selected Dietary Phytochemicals on LPS-induced NO Generation in RAW264.7 Macrophages.** Panel A, the concentrations of  $\text{NO}_2^-$  in the media 24 hours after LPS-exposure. Panel B, cell viability as detected by MTT assay. Panel C, the values of  $\text{NO}_2^-$  concentrations/cell viability which was corrected with the LPS-treated cells as 100. BL, blank; CA, chebulagic acid (20 and 100  $\mu\text{M}$ ); RES, resveratrol (20 and 100  $\mu\text{M}$ ); MHS, 3-methoxy-5-hydroxystilbene (20 and 100  $\mu\text{M}$ ); CUR, curcumin (4, 20 and 100  $\mu\text{M}$ );  $\beta\text{AT}$ ,  $\beta$ -atlantone (4, 20 and 100  $\mu\text{M}$ );  $\alpha\text{TM}$ ,  $\alpha$ -turmerone (4, 20 and 100  $\mu\text{M}$ );  $\beta\text{TM}$ ,  $\beta$ -turmerone (4, 20 and 100  $\mu\text{M}$ ); ATM, ar-turmerone (4, 20 and 100  $\mu\text{M}$ ); GER, germacrone (4, 20 and 100  $\mu\text{M}$ ).

xanthine oxidase (Fogliani, 2005), to exhibit tumor cell toxicity (Kashiwada, 1992), and to suppress collagen-induced arthritis (Lee, 2005). Together, both the sesquiterpenoids and chebulagic acid are attractive new agents which deserve evaluation of their cancer preventive efficacy.

### Concluding Remarks

In conclusion, we have shown that Okinawan food items have pronounced potentials to search for effective anti-oxidative new compounds as compared with commonly ingested food items in Japanese main island. In particular, the Gramineae and Zingiberaceae plants appear highly suppressive. Also, chebulagic acid, 3-methoxy-5-hydroxystilbene, and sesquiterpenes have been isolated as NO generation inhibitors from effective plants. Further efforts for identification and chemopreventive studies using those agents are underway in our laboratory.

### Acknowledgements

This work was supported by a grant from the Ministry of Agriculture, Forestry, and Fisheries (MAFF) Food Research Project 'Integrated Research on Safety and Physiological Function of Food' (To H.O.) and by a Grant-in-Aid for Cancer Research from the Ministry of Health, Labor and Welfare of Japan (To A.M).

### References

Ambs S, Merriam WG, Bennett WP, et al (1998). Frequent nitric oxide synthase-2 expression in human colon adenomas: implication for tumor angiogenesis and colon cancer progression. *Cancer Res*, **58**, 334-41.

Babbs CF (1992). Oxygen radicals in ulcerative colitis. *Free Radic, Biol Med*, **13**, 169-81.

Barrett WC, De Gnore JP, Keng YF, et al (1999). Roles of superoxide radical anion in signal transduction mediated by reversible regulation of protein-tyrosine phosphatase 1B. *J Biol Chem*, **274**, 34543-6.

Brouet I, Ohshima H (1995). Curcumin, an anti-tumour promoter and anti-inflammatory agent, inhibits induction of nitric oxide synthase in activated macrophages. *Biochem Biophys Res Commun*, **206**, 533-40.

Cho DI, Koo NY, Chung WJ, et al (2002). Effects of resveratrol-related hydroxystilbenes on the nitric oxide production in macrophage cells: structural requirements and mechanism of action. *Life Sci*, **71**, 2071-82.

Fogliani B, Raharivelomanana P, Bianchini JP, Bouraima-Madjebi S, Hnawia E (2005). Bioactive ellagitannins from *Cunonia macrophylla*, an endemic Cunoniaceae from New Caledonia. *Phytochemistry*, **66**, 241-7.

Golding BT, Pombo-Villar E (1992). Structures of  $\alpha$ - and  $\beta$ -turnerones. *J Chem Soc Perkin Trans*, **1**, 1519-24.

Goto T, Haruma K, Kitadai Y, et al (1999). Enhanced expression of inducible nitric oxide synthase and nitrotyrosine in gastric mucosa of gastric cancer patients. *Clin Cancer Res*, **5**, 1411-5.

Hecht SM, Berry DE, MacKenzie LJ, Busby RW, Nasuti CA (1992). A strategy for identifying novel, mechanistically unique

inhibitors of topoisomerase I. *J Nat Prod*, **55**, 401-13.

Hong CH, Noh MS, Lee WY, Lee SK (2002). Inhibitory effects of natural sesquiterpenoids isolated from the rhizomes of *Curcuma zedoaria* on prostaglandin E2 and nitric oxide production. *Planta Med*, **68**, 545-7.

Ichikawa H, Takada Y, Murakami A, Aggarwal BB (2005). Identification of a novel blocker of I $\kappa$ B kinase that enhances cellular apoptosis and inhibits cellular invasion through suppression of NF $\kappa$ B regulated gene products. *J Immunol*, **174**, 7383-92.

Ito K, Nakazato T, Murakami A, et al (2004). Induction of apoptosis in human myeloid leukemic cells by 1'-acetoxychavicol acetate through a mitochondrial- and Fas-mediated dual mechanism. *Clin Cancer Res*, **10**, 2120-30.

Ito K, Nakazato T, Xian MJ, et al (2005). 1'-Acetoxychavicol acetate (ACA) is a novel NF- $\kappa$ B inhibitor with significant activity against multiple myeloma in vitro and in vivo. *Cancer Res*, **65**, 4417-24.

Itokawa H, Hirayama F, Funakoshi K, Takeya K (1985). Studies on the antitumor bisabolane sesquiterpenoids isolated from *Curcuma xanthorrhiza*. *Chem Pharm Bull*, **33**, 3488-92.

Kamp DW, Dunne M, Weitzman, SA, Dunn MM (1989). The interaction of asbestos and neutrophils injures cultured human pulmonary epithelial cells: role of hydrogen peroxide. *J Lab Clin Med*, **114**, 604-12.

Kashiwada Y, Nonaka G, Nishioka I, Chang JJ, Lee KH (1992). Antitumor agents, 129. Tannins and related compounds as selective cytotoxic agents. *J Nat Prod*, **55**, 1033-43.

Kim OK, Murakami A, Nakamura Y, Ohigashi H (1998). Screening of edible Japanese plants for nitric oxide generation inhibitory activities in RAW 264.7 cells. *Cancer Lett*, **125**, 199-207.

Kim HW, Murakami A, Nakamura Y, Ohigashi H (2002). Screening of edible Japanese plants for suppressing effects on phorbol ester-induced superoxide generation in differentiated HL-60 cells and AS52 cells. *Cancer Lett*, **176**, 7-16.

Kobayashi T, Tsunawaki S, Seguchi H (2001). Evaluation of the process for superoxide production by NADPH oxidase in human neutrophils: evidence for cytoplasmic origin of superoxide. *Redox Rep*, **6**, 27-36.

Kobayashi Y, Nakae D, Akai H (1998). Prevention by 1'-acetoxychavicol acetate of the induction but not growth of putative preneoplastic, glutathione S-transferase placental form-positive, focal lesions in the livers of rats fed a choline-deficient, L-amino acid-defined diet. *Carcinogenesis*, **19**, 1809-14.

Koon-Sin N, Geoffery DB (1998). Stilbenes, monoterpenes, diarylheptanoids, labdanes, and chalcones from *Alpinia katsumadai*. *Phytochemistry*, **47**, 1117-23.

Koshimizu K, Ohigashi H, Tokuda H, Kondo A, Yamaguchi K (1988). Screening of edible plants against possible anti-tumor promoting activity. *Cancer Lett*, **39**, 247-57.

Kuroda M, Mimaki Y, Nishiyama T, et al (2005). Hypoglycemic effects of turmeric (*Curcuma longa* L. Rhizomes) on genetically diabetic KK-Ay mice. *Biol Pharm Bull*, **28**, 937-9.

Lee SL, Wang WW, Fanburg BL (1998). Superoxide as an intermediate signal for serotonin-induced mitogenesis. *Free Radic Biol Med*, **24**, 855-8.

Lee SK, Hong CH, Huh SK, et al (2002). Suppressive effect of natural sesquiterpenoids on inducible cyclooxygenase (COX-2) and nitric oxide synthase (iNOS) activity in mouse macrophage cells. *J Environ Pathol Toxicol Oncol*, **21**, 141-8.

Lee SI, Hyun PM, Kim SH, et al (2005). Suppression of the onset and progression of collagen-induced arthritis by chebulagic acid screened from a natural product library. *Arthritis Rheum*,

- 52, 345-53.
- Li A, Yue G, Li Y, Pan X, Yang TK (2003). Total asymmetric synthesis of (7*S*,9*R*)-(+)-bisacumol. *Tetrahedron: Asymmetry*, **134**, 75-8.
- Maeda H, Akaike T (1998). Nitric oxide and oxygen radicals in infection, inflammation, and cancer. *Biochemistry (Mosc)*, **63**, 854-65.
- Matsuda H, Ninomiya K, Morikawa T, Yoshikawa M (1998). Inhibitory effect and action mechanism of sesquiterpenes from *Zedoariae Rhizoma* on D-galactosamine/lipopolysaccharide-induced liver injury. *Bioorg Med Chem Lett*, **8**, 339-44.
- Miyauchi M, Nishikawa A, Furukawa F, et al (2000). Inhibitory effects of 1'-acetoxychavicol acetate on *N*-nitrosobis(2-oxopropyl)-amine-induced initiation of cholangiocarcinogenesis in Syrian hamsters. *Jpn J Cancer Res*, **91**, 477-81.
- Moffat J, Hashimoto M, Kojima A, et al (2000). Apoptosis induced by 1'-acetoxychavicol acetate in Ehrlich ascites tumor cells is associated with modulation of polyamine metabolism and caspase-3 activation. *Carcinogenesis*, **21**, 2151-7.
- Moffat J, Kennedy DO, Kojima A, et al (2002). Involvement of protein tyrosine phosphorylation and reduction of cellular sulfhydryl groups in cell death induced by 1'-acetoxychavicol acetate in Ehrlich ascites tumor cells. *Chem Biol Interact*, **139**, 215-30.
- Morikawa T, Matsuda H, Ninomiya K, Yoshikawa M (2002). Medicinal foodstuffs. XXIX. Potent protective effects of sesquiterpenes and curcumin from *Zedoariae Rhizoma* on liver injury induced by D-galactosamine/lipopolysaccharide or tumor necrosis factor- $\alpha$ . *Biol Pharm Bull*, **25**, 627-31.
- Murakami A, Jiwajinda S, Koshimizu K, Ohigashi H (1995). Screening for in vitro anti-tumor promoting activities of edible plants from Thailand. *Cancer Lett*, **95**, 139-46.
- Murakami A, Ohura S, Nakamura Y, Koshimizu K, Ohigashi H (1996). 1'-Acetoxychavicol acetate, a superoxide anion generation inhibitor, potently inhibits tumor promotion by 12-*O*-tetradecanoylphorbol-13-acetate in ICR mouse skin. *Oncology*, **53**, 386-91.
- Murakami A, Morita H, Safitri R, et al (1998). Screening for in vitro anti-tumor-promoting activities of edible plants from Indonesia. *Cancer Detec Prev*, **22**, 516-25.
- Murakami A, Gao G, O.-K. Kim, et al (1999). Identification of coumarins from the fruit of *Citrus hystrix* DC as inhibitors of nitric oxide generation in mouse macrophage RAW 264.7 cells. *J Agric Food Chem*, **47**, 333-9.
- Murakami A, Manaf A, Kamaruddin MS, Koshimizu K, Ohigashi H (2000a). Screening for the in vitro anti-tumor-promoting activities of edible plants from Malaysia. *Biosci Biotechnol Biochem*, **64**, 9-16.
- Murakami A, Gao G, Omura M, et al (2000b). 1,1-Dimethylallylcoumarins potently suppress both lipopolysaccharide- and interferon- $\gamma$ -induced nitric oxide generation in mouse macrophage RAW 264.7 cells. *Bioorg Med Chem Lett*, **10**, 59-62.
- Murakami A, Takahashi D, Kinoshita T, et al (2002). Zerumbone, a Southeast Asian ginger sesquiterpene, markedly suppresses free radical generation, proinflammatory protein production, and cancer cell proliferation accompanied by apoptosis: the  $\alpha,\beta$ -unsaturated carbonyl group is a prerequisite. *Carcinogenesis*, **23**, 795-802.
- Murakami A, Hayashi R, Takana T, et al (2003). Suppression of dextran sodium sulfate-induced colitis in mice by zerumbone, a subtropical ginger sesquiterpene, and nimesulide separately and in combination. *Biochem Pharmacol*, **66**, 1253-61.
- Murakami A, Tanaka T, Lee JY (2004). Zerumbone, a sesquiterpene in subtropical ginger, suppresses skin tumor initiation and promotion stages in ICR mice. *Int J Cancer*, **110**, 481-90.
- Nakamura Y, Gindhart TD, Winterstein D, et al (1988). Early superoxide dismutase-sensitive event promotes neoplastic transformation in mouse epidermal JB6 cells. *Carcinogenesis*, **9**, 203-7.
- Nakamura Y, Murakami A, Ohto Y, et al (1998). Suppression of tumor promoter-induced oxidative stress and inflammatory responses in mouse skin by a superoxide generation inhibitor 1'-acetoxychavicol acetate. *Cancer Res*, **58**, 4832-9.
- Nakamura Y, Kawamoto N, Ohto Y (1999). A diacetylenic spiroketal enol ether epoxide, AL-1, from *Artemisia lactiflora* inhibits 12-*O*-tetradecanoyl-phorbol-13-acetate-induced tumor promotion possibly by suppression of oxidative stress. *Cancer Lett*, **140**, 37-45.
- Nakamura Y, Murakami A, Ohigashi H (2000). Search for naturally-occurring antioxidative chemopreventers on the basis of the involvement of leukocyte-derived reactive oxygen species in carcinogenesis. *Asia Pacific J Cancer Prev*, **1**, 115-120.
- Nieto N, Torres MI, Fernandez MI, et al (2000). Experimental ulcerative colitis impairs antioxidant defense system in rat intestine. *Dig Dis Sci*, **45**, 1820-7.
- Nishiyama T, Mae T, Kishida H, et al (2005). Curcuminoids and sesquiterpenoids in turmeric (*Curcuma longa* L.) suppress an increase in blood glucose level in type 2 diabetic KK-Ay mice. *J Agric Food Chem*, **53**, 959-63.
- Ohata T, Fukuda K, Murakami A (1998). Inhibition by 1'-acetoxychavicol acetate of lipopolysaccharide- and interferon- $\gamma$ -induced nitric oxide production in RAW264 cells. *Carcinogenesis*, **19**, 1007-12.
- Purohit S, Kuma GP, Laloraya M, Laloraya MM (1994). Involvement of superoxide radical in signal transduction regulating stomatal movements. *Biochem Biophys Res Commun*, **205**, 30-7.
- Sakai H, Takeguchi N (1994). A GTP-binding protein inhibits a gastric housekeeping chloride channel via intracellular production of superoxide. *J Biol Chem*, **269**, 23426-30.
- Schwarz M, Peres G, Kunz W, et al (1984). On the role of superoxide anion radicals in skin tumour promotion. *Carcinogenesis*, **5**, 1663-70.
- Shacter E, Beecham EJ, Covey JM, Kohn KW, Potter M (1988). Activated neutrophils induce prolonged DNA damage in neighboring cells. *Carcinogenesis*, **9**, 2297-304.
- Shen Z, Wu W, Hazen SL (2000). Activated leukocytes oxidatively damage DNA, RNA, and the nucleotide pool through halide-dependent formation of hydroxyl radical. *Biochemistry*, **39**, 5474-82.
- Takada Y, Murakami A, Aggarwal BB (2005). Zerumbone abolishes NF- $\kappa$ B and I $\kappa$ B kinase activation leading to suppression of antiapoptotic and metastatic gene expression, upregulation of apoptosis, and downregulation of Invasion. *Oncogene*, **24**, 6957-69.
- Tanaka T, Makita H, Kawamori T, et al (1997a). A xanthine oxidase inhibitor 1'-acetoxychavicol acetate inhibits azoxymethane-induced colonic aberrant crypt foci in rats. *Carcinogenesis*, **18**, 1113-8.
- Tanaka T, Kawabata K, Kakumoto M, et al (1997b). Chemoprevention of azoxymethane-induced rat colon carcinogenesis by a xanthine oxidase inhibitor, 1'-acetoxychavicol acetate. *Jpn J Cancer Res*, **88**, 821-30.
- Tanaka T, Shimizu M, Kohno H, et al (2001). Chemoprevention

- of azoxymethane-induced rat aberrant crypt foci by dietary zerumbone isolated from *Zingiber Zerumbet*. *Life Sci*, **69**, 1935-45.
- Tardieu D, Jaeg JP, Deloly A, et al (2000). The COX-2 inhibitor nimesulide suppresses superoxide and 8-hydroxydeoxyguanosine formation, and stimulates apoptosis in mucosa during early colonic inflammation in rats. *Carcinogenesis*, **21**, 973-6.
- Trush MA, Seed JL, Kensler TW (1985). Oxidant-dependent metabolic activation of polycyclic aromatic hydrocarbons by phorbol ester-stimulated human polymorphonuclear leukocytes: possible link between inflammation and cancer. *Proc Natl Acad Sci USA*, **82**, 5194-8.
- Yamada T, Grisham MB (1991). Role of neutrophil-derived oxidants in the pathogenesis of intestinal inflammation. *Klin Wochenschr*, **69**, 988-94.
- Weitzman SA, Weitberg AB, Clark EP, Stossel TP (1985). Phagocytes as carcinogens: malignant transformation produced by human neutrophils. *Science*, **227**, 1231-3.
- Yang FQ, Li SP, Chen Y, et al (2005). Identification and quantitation of eleven sesquiterpenes in three species of *Curcuma* rhizomes by pressurized liquid extraction and gas chromatography-mass spectrometry. *J Pharm Biomed Anal*, **39**, 552-8.
- Weitzman SA, Gordon LI (1990). Inflammation and cancer: role of phagocyte-generated oxidants in carcinogenesis. *Blood*, **76**, 655-63.