

## RESEARCH COMMUNICATION

# Stage at Diagnosis and Relative Differences in Breast and Prostate Cancer Incidence in India: Comparison with the United States

James R Hebert<sup>1,2\*</sup>, Santosh S Ghumare<sup>1,2,3</sup>, Prakash C Gupta<sup>4</sup>

### Abstract

**Purpose:** To examine and reconcile differences in incidence rates and stage-at-initial-presentation of prostate and breast cancers in India, a country in epidemiologic transition. **Methods:** Age-adjusted prostate and female breast cancer incidence rates and proportion of cases by stage-at-diagnosis were compared. Data were derived from the National Cancer Registry Program of India, other Indian registries, the International Agency for Research on Cancer, and the US/ NCI Surveillance, Epidemiology, and End Results (SEER) Program. **Results:** Average annual cancer incidence rates in India ranged from 5.0 to 9.1 per 100,000/year for prostate and 7.2 to 31.3 per 100,000/year for female breast. Comparative rates in the US for prostate cancer are 110.4 for Whites and 180.9 for Blacks; for female breast, the rates are 86.6 for Blacks and 96.4 for Whites. Notable differences were observed between rural and urban areas in India, while such differences by rurality appear to be much smaller in the US. Overall, about 50-55% of breast cancer cases and about 85% of prostate cancers were detected at late (III and IV) stage; in contrast to the US where 15% of either cancer is diagnosed at late stage. **Conclusions:** Differences in stage-at-diagnosis help explain variations in incidence rates among cancer registries in India and rate differences between India and the US. These findings indicate that erroneous inferences will result from incidence-rate comparisons that do not take into account stage-at-diagnosis. Results also point to epidemiologic studies that could be conducted to deepen understanding of the etiology of these cancers. By enhancing data on staging, the Indian cancer registries could widen the scope of collaborative, cross-national research.

**Key Words:** Prostate cancer - breast cancer - epidemiology - incidence - India - stage at diagnosis - screening

*Asian Pacific J Cancer Prev*, 7, 547-555

### Introduction

India, a vast nation with a human population of over one billion, is the largest country on the Indian subcontinent in South Asia and home to about 16% of the world's population (Desai, 2002). India also is home to a variety of unique linguistic, religious, and culinary traditions. Some of these, most notably those related to diet, have been associated with cancer-related outcomes (dos Santos Silva et al., 2004; Gupta et al., 1999; Gupta et al., 1998; Hebert et al., 2002; Krishnaswamy, 1996; Messer, 1997; Sengupta et al., 2000; 2004b; Sinha, Anderson, McDonald, & Greenwald, 2003). Indeed, there is a vast literature on the role of specific components of Indian diet on cancer in laboratory models of carcinogenesis (Manju & Nalini, 2005; Sengupta et al., 2004a). With occasional additions of foods from other

cultures, most notably during the 16th and 17th centuries, the essential elements of the Indian traditional diet has been a distinct feature of Indian populations for centuries. These diets, which vary regionally and by socio-religious group, have been characterized by being primarily, and exclusively in many instances, vegetarian (Hebert et al., 1998; 1999; 2000). As part of religious 'prescriptions' diet also is part of larger lifestyle patterns that include yoga, massage, sleep hygiene, and rules of ethical behavior (Aruna & Sivaramakrishnan, 1992; Gogtay et al., 2002; Malhotra, 1967; Popkin, 2002; Popkin, Horton, Kim, & Mahal, 2001; Shetty, 2002; Sinha et al., 2003; Storer, 1977). This general pattern of eating (and associated behaviors) is consistent with scientific evidence indicating an association with lower rates of chronic diseases, such as cancer, and is hypothesized to be the explanation for relatively low occurrence of these

<sup>1</sup>South Carolina Statewide Cancer Prevention and Control Program, University of South Carolina, 2221 Devine Street Room 244A Columbia, SC 29208, USA; <sup>2</sup>Department of Epidemiology and Biostatistics, Arnold School of Public Health, University of South Carolina, Columbia, SC, USA, <sup>3</sup>Current Address: National Cancer Institute, Bethesda, MD, USA, <sup>4</sup>Healis-Sekhsaria Institute of Public Health, Navi Mumbai, India \*For Correspondence: Fax: 1-803-734-5259 Email: jhebert@sc.edu

diseases in India (Aruna & Sivaramakrishnan, 1992; Popkin et al., 2001; Sinha et al., 2003). However limitations in data quality and comparability make it difficult to draw inferences with confidence.

Though traditionally agrarian and rural, with rapid industrialization and urbanization, India is changing from a developing to a developed country. Changes in diet, lifestyle, and physical activity patterns, which constitute the basis of this demographic and epidemiologic transition (Popkin, 2002; Popkin et al., 2001), are being observed along with what appears to be concurrent rapid increase in the rates of chronic diseases such as cancer (Shetty, 2002). Presently rates of most cancers in India are lower than those in more developed countries such as the United States (Sinha et al., 2003). However, it is projected that the gap will be narrowing over the next decade or so (Pal & Mittal, 2004). As we show it appears to have begun to do so.

Cancer is responsible for about 20% of all deaths in industrialized countries and 10% of deaths in developing nations (Jones et al., 2006; "Stat bite: Estimated worldwide cancer mortality among men, 2002"; 2005). However the cancer burden in developing countries, including India, is expected to increase (Boutayeb & Boutayeb, 2005). Information about the frequency and patterns of cancer is an essential prerequisite for understanding the epidemiology of specific cancers. As we have discussed, descriptive epidemiology based on excellent cancer registry data can reveal important discrepancies in evidence from analytic epidemiologic studies that could shed light on the underlying causes of cancer (Hebert, 2005; Hebert et al., 2006b). After all, competent population-based cancer registries do not suffer from selection biases and resultant problems with inference that typically plague epidemiologic studies (Greenland & Robins, 1986; Rothman, 1988). Such information also is crucial for planning cancer control programs. In addition, India is an important country because of its size, strategic importance, and as a model for development as it undergoes a historic demographic and epidemiologic transition (Gopalan, 1999; "Health and Development Initiative India. World Health Day 2002" 2002; Nandakumar et al., 2004).

This paper reviews the recent trends in breast and prostate cancer incidence observed in various regions in India. We chose to focus on these two cancer sites in the context of the demographic and epidemiologic transitions because of their proven historical importance in countries that have undergone these transitions (Haynes, 1986; Kodama et al., 1992; Zheng et al., 2005), their public health relevance globally (Brudnak & Hoener, 2003; Ferlay et al., 2001; Food, Nutrition and the Prevention of Cancer: A Global Perspective, 1997; Ries et al., 2005; US Cancer Statistics Working Group, 2005), and their putative importance for assessing future trends in cancer risk (Parkin et al., 2001). We present an overview of the trends in these two cancer sites, by geographical region, and registry location (rural versus urban). In this, we emphasize differences observed in stage at initial presentation. Finally the possible role of



**Figure 1. Locations of Cancer Registries in India (Courtesy of Dr A Nandakumar)**

screening in order to identify a larger number of cases at an earlier disease stage is explored.

## Materials and Methods

### Indian Cancer Registries:

Cancer registration and data abstraction in India is performed mainly by the cancer registries situated in various regions of India (Figure 1). Many of these cancer registries were established by the Indian Council for Medical Research in 1982 under the National Cancer Registry Programme (Nandakumar et al., 2004). In addition to these, there are some additional registries which are not under the National Cancer Registry Programme, but collect and provide important cancer data. Because they are linked to real, geographically defined, population bases, the population-based cancer registries are able to compute cancer incidence rates. Therefore, we used reports published by all population-based cancer registries in order to obtain age-adjusted incidence rates. On the other hand, the hospital-based registries are constrained to provide just relative incidence. However, they provide valuable information on other aspects such as stage at diagnosis, which we incorporated in this study. Data on stage at initial presentation were obtained from hospital-based cancer registries and from previous publications, as cited.

### Method of Reporting and Calculation of Incidence Rates:

Usually cancer incidence rates are reported as the number of newly incident cases arising in the covered population in a given period of time, typically one calendar year. The convention, used here, is to express the rate as number of

incident cases per 100,000 population per year. Often the base population is a geopolitical unit (e.g., a state, one or more districts, or well-defined metropolitan area) and the rate is age-adjusted to a standard population for purposes of comparison. Usually when data are available for more than one calendar year, the average annual rates are reported (Nandakumar et al., 2004).

Because cancer is not a notifiable disease in India, an active method of data collection is used. The registrars receive advance training and administer the workers from the registries who are involved in processes such as scanning hospital records in the base population area, clarifying incomplete or contradictory information, and abstracting data from several different potential data sources (Nandakumar et al., 2004). The data thus obtained through the Indian Cancer Registries, as well as through the published studies conducted in Indian populations, were summarized and compared to discern if there were differences in stage at initial presentation.

*Other Sources of Incidence Rate Statistics:*

We used International Agency for Research on Cancer (IARC)'s publication, 'Cancer Incidence in Five Continents-Vol. VIII', and USA Surveillance, Epidemiology, and End Results (SEER) program statistics as other sources of incidence data. These were used to make comparisons between India and the US. Incidence rates in all populations described in this study were age-adjusted to the world standard population 2001 (Nandakumar et al., 2004).

*Data on Stage at Diagnosis:*

Data on stage at diagnosis in India were obtained mainly through hospital-based registries and published studies. These sources use the 'clinical extent of disease' system and classify the information on stage into three categories- 'localized', 'regional' and 'distant'. This differs from the Tumor Node Metastasis (TNM) system, which classifies the cancers into stage I through IV. Hence, we had to make certain assumptions and use references in the literature to present the stage information in TNM style.

*Years Covered:*

The results presented in this study are based mainly on the statistics for the period 1993-97, and 1997-98, and are specified in the results section. For comparability, we used the IARC and SEER statistics for approximately the same time period.

*Methods of Comparison:*

Because the focus is on a descriptive comparison of rates across populations, no formal statistical methods of hypothesis testing are presented.

**Results**


Based on data from the India population-based registries, we observed a steady increase in age-adjusted incidence rates

**Table 1. Average Annual Age-adjusted Incidence Rates for all Sites per 100,000 in Indian Population-based Cancer Registries (1997)**

Registry	Setting	Years Covered	Males	Females
Bangalore*	Urban	1990-1996	97.8	122.1
Barshi*	Rural	1990-1996	46.2	57.7
Bhopal*	Urban	1990-1996	100.4	92.2
Chennai*	Metropolitan	1990-1996	104.6	115.3
Delhi*	Metropolitan	1990-1996	121.9	135.3
Mumbai*	Metropolitan	1990-1996	115.4	119.1
Ahmedabad	Urban	1993-1997	107.2	82.9
Karunagapalli	Semi-urban	1993-1997	102.6	76.0
Kolkata	Metropolitan	1998-1999	102.1	114.6
Nagpur	Urban	1993-1997	118.4	118.8
Pune	Urban	1993-1997	103.9	115.3
Thiruvananthapuram	Urban	1993-1997	87.8	81.1

\* NCRP registries Source: 1. National Cancer Registry Programme-First All India Report 2001-2002. 2. Individual reports from non-NCRP registries

of most cancers over the years covered. The all-sites age-standardized (to world standard population) cancer incidence rates in urban NCRP cancer registries in India for the period 1990-96 ranged from 97.8 to 121.9 per 100,000 for men and from 92.2 to 135.7 per 100,000 for women (Table 1). The Delhi registry recorded the highest incidence for both men and women, whereas the rates from the rural population-based registry in Barshi (in the Western Indian state of Maharashtra) were the lowest, at 46.2 and 57.7 per 100,000 for men and women, respectively (Nandakumar et al., 2004). In comparison, the corresponding rates for the US range from 291.4 to 538.6 per 100,000 for men and from 175.3 to 311.5 per 100,000 for women (SEER all sites rates for 1992-1997 adjusted to world standard population). Since 1985, the cancer registries in India have reported a 12% increase in cancer cases (Pal & Mittal, 2004), which is much higher than the rate of increase in the US. The Indian rates for tobacco-related cancer sites such as oral and esophageal cancer are among the highest in the world, whereas for other sites, such as prostate, they are among the lowest recorded



**Figure 2. Breast and Prostate Cancer Incidence Rates in India (1997).** Source: National Cancer Registry Programme-First All India report 2001-2002

**Table 2. Combined Breast Cancer and Prostate Cancer Incidence and Mortality Rates in Indian Population-based Cancer Registries (1990-1998)**

Site	Incidence	Mortality	Prevalence	
	Age standardized rates		1-year	5-year
Prostate	4.6	3.0	12,988	42,521
Breast	19.1	10.4	71,493	269,470

Source: National Cancer Registry Programme-First All India Report 2001-2002

(Jones et al., 2006; Sinha et al, 2003; “Stat bite: Estimated worldwide cancer mortality among men, 2002”, 2005).

The overall age-standardized incidence rate from all registries was 4.6 per 100,000 for prostate cancer and 19.1 per 100,000 for breast cancer. Again, the range in incidence rates of both these sites (1.5 to 7.1 per 100,000 for prostate, 8.8 to 28.6 per 100,000 for breast) varied by registry regions and a clear difference was observed between the rural registry area (Barshi) and other registries (located in or near urban areas) (Figure 2). Mortality and prevalence rates of the two cancer sites differed noticeably in magnitude as well (Table 2).

A ten-year report (1983-1993) of the hospital-based cancer registries suggests that at the time of diagnosis, the percentage of patients with localized disease for all cancer sites combined was only 24.3 % in men and 22.8 % in women, about the inverse of the situation in the United States in the corresponding period (Parker, Tong, Bolden, & Wingo, 1997) and indicating an overall late detection phenomenon in India.


*Trends in Prostate Cancer Incidence:*

Prostate cancer is a much more commonly diagnosed disease in the West than in India; it is the most prevalent cancer in American men and those from other Western countries (Drake et al., 2006; Hebert et al., 1998). For example, the age-adjusted incidence in US Black population is more than 18 times higher than that observed in the population in Mumbai, India (Hsing et al., 2000). However

**Table 3. Age-adjusted Incidence Rates of Prostate and Breast Cancer per 100,000 in Indian Population-based Cancer Registries (1997)**

Registry	Female Breast	Prostate
Bangalore*	22.1	4.3
Barshi*	8.8	1.5
Bhopal*	19.9	5.1
Chennai*	21.6	3.8
Delhi*	28.1	6.5
Mumbai*	28.6	7.1
Ahmedabad	19.1	3.6
Karunagapalli	15.0	-
Kolkata	25.1	5.3
Nagpur	24.2	-
Pune	26.9	6.6
Thiruvananthapuram	19.7	4.0

\* NCRP Registries Source: National Cancer Registry Programme-First All India Report 2001-2002



**Figure 3. Breast and Prostate Cancer Incidence: Time Trends in Two Urban Registries.** Source: National Cancer Registry Program-First All India report 2001-2002

over the last two decades the incidence of prostate cancer in India has been increasing across most registry regions (Banerjee et al., 2003; Ferlay et al., 2004; Nandakumar et al., 2004), as illustrated by the example of Chennai and Mumbai registries (Figure 3). Large differences exist in the age-standardized prostate cancer rates in India. The rural population-based registry at Barshi, in Western Maharashtra, recorded the lowest age-standardized incidence of 1.5 per 100,000; whereas the Mumbai registry had a higher incidence rate of 7.1 per 100,000 (Table 3). A study reported the highest incidence of prostate cancer in the country (11.6 per 100,000) at Jaipur in Rajasthan state (Sharma et al., 1994). The world age-standardized US rates for the same period as reported by SEER registries were 110.4 for Whites and 180.9 for Blacks.

*Prostate Cancer Stage at Diagnosis:*

As noted, prostate cancer is generally not diagnosed in early stages in India. The proportion of cases classified as ‘regional’ (which partially represents stage III) and ‘distant’ (Stage IV) illustrate this fact (Table 4). A study conducted

**Table 4. Indian Hospital-Based Registries: Comparison of Stage at Initial Presentation**

Registry (years)	Cancer site	Stage distribution (%)			
		Loc	Reg	Dist	Other
Chandigarh (84-89)	Prostate	18.4	58.2	23.4	0.0
	Breast	14.3	71.5	14.3	0.0
Mumbai (84-93)	Prostate	14.1	22.1	58.9	4.9
	Breast	31.9	51.9	12.1	4.1
Bangalore (84-93)	Prostate	20.0	20.0	54.7	5.3
	Breast	16.0	62.0	20.7	1.3
Chennai (84-93)	Prostate	5.6	45.1	49.3	0.0
	Breast	2.6	78.6	18.7	0.1
Thiruvananthapuram (84-93)	Prostate	41.1	41.1	16.4	1.4
	Breast	18.2	71.2	10.0	0.4

Source: National Cancer Registry Programme-Hospital-based registries report

at the Bombay Hospital Institute of Medical Sciences suggests that the proportion of prostate cancer cases diagnosed at an advanced stage may be as high as 84% at initial presentation (Srinivas et al., 1995). This is in stark contrast to the situation in the US, where the converse is true; i.e., only about 15% of patients are diagnosed at an advanced stage (Parker et al., 1997).

*Trends in Breast Cancer Incidence:*


As for prostate cancer, age-standardized breast cancer incidence in India is significantly lower than in the United States and Europe (Ferlay et al., 2004; Raina et al., 2005). However, most registries indicate an increasing trend over the years. The data from the Chennai and Mumbai registries demonstrates this trend (Figure 3). Breast cancer is now replacing cervical cancer as the leading cancer site in most registries, especially in urban areas (Nandakumar et al., 2004), although the trend can be seen even in predominantly rural areas such as Eastern Rajasthan (Sharma et al., 1994). It is emerging as the leading cause of cancer mortality in Indian women. Nearly 80,000 new cases of breast cancer are diagnosed annually in India (Sinha et al., 2003). This number is projected to surpass 100,000 by the year 2010 (Saxena et al., 2002).

Age-standardized female breast cancer rates in India evince about four-fold range in the population-based cancer registries. Again, a notable difference is observed between the rural and urban areas (Table 3). The rural registry at Barshi registered the lowest incidence rate of 8.8 per 100,000; the semi urban registry at Karunagapalli in Southern India had an intermediate rate of 15.3 while the registries in urban regions such as Delhi and Mumbai had higher incidence rates of around 28 per 100,000 (Nandakumar et al., 2004). The Kolkata Registry in the Eastern region showed an incidence rate of 25.1 (Chopra, 2001; Sen & Sankaranarayanan, 2002). By comparison, the US rates were 86.6 and 96.4 per 100,000 for Blacks and Whites, respectively.

Reflecting an interesting trend, age-specific female breast cancer incidence rates from most of the urban registries show a steep increase until about the time of menopause (age 49 years), and then the rates plateau. This is very dissimilar to the general pattern observed in Western women, in whom breast cancer rates increase sharply after menopause. Recent trends, especially among urban Indian women, indicate that the increase in incidence is higher post-menopausally (Nandakumar et al., 2004; Yeole & Kurkure, 2003). This observation also alludes to the other possible factors responsible for a relatively greater incidence of breast cancer in India (compared to prostate cancer), beyond the difference explained by late stage at diagnosis.

*Breast Cancer Stage at Diagnosis:*

Breast cancer constitutes nearly one fifth of all female cancers in India, and many patients present with advanced disease. On average, 57% of breast cancer cases in India present at late stage (stage III and IV) (Chopra, 2001), and



**Figure 4. Proportion (%) of Cancer Cases by Stage at Initial Presentation in India.** Sources: 1. National Cancer Registry Program-First All India report 2001-2002 2. Srinivas et al., 1995

in some areas as many as 70% of breast cancer cases are locally advanced when diagnosed (Table 4) (Singletary & Connolly, 2006). Although much higher than the 12% figure observed in the US (Goel et al., 1995), this is still much lower than the 85% observed for prostate cancer.

*Comparing Breast and Prostate Cancers*

For both of these cancers, the age-adjusted incidence rates in India were much lower than those reported by SEER in the US for approximately the same period (1992-1997). The difference was much more striking for prostate (up to 18-fold) than breast (3- to 4-fold) (Ries et al., 2005). However, it should be kept in mind that in the US, both of these cancers are subject to extensive population-based and diagnostic screening as compared to India.

For both breast and prostate cancers noticeable differences could be observed by rurality (i.e., rural versus urban), hospital-based versus population-based registry, and by geographic region. For example, data from the population-based registry in Mumbai indicate that 55% of incident prostate cancer cases were in men > 70 years of age, whereas for the Mumbai hospital-based registry, the proportion in that age group was 33%. The proportion of late-stage prostate cancer cases was very different as well; i.e., 62% for population-based registries and 32% for hospital-based registries. In general though, the proportion of late-stage breast cancer cases was much lower than the proportion of late-stage prostate cancer cases (Figure 4).

**Discussion**

Reports from Indian cancer registries and epidemiologic studies on cancer incidence in India reveal intriguing trends in both breast and prostate cancers. Across regions, either within India or comparing India to the US, there appears to be a generally inverse relationship between recorded incidence of each cancer and virulence (as measured by stage). This suggests that potentially treatable, early-stage disease is going undetected until very late in the natural

history of the cancer.

These findings also point to interesting areas on which to focus future research. Potential studies range from those focused on cancer etiology to ones focusing on provision of health services. Based primarily on ecological and laboratory evidence, the traditional Indian diet, though varying regionally and across socio-religious groups, would be expected to be associated with low rates of non-tobacco-related epithelial cancers (Branca & Lorenzetti, 2005; Food, Nutrition and the Prevention of Cancer: A Global Perspective, 1997; Hebert, 2004; Hebert et al., 1998; Hebert & Rosen, 1996; Sinha et al., 2003). India is in demographic transition, a process which began over 50 years ago and has been uneven across the population (Dasgupta, 1995; Purohit, 2004) evincing a pattern typical of much of the current developing world (Bardhan, 2006; Wade, 2004). Recent changes in the economic status of many Indians, including a burgeoning middle class and a general pattern of urbanization (Misra et al., 2001; Purohit, 2004; Tiwari, Kumar, & Kumar, 2005; Wade, 2004), has in fact accelerated the epidemiologic transition (Chopra, 2001; Pal & Mittal, 2004).

India provides an extraordinarily interesting venue for examining the effect of lifestyle, most notably diet, and changes in those factors on both cancer rates and changes in rates. This is because dietary behavior and adherence to dietary traditions are not entirely under the dominance of economic influences (as they had been in Western cultures); in India these behaviors are largely determined by socio-religious factors (Chadha, 1995; Sinha et al., 2003). Therefore, the kinds of potential for confounding of the diet-cancer relationship that might exist in other cultures probably would be substantially reduced in India. This creates interesting opportunities to investigate the role of diet and other risk factors, such as physical activity, that we would expect to co-vary in unusual and potentially illuminating ways.

The issue of variability of cancer rates in India, especially when contrasting the urban and rural registries, is fascinating. This urban-rural contrast, which appears to be proportionally as large as US-India difference in some instances, may hold the key to understanding the relationship between low incidence and late stage of these cancers. Additionally, it possibly throws light on the trends observed in transitioning nations such as India. The importance of the urban-rural contrast is illustrated by the fact that that in the US, SEER rates (which represent virtually no African Americans living in rural areas) are often different from those derived from the National Program of Cancer Registries data (which does represent the 55% of African Americans who live in the rural Southeast) (Hebert, 2003; Hebert, 2005; Hebert et al., 2006a; 2006b; Ries et al., 2005; US Cancer Statistics Working Group, 2005). For example, prostate cancer incidence rates among African Americans in the South Atlantic region are generally about 10-15% higher than they are nationally (US Cancer Statistics Working Group, 2005; Drake et al., 2006). This raises interesting questions about the possible

role of genetic factors related to admixture (Parra et al., 2001) that may relevant to the situation in India (and other parts of the world).

Using registry-derived data of uniformly very high quality we are able to comment on racial differences in cancer rates in very much smaller geopolitical units; e.g., the State of South Carolina (Hebert et al., 2006a; South Carolina Cancer Alliance, 2005). The results from our exploration are intriguing and point to the need for both improved and expanded data collection methods and research to understand the causes for the observed trends, both real and artificial. Indeed, it is important to note that the trends should be interpreted with caution for several reasons. The establishment of cancer registries and collection of cancer data in India is a relatively recent phenomenon and the facilities are still in the developing stages. Additionally, the existing registries cover limited geographical regions. Consequently, large areas of the population, particularly the rural areas, are still not covered sufficiently and the patterns of cancer in these areas remain largely unknown.

This work underlines the need for improvements in standardizing collection techniques across areas and understanding, within the constraints of the data, how differences in access to care, knowledge, attitudes, and beliefs can lead to differences in the overall detection rate of disease. It also may point to an explanation for the observed differences in incidence and virulence between what are generally more occult prostate cancers and more palpable breast cancers.

Clearly, the use of registry data in the United States; where we are able to categorize according to histologic and anatomic subtype, stage, and grade of disease, has allowed us to make significant progress in understanding racial differences in the descriptive epidemiologies of several cancers, most notably squamous cell carcinomas of the esophagus (Hebert et al., 2006b). Data on staging reported by most cancer registries in India are not presented according to the staging systems more commonly used clinically, such as the American Joint Committee on Cancer (AJCC)/UICC Tumor Node Metastases (TNM) stage grouping. Instead 'clinical extent of disease' classified as 'localized', 'regional', and distant is reported. References suggest that these represent 'stage 0 and I', 'stage II and III', and 'stage IV' of the TNM system, respectively (Reynolds, 2002; Singletary & Connolly, 2006). But even this information is often incomplete, and many cases are classified as 'stage unknown'. This leads to an approach of making reasonable assumptions and then drawing inferences. However this in turn limits direct comparisons with data from the West. We suggest that Indian cancer registries take appropriate steps to resolve this impediment to both epidemiologic research and healthcare planning.

We were able to access the data from the first All India Report on Cancer Incidence in India published by the National Cancer Registries Program, which covers data up to year 1997 as well as the updated data available from some

individual registries through their 2001 reports (Nandakumar et al., 2004). A more recent report based on data from the population-based registries for 1999-2000 has been published by the National Cancer Registries Program, which also has published information on cancer cases diagnosed in areas beyond those served by the registries (Nandakumar et al., 2005). However we chose to use the published data for reasons of accessibility by other interested parties.

In conclusion, unlike the observed similarities in both overall incidence and stage-at-diagnosis in the US, incidence and stage-at-presentation of breast cancer and prostate cancer in India differ notably from each other. The nearly three-fold difference in the proportion of prostate versus breast cancer cases detected in early stages (stage I and II) correlates with the three-fold difference observed in the age-standardized incidence rates of prostate and breast cancers. This suggests that the actual difference in the incidence rates of these two cancers in India may not be as high as the observed incidence rates would indicate. In fact, to a large extent, they may reflect differences in stage at initial presentation (and corollary issues such as seeking care because of signs and symptoms of disease). This, in turn, may be a concomitant of the paucity of programs aimed at early detection and very uneven distribution of wealth (Dasgupta, 1995; Purohit, 2004).

Both prostate and breast cancers can be detected in early stages by screening tests; e.g., prostate specific antigen (PSA) screening for prostate cancer and mammography for breast cancer, that are widely used in the West. Despite its shortcomings (Lehrer et al., 2002; Drake et al., 2006), PSA is the most commonly used and clinically useful tumor marker in the diagnosis of prostate cancer (Saw & Aw, 2000). The prostate cancer mortality rate has decreased slightly in the US over the past decade, and it is suggested that a primary reason may be increased use of the PSA test to detect prostate cancer early, especially in men with virulent disease (Chu et al., 2003). One concern, however is that very indolent diseases that are picked up by PSA screening are not interesting from an epidemiologic perspective (Hardie et al., 2005; Johansson et al., 2004; von Eschenbach, 1996). Another is that detection of these cancers also creates economic and ethical dilemmas (Klotz, 2005; Lefevre, 1998), especially regarding treatment.

While large-scale, population-based screening for prostate cancer and breast cancer in India is infeasible, instituting small-scale demonstration projects would serve two purposes. First, it would allow us to test the effect of down staging on incidence. Second, if conducted on a large scale, it would provide a 'platform' on which observational epidemiologic studies could be conducted for research into etiologic factors that may be operative in a country in the process of an epidemiologic transition. Well-conceived and designed research would deepen our understanding of the underlying causes of differences in observed rates of these (and other) cancers across distinct areas (e.g., urban vs. rural). Third, it would allow an estimation of how many breast and prostate cancers might be prevented entirely.

## Acknowledgements

We are grateful to Drs. Balkrishna Yeole (Indian Cancer Society), Ganesh Balasubramaniam, and A. Nandakumar (NCRP) for providing valuable input from India on recent cancer statistics. We are also very appreciative of Dr. R. Sankaranarayanan (Head, Screening Group, International Agency for Research on Cancer, Lyon, France), for links to data sources and with periodic comments on the project. Special thanks also to Dr Ivo Foppa (Department of Epidemiology and Biostatistics, Arnold School of Health, University of South Carolina) for his comments on the draft.

## References

- Aruna K, Sivaramakrishnan VM (1992). Anticarcinogenic effects of some Indian plant products. *Fd Chem Toxicol*, **30**, 953-6.
- Banerjee AG, Liu J, Yuan Y, et al. (2003). Expression of biomarkers modulating prostate cancer angiogenesis: differential expression of annexin II in prostate carcinomas from India and USA. *Mol Cancer*, **2**, 34.
- Bardhan P (2006). Does globalization help or hurt the world's poor? *Scientific American*, **294**, 84-91.
- Boutayeb A, Boutayeb S (2005). The burden of non communicable diseases in developing countries. *Int J Equity Health*, **4**, 2.
- Branca F, Lorenzetti S (2005). Health effects of phytoestrogens. *Forum Nutr*, **57**, 100-111.
- Brudnak MA, Hoener SG (2003). Plato, Sun Tsu, and the Art of War (on cancer). Can we learn from history? *Medical Hypotheses*, **60**, 603-6.
- Chadha SL (1995). Dietary profile of adults in an urban and a rural community. *Ind J Med Res*, **101**, 258-67.
- Chopra R (2001). The Indian scene. *J Clin Oncol*, **19(18 Suppl)**, 106S-111S.
- Chu KC, Tarone RE, Freeman HP (2003). Trends in prostate cancer mortality among black men and white men in the United States. *Cancer*, **97**, 1507-16.
- Dasgupta P (1995). *An Inquiry into Well-Being and Destitution*. Oxford: Oxford University Press.
- Desai PB (2002). Cancer control efforts in the Indian subcontinent. *Jpn J Clin Oncol*, **32 Suppl**, S13-16.
- dos Santos Silva I, Mangtani P, McCormack V, et al (2004). Phytoestrogen intake and breast cancer risk in South Asian women in England: findings from a population-based case-control study. *Cancer Causes Control*, **15**, 805-18.
- Drake BF, Keane TE, Mosley CM, et al (2006). Prostate cancer disparities in South Carolina: Early detection, special programs, and descriptive epidemiology. *J South Carolina Med Assoc*, **102**, 241-9.
- Ferlay J, Bray F, Pisani P, Parkin DM (2001). *Cancer Incidence, Mortality and Prevalence Worldwide*. On GLOBOCAN 2000: International Agency for Research on Cancer, World Health Organization.
- Ferlay J, Bray F, Pisani P, Parkin DM (2004). *Globocan 2002: Cancer Incidence, Mortality and Prevalence Worldwide*. IARC Cancer Base, 5 (2.0).
- Food, Nutrition and the Prevention of Cancer: A Global Perspective. (1997). Washington, DC: American Institute for Cancer Research.
- Goel AK, Seenu V, Shukla NK, Raina V (1995). Breast cancer presentation at a regional cancer centre. *Natl Med J India*, **8**, 6-9.

- Gogtay NJ, Bhatt HA, Dalvi SS, Kshirsagar NA (2002). The use and safety of non-allopathic Indian medicines. *Drug Saf*, **25**, 1005-19.
- Gopalan C (1999). Nutrition and Developmental Transition: Lessons from Asian Experience.
- Greenland S, Robins JM (1986). Identifiability, exchangeability, and epidemiological confounding. *Int J Epidemiol*, **15**, 413-9.
- Gupta PC, Hebert JR, Bhonsle RB, et al (1999). Influence of dietary factors on oral precancerous lesions in a population-based case-control study in Kerala, India. *Cancer*, **85**, 1885-93.
- Gupta PC, Hebert JR, Bhonsle RB, et al (1998). Dietary factors in oral leukoplakia and submucous fibrosis in a population-based case-control study in Gujarat, India. *Oral Dis*, **4**, 200-206.
- Hardie C, Parker C, Norman A, et al (2005). Early outcomes of active surveillance for localized prostate cancer. *BJU Int*, **95**, 956-90.
- Haynes R (1986). Cancer mortality and urbanization in China. *Int J Epidemiol*, **15**, 268-71.
- Health and Development Initiative India. World Health Day 2002. (2002).
- Hebert JR (2003). Invited commentary: menthol cigarettes and risk of lung cancer. *Am J Epidemiol*, **158**, 617-20.
- Hebert JR (2004). Epidemiologic studies of diet and cancer: The case for international collaboration (Commissioned as part of the Eminent Scientist of the Year 2004 Award by the World Science Forum). *Austro-Asian J Cancer*, **5** (Special Issue - Recent Advances and Research Updates), 140-53.
- Hebert JR (2005). Epidemiologic studies of diet and cancer: The case for international collaboration. *Austro-Asian J Cancer*, **4**, 125-34.
- Hebert JR, Elder K, Ureda JR (2006b). Meeting the challenges of cancer prevention and control in South Carolina: Focus on seven cancer sites, engaging partners. *J South Carolina Med Assoc*, **102**, 177-82.
- Hebert JR, Gupta PC, Bhonsle RB, et al (2002). Dietary exposures and oral precancerous lesions in Srikakulam District, Andhra Pradesh, India. *Public Health Nutr*, **5**, 303-12.
- Hebert JR, Gupta PC, Bhonsle RB, et al (1998). Development and testing of a quantitative food frequency questionnaire for use in Kerala, India. *Public Health Nutr*, **1**, 123-30.
- Hebert JR, Gupta PC, Bhonsle RB, et al (1999). Development and testing of a quantitative food frequency questionnaire for use in Gujarat, India. *Public Health Nutr*, **2**, 39-50.
- Hebert JR, Gupta PC, Mehta H, et al (2000). Sources of variability in dietary intake in two distinct regions of rural India: implications for nutrition study design and interpretation. *Eur J Clin Nutr*, **54**, 479-86.
- Hebert JR, Adams SA, Daguise VG, et al (2006b). Esophageal cancer disparities in South Carolina: Early detection, special programs, and descriptive epidemiology. *J South Carolina Med Assoc*, **102**, 201-9.
- Hebert JR, Hurley TG, Olendzki B, et al (1998). Nutritional and socioeconomic factors in relation to prostate cancer mortality: A cross-national study. *J Natl Cancer Inst*, **90**, 1637-47.
- Hebert JR, Rosen A (1996). Nutritional, socioeconomic, and reproductive factors in relation to female breast cancer mortality: findings from a cross-national study. *Cancer Detect Prevent*, **20**, 234-44.
- Hsing AW, Tsao L, Devesa SS (2000). International trends and patterns of prostate cancer incidence and mortality. *Int J Cancer*, **85**, 60-7.
- Johansson JE, Andren O, Andersson SO, et al (2004). Natural history of early, localized prostate cancer. *JAMA*, **291**, 2713-9.
- Jones LA, Chilton JA, Hajek RA, Iammarino NK, Laufman L (2006). Between and within: international perspectives on cancer and health disparities. *J Clin Oncol*, **24**, 2204-8.
- Klotz LH (2005). Active surveillance with selective delayed intervention: walking the line between overtreatment for indolent disease and undertreatment for aggressive disease. *Can J Urol*, **1**, 53-7.
- Kodama M, Kodama T, Murakami M, et al (1992). Interrelation between Western type cancers and non-Western type cancers as regards their risk variation in time and space. IV. Hormonal transition of Japanese women from the pro-cervical cancer age through the pro-endometrial cancer age to the pro-hypogonadism age. *Anticancer Res*, **12**, 693-704.
- Krishnaswamy K (1996). Indian functional foods: role in prevention of cancer. *Nutrition Reviews*, **54**, S127-131.
- Lefevre ML (1998). Prostate cancer screening: more harm than good? *Am Fam Phys*, **58**, 432-8.
- Lehrer S, Stone NN, Droller MJ, Stock RG (2002). Association between American Urologic Association (AUA) urinary symptom score and disease stage in men with localized prostate cancer. *Urol Oncol*, **7**, 73-6.
- Malhotra SL (1967). Geographical aspects of acute myocardial infarction in India with special reference to patterns of diet and eating. *Br Heart J*, **29**, 337-44.
- Manju V, Nalini N (2005). Chemopreventive efficacy of ginger, a naturally occurring anticarcinogen during the initiation, post-initiation stages of 1,2 dimethylhydrazine-induced colon cancer. *Clin Chim Acta*, **358**, 60-7.
- Messer E (1997). Intra-household allocation of food and health care: Current findings and understandings--introduction. *Soc Sci Med*, **44**, 1675-1684.
- Misra A, Sharma R, Pandey RM, Khanna N (2001). Adverse profile of dietary nutrients, anthropometry and lipids in urban slum dwellers of northern India. *Eur J Clin Nutr*, **55**, 727-34.
- Nandakumar A, Gupta PC, Gangadharan P, Visweswara RN (2004). Development of an Atlas of Cancer in India, First All India Report: 2001-2002. Bangalore, India: National Cancer Registry Programme (ICMR).
- Nandakumar A, Gupta PC, Gangadharan P, Visweswara RN (2004). First All India Report: 2001-2002: Mapping Patterns of Cancer. Bangalore: National Cancer Registry Programme (ICMR).
- Nandakumar A, Gupta PC, Gangadharan P, Visweswara RN, Parkin DM (2005). Geographic pathology revisited: development of an atlas of cancer in India. *Int J Cancer*, **116**, 740-54.
- Pal SK, Mittal B (2004). Improving cancer care in India: prospects and challenges. *Asian Pac J Cancer Prev*, **5**, 226-8.
- Parker SL, Tong T, Bolden S, Wingo PA (1997). Cancer statistics, 1997. *CA Cancer J Clin*, **47**, 5-27.
- Parkin DM, Bray FI, Devesa SS (2001). Cancer burden in the year 2000. The global picture. *Eur J Cancer*, **37 Suppl 8**, S4-66.
- Parra EJ, Kittles RA, Argyropoulos G, et al. (2001). Ancestral proportions and admixture dynamics in geographically defined African Americans living in South Carolina. *Am J Phys Anthropol*, **114**, 18-29.
- Popkin BM (2002). An overview on the nutrition transition and its health implications: the Bellagio meeting. *Public Health Nutr*, **5**, 93-103.
- Popkin BM, Horton S, Kim S, Mahal A (2001). Trends in diet, nutritional status, and diet-related noncommunicable diseases in China and India: the economic costs of the nutrition transition. *Nutr Rev*, **59**, 379-90.
- Purohit BC (2004). Inter-state disparities in health care and financial burden on the poor in India. *J Health Soc Policy*, **18**, 37-60.



- Raina V, Bhutani M, Bedi R, et al (2005). Clinical features and prognostic factors of early breast cancer at a major cancer center in North India. *Ind J Cancer*, **42**, 40-5.
- Reynolds T (2002). Updates to staging system reflect advances in imaging, understanding. *J Natl Cancer Inst*, **94**, 1664-6.
- Ries LAG, Eisner MP, Kosary CL, et al (2005). Cancer Statistics Review 1975-2002. Bethesda, MD: National Cancer Institute.
- Ries LAG, Eisner MP, Kosary CL, et al (2005). SEER Cancer Statistics Review, 1975-2002. Bethesda, MD: National Cancer Institute.
- Rothman K (Ed.). (1988). *Casual Inference*. Chestnut Hill, MA: Epidemiology Resources.
- Saw S, Aw TC (2000). Age-related reference intervals for free and total prostate-specific antigen in a Singaporean population. *Pathology*, **32**, 245-9.
- Saxena S, Szabo CI, Chopin S, et al (2002). BRCA1 and BRCA2 in Indian breast cancer patients. *Hum Mutat*, **20**, 473-4.
- Sen U, Sankaranarayanan R (2002). Cancer patterns in eastern India: the first report of the Kolkata Cancer Registry. *Int J Cancer*, **100**, 86-91.
- Sengupta A, Ghosh S, Bhattacharjee S, Das S (2004a). Allium vegetables in cancer prevention: An overview. *Asian Pac J Cancer Prev*, **5**, 237-45.
- Sengupta A, Ghosh S, Bhattacharjee S, Das S (2004b). Indian food ingredients and cancer prevention: An experimental evaluation of anticarcinogenic effects of garlic in rat colon. *Asian Pac J Cancer Prev*, **5**, 126-132.
- Sharma RG, Ajmera R, Saxena O (1994). Cancer profile in eastern Rajasthan. *Ind J Cancer*, **31**, 160-73.
- Shetty PS (2002). Nutrition transition in India. *Public Health Nutr*, **5**, 175-82.
- Singletary SE, Connolly JL (2006). Breast cancer staging: working with the sixth edition of the AJCC Cancer Staging Manual. *CA Cancer J Clin*, **56**, 37-47; quiz 50-31.
- Sinha R, Anderson DE, McDonald SS, Greenwald P (2003). Cancer risk and diet in India. *J Postgrad Med*, **49**, 222-8.
- South Carolina Cancer Alliance. (2005). South Carolina Cancer Report Card, 2004. Columbia, SC.
- Srinivas V, Mehta H, Amin A, et al (1995). Carcinoma of the prostate--state at initial presentation. *Int Urol Nephrol*, **27**, 419-22.
- Stat bite: Estimated worldwide cancer mortality among men, 2002. (2005). *J Natl Cancer Inst*, **97**, 1402.
- Storer J (1977). 'Hot' and 'cold' food beliefs in an Indian community and their significance. *J Human Nutr*, **31**, 33-40.
- Tiwari SC, Kumar A, Kumar A (2005). Development & standardization of a scale to measure socio-economic status in urban & rural communities in India. *Ind J Med Res*, **122**, 309-14.
- US Cancer Statistics Working Group. (2005). United States Cancer Statistics: 2002 Incidence and Mortality. Atlanta (GA): US DHHS/CDC/NIH-NCI.
- von Eschenbach AC (1996). The biologic dilemma of early carcinoma of the prostate. *Cancer*, **78**, 326-9.
- Wade RH (2004). Is globalization reducing poverty and inequality? *Int J Health Serv*, **34**, 381-414.
- Yeole BB, Kurkure AP (2003). An epidemiological assessment of increasing incidence and trends in breast cancer in Mumbai and other sites in India, during the last two decades. *Asian Pac J Cancer Prev*, **4**, 51-6.
- Zheng W, Chow WH, Fan J, et al (2005). The Shanghai Women's Health Study: rationale, study design, and baseline characteristics. *Am J Epidemiol*, **162**, 1123-31.