

RESEARCH COMMUNICATION

Do Turkish Nursing and Midwifery Students Teach Breast Self-Examination to Their Relatives?

Arife Özkan¹, Arzu Tuna Malak¹, Aysel Gürkan², Ayse San Turgay¹

Abstract

Aim: To describe health beliefs and breast self-examination (BSE) practice of Turkish female nursing and midwifery students and assess teaching of BSE to their mothers, sisters, relatives. **Methods:** The study was designed as a definition survey, with data obtained from 113 participants, in third and fourth class and their mothers, sisters, relatives in Turkey. Data were collected by using a personal data form, knowledge evaluation form for BSE (Maurer 1997) and the Champion's Health Belief Model Scale. **Results:** Students had learned about breast cancer and BSE in their lessons one or two years previous to the study. Knowledge level scores of the students were 52.3 ± 9.63 (min:25, max: 75). Rate of regular BSE was 32.7%. When health belief scale assessed, the average susceptibility was 7.52 ± 2.62 , seriousness was 21.8 ± 5.30 , benefit was 16.7 ± 4.45 , barrier was 22.3 ± 6.44 , confidence was 40.3 ± 6.67 and medical motivation was 26.6 ± 4.22 . A statistically significant difference in the rate of having regular BSE and benefit, barrier scores was noted ($p < 0.05$). Regarding BSE training, 91.3% (n = 106) gave assistance to their mother and sisters, 42.6% (n = 48) to relatives, 6.2% (n = 7) to friends, and 5.4% (n = 6) to patients. **Conclusions:** Knowledge about breast cancer and BSE repetition training programs should be planned for nursing/midwifery students. Their susceptibility, belief and attitudes, medical motivation with BSE should thereby increase.

Keywords: Breast self examination - female nursing/midwifery students - training - family and friends

Asian Pacific J Cancer Prev, 11, 1569-1573

Introduction

Breast cancer is the most common type of cancer among women across the world (Darendeliler & Ağaoğlu, 2003; Ozmen, 2008). In Turkey, the incidence of breast cancer has increased recently and it is estimated that there will be more than 51.000 breast cancer cases by 2012 (Ozmen, 2008).

While breast cancer is encountered so commonly, it has a slow growth rate and notable treatment outcomes can be achieved by early diagnosis. Early diagnosis and treatment of breast cancer can both be effective in extending life expectancy, reducing mortality, increasing quality of life, and preventing physical pain and psychosocial problems in women (İğci & Asoğlu, 2003).

Screening methods such as mammography, clinical breast examination, and breast self-examination (BSE) are described as health improvement activities and play important roles in the early diagnosis of breast cancer (Semiglazov et al., 1999; Smith et al., 2003; Ministry of Health in Turkey, 2006). While mammography is the sole effective diagnostic method for reduction of mortality in breast cancer, it is not regarded as a suitable modality for poor countries due to its costly nature and requirement of technical specialty along with man power. Therefore,

development of awareness and consciousness on breast health among women is considered to be the most effective and important action (Anderson et al., 2003; Smith et al., 2003; McCready et al., 2005). BSE is recommended to be performed routinely on a monthly basis in all the women aged above 20 years and the importance of raising awareness on breast cancer via BSE is noted (Smith et al., 2003).

Currently, while investigating the roles of beliefs and perceptions in preventive health behaviors such as BSE practice and undergoing mammography for breast cancer screening, the theoretical structure of Health Belief Model (HBM) is used (Jane, 1995; Wu, 2003; Gözümlü et al., 2004). Particularly in countries where Islam is the predominant religion, such as Turkey, generally women abstain from touching even their own breasts, do not want to go to a physician for breast examination, delay their visits for breast examination, and feel embarrassed of undergoing mammography and being examined by a physician (Rajaram & Rashidi, 1999).

In Turkey, where the vast majority of people are muslim, the aim of the education delivered to the students in nursing and midwifery schools at universities, is to raise the awareness of young girls towards their own body, establish a regular BSE pattern among them by having a

¹ Nursing Department, School of Health Sciences, Çanakkale Onsekiz Mart University, ² Nursing Department, Faculty of Health Sciences, Marmara University, Turkey *For correspondence : arzutunam@gmail.com

positive influence over their health beliefs, and encourage them to share the related knowledge with their relatives and the public for maintaining a healthier community (Aydın Avcı & Keskin, 2005; Beydağ & Karaoğlan, 2007; Aydın Avcı et al., 2008).

In a study where the effect of health beliefs of midwifery students over BSE were analyzed, as the perceived susceptibility and confidence (self-reliability) were found to be at moderate level, health motivation and perceived benefits were found to be high, while perceived barriers was observed to be at low levels (Aydın Avcı et al., 2008).

Nursing students undergoing breast cancer and BSE education, perform BSE more regularly than those who do not take these lessons. However, there are also educated students who still perform BSE irregularly or not at all because of forgetfulness, embarrassment, fear and lack of knowledge, considering one's age early for starting breast examination, or disregard (Uzun et al., 2004; Aydın Avcı & Keskin, 2005).

Nurses and midwives who are taught with a philosophy of maintaining and developing public health, play an important role in teaching and promoting BSE among women. Since the nursing and midwifery students trained on BSE are around their twenties, it is expected from them to perform BSE each month and begin to share their knowledge on this subject with the public during their undergraduate years. Defining the beliefs of students with regard to breast cancer may shed a light on future studies focusing on changing wrong beliefs and increasing the efficiency of public health trainings that will be given by them. Therefore, this is a descriptive survey investigating health beliefs and BSE teaching status of nursing and midwifery undergraduate students who are taught on breast cancer and BSE.

Materials and Methods

Our survey was carried out on 113 third-year and fourth-year nursing and midwifery students between September - December 2009. Required permissions and approvals were obtained from the institutional governing bodies and students. Our study population had received information about breast cancer and BSE 1 or 2 years previously.

As a data collection method, sociodemographic characteristics of the 113 students were determined. This form included information on students involving their age, university term, family history of breast cancer, frequency of BSE practice, and the factors influencing that frequency. Moreover, the students were evaluated in terms of providing information on BSE and breast cancer and teaching BSE.

The study population was consisted of 113 undergraduate students in a School of Health. 59.3% of the study group were third-year and fourth-year midwifery students (n=46), whereas 40.7% were students of nursing (n=67). Sociodemographic characteristics were analyzed by reviewing the related literature. We used the 20-item breast cancer and BSE knowledge form (8 on breast cancer and 12 on BSE) with a score range of 0-100 which

was created by Maurer in 1997. This form was adapted to Turkish language by Tuna Malak and Dicle in 2007 (Maurer, 1997; Tuna Malak & Dicle, 2007).

Champion's Health Belief Model Scale (CHBMS) was used in order to examine the influence of health beliefs of students over BSE practice.

Champion's Health Belief Model Scale (CHBMS) is an instrument for measuring the beliefs and attitudes of women towards breast cancer and breast self-examination. This 42-item test was developed by Champion in 1984 and revised in the following studies (1993, 1997, 1999).

Susceptibility domain (or subscale) has 3 items and is about the degree of perceived risks and vulnerability. Seriousness domain has 7 items and indicates the degree of perceived consequences of developing breast cancer. Benefits domain has 4 items and indicates perceived advantages associated with BSE. Barriers domain has 11 items and indicates the perceived barriers associated with BSE. Confidence domain has 10 items and indicates the perceived capability or efficiency in practicing BSE for detection of breast masses. Health motivation domain has 7 items and indicates the general interest and concerns towards personal health status.

Reliability and validity studies of the original scale was carried out by Victoria Champion. Cronbach alpha coefficient (reliability coefficient) of the scale was found to be 0.69-0.90 for the subscales. Test-retest for the subscales of the survey varies between 0.45-0.70. The scale shows construct and content validity (Champion, 1993; Champion & Scott, 1997; Champion, 1999). Adaptation to Turkish language was performed by Gozum & Aydin (2004), Karayurt & Dramali (2007), and Secginli & Nahcivan (2004).

In this study, we preferred to use the Turkish form of CHBMS created by Karayurt Dramali (2007). In order to evaluate the reliability of this scale, item statistics and Cronbach alpha coefficients were analyzed. Since item-total correlations were above 0.30, no item was excluded. The consistency of the scale over time was evaluated by test-retest correlation and was observed to be between 0.89-0.99 for the subscales. Cronbach alpha reliability coefficients varied between 0.58-0.89. Confirmatory factor analysis was performed for evaluation of the construction validity and 6 factors similar to the ones in the original scale were found. Turkish adaptation of CHBMS was found to be a reliable and valid instrument for determination of beliefs and behaviors associated with breast cancer and BSE.

Six constructs of the health belief model formed the 6 subscales of the scale. Items were formatted with a 5-point Likert scale consisted of the following: (1) strongly disagree, (2) disagree, (3) undecided, (4) agree, (5) and strongly agree.

Each domain of the scale is evaluated separately and they are not combined to obtain a total score. Thus, a score is acquired for each of the domains. While the length of time required to complete the scale depends on the number of domains used, generally it is finished within 12-15 minutes when all domains are applied. The data were analyzed by SPSS 15.0 program. Sociodemographic data and BSE teaching status of students to the people around

them were expressed by percentages. Health beliefs of students was evaluated with t test.

Results

Personal characteristics of the students included in our study are shown in Table 1. Study population was comprised of students and the mean age was 21.4±1.43. All of the students were unwed. The total number of mothers, sisters, relatives, friends, and patients taught about BSE practice by all the students, was 167. While

Table 1. Data on Student Characteristics

| Characteristics (n=113) | n | % |
|--|-----|------|
| Third- and fourth-year students | | |
| Nursing students | 67 | 59.3 |
| Midwifery students | 46 | 40.7 |
| Family history of breast cancer | | |
| Positive | 3 | 2.7 |
| Negative | 110 | 97.3 |
| History of symptoms | | |
| Benign breast mass | 2 | 1.8 |
| Discharge | 1 | 0.9 |
| Negative | 110 | 97.3 |
| Knowledge on breast cancer and BSE | | |
| 52.34 ± 9.63 (range: 25 - 75) | | |
| BSE practice pattern | | |
| Regular | 37 | 32.7 |
| Irregular | 76 | 67.3 |
| Reasons behind irregular BSE practice* | | |
| Sparing no time | 14 | 18.4 |
| Perception of no susceptibility | 3 | 3.9 |
| Feeling no need | 17 | 22.3 |
| Fear of breast cancer diagnosis | 23 | 30.2 |
| Carelessness towards health | 19 | 25.0 |

* (n=76)

Table 2. People Who Received Instructions on BSE from the Students

| Person who received instructions | n | % |
|----------------------------------|-----|------|
| Mother/Sister | 106 | 63.5 |
| Relative | 48 | 28.7 |
| Friend | 7 | 4.19 |
| Patient | 6 | 3.59 |
| Total number of people | 167 | 100 |

Table 3. Subscale Scores of Health Belief Model Scale for BSE Among Nursing and Midwifery Students

| Subscales | Number of items | Range of score | $\bar{X} \pm sd$ |
|-------------------|-----------------|----------------|------------------|
| Susceptibility | 3 | 3-15 | 7.52±2.62 |
| Seriousness | 7 | 6-30 | 21.83±5.30 |
| Health motivation | 7 | 5-25 | 26.55±4.22 |
| BSE benefits | 4 | 4-20 | 16.73±4.45 |
| BSE barriers | 11 | 8-40 | 22.33±6.44 |
| BSE confidence | 10 | 10-50 | 40.28±6.67 |

Table 4. Comparison of Health Beliefs with Regard to Regular Bse Practice Among Students (N = 113)

| Regular BSE practice | Susceptibility | Seriousness | Health motivation | BSE benefits | BSE barriers | BSE confidence |
|----------------------|------------------|------------------|-------------------|------------------|------------------|------------------|
| | $\bar{X} \pm sd$ | $\bar{X} \pm sd$ | $\bar{X} \pm sd$ | $\bar{X} \pm sd$ | $\bar{X} \pm sd$ | $\bar{X} \pm sd$ |
| Yes (n=37) | 8.45±3.61 | 21.97±5.26 | 27.45±4.35 | 18.29±3.65 | 19.56±8.31 | 42.13±5.72 |
| No (n=76) | 7.06±1.83 | 21.76±5.35 | 26.11±4.12 | 15.97±4.63 | 23.68±4.82 | 39.38±6.94 |
| t | 2.72 | 0.19 | 1.59 | 2.67 | 3.32 | 2.08 |
| p* | 0.002* | 0.955 | 0.936 | 0.069 | 0.024* | 0.046* |

63.5% (n=106) taught BSE to their mothers and sisters, 28.7% (n=48) and 59% (n=6) taught their relatives and patients, respectively (Table 2). Results of evaluation of health belief scale are shown in Table 3. There was a statistically significant difference between the regular BSE practice and susceptibility, barriers, and confidence domains (p<0.05). Students who practiced BSE regularly had higher susceptibility than those who did not. Students with high susceptibility were found to have taught BSE practice to their relatives and patients. Students with high confidence values were found to practice BSE regularly, as well. Students who had positive family and symptoms history, were also observed to practice BSE regularly and have high confidence values.

BSE barrier scores of students who considered themselves as incapable to practice regular BSE were higher than those who practiced regularly. The significant difference in perceived BSE barriers was originating from the ones who did not practice BSE regularly. There was no significant difference between the perceived seriousness, health motivation, and BSE benefits in the entire study population (p>0.05) (Table 4).

Discussion

The percentage of nursing and midwifery students in Turkey who practice BSE regularly each month, vary between 31-75.4% (Uzun et al., 2004; Kılıç et al., 2006; Aydın Avcı et al., 2008). Studies on nursing and midwifery students reveal that despite having knowledge on BSE, they fail to practice BSE on a regular basis (Alsaif, 2004; Plesnicar et al., 2004). The information gained throughout their education, influences students' knowledge on BSE. Studies show that education has a positive effect over processes of knowing and acting (Attia et al., 1997; Thomas et al., 2002).

However, students can pass on their knowledge on BSE more easily to their relatives. Although it is important that students teach their patients about BSE in terms of maintenance and development of public health, the rate of students accomplishing this objective was very low (3.59%) in our study. In a study which supports our results, after providing BSE education to the students of a health school, the rate of BSE teaching to close relatives such as mother and sisters was observed to be higher than the rates determined prior to the BSE education (Gürsoy et al., 2009).

The susceptibility of regular BSE practicers was higher. Those students with higher perceived susceptibility taught their patients and relatives how to practice BSE. It is an expected behavior for a student with high perceived susceptibility to demonstrate care for public health and

teach BSE to the relatives (Aydın Avcı et al., 2008). Other studies have also revealed the fact that Turkish students considered beginning BSE practice during university as early (Uzun et al., 2004; Memis et al., 2009).

In a study performed on a group with a culture other than Turkish, the BSE beliefs of nursing students were found to have high perceived benefits, susceptibility, and health motivation scores, while exhibiting low perceived barriers (Budden, 1999). This result isn't consistent with our results. It is evident that nursing and midwifery students from different cultures may have different health beliefs for BSE practice.

Although the beliefs of students (n=113) towards practicing BSE on themselves vary depending on their cultural backgrounds, their efforts to teach BSE and raise awareness towards importance of early diagnosis among their relatives and family, are valuable. This result shows that students who receive healthcare education can be helpful in maintaining and developing public health even during their school years. The reason behind informing their relatives and family about breast cancer, BSE, and importance of early diagnosis was their fear that their relatives would contract it. However, nursing and midwifery students clearly did not show the same behavior towards their patients (n=6). When students were asked about this biased and contradictory behavior, they mentioned differing individual priorities of their patients or failure to remember under working conditions.

Those expressions show that nursing and midwifery students tend to be more efficient in teaching positive health behaviors to their first-degree relatives, while experiencing difficulties in teaching those behaviors, such as BSE, to patients with poor health status and being inefficient to instruct them about health behaviors unrelated to their priorities. In light of the results of our study, we can say that regular training sessions underscoring the importance of breast cancer and self-breast examination, will improve perceived susceptibility in students and their patients/relatives, while showing a positive impact over the health beliefs of students. Repeating the importance of regular BSE practice frequently to students and using warning posters and videos about BSE practice, may help to elevate the perceived susceptibility among students. Students should be encouraged to provide trainings to patients for teaching and promoting positive health behavior.

References

- Alsaiif AA (2004). Breast self examination among Saudi female nursing students in Saudi Arabia. *Saudi Med J*, **25**, 1574-8.
- Anderson BO, Braun S, Lim S, et al (2003). Early detection of breast cancer in countries with limited resources. *Breast J*, **9**, 51-9.
- Attia AK, Rahman DAMA, Kamel LI (1997). Effect of an educational film on health belief model and breast self examination practice. *East Med Hlth J*, **3**, 435-43.
- Aydın Avcı İ, Altay B, Kocaturk B (2008). Midwifery students' health beliefs intended for breast self examination. *Meme Sağlığı Dergisi*, **4**, 25-8.
- Aydın Avcı İ, Keskin T (2005). Nursing students' health beliefs intended for breast self examination. *O.M.U.Tip Dergisi*, **22**, 146-50.
- Beydağ K, Karaoğlu H (2007). Effect of breast self examination education to the knowledge and attitudes of female students. *TSK Koruyucu Hekimlik Bülteni*, **6**, 106-11.
- Budden L (1999). Student nurses' breast self-examination health beliefs, attitudes, knowledge, and performance during the first year of a preregistration degree program. *Cancer Nurs*, **22**, 430-7.
- Champion V (1999). Revised susceptibility, benefits and barriers scale for mammography screening. *Res Nurs Hlth*, **22**, 341-8.
- Champion VL, Scott C (1997). Reliability and validity of breast cancer screening belief in African American women. *Nurs Res*, **46**, 331-7.
- Champion VL (1993). Instrument refinement for breast cancer screening behaviors. *Nurs Res*, **42**, 139-43.
- Darendeliler E, Ağaoğlu FY (2003). Meme Kanserinin Epidemiyolojisi ve Etiyolojisi. Eds: Topuz E, Aydın A, Dincer M. *Meme Kanseri.İstanbul: Nobel Tıp Kitabevleri*, 13-33.
- Gözüm S, Aydın I (2004). Validation evidence for Turkish adaptation of champion's health belief model scales. *Cancer Nurs*, **6**, 491-8.
- Gözüm S, Karayurt Ö, Aydın İ (2004). The results of Turkish adaptations of champion's health belief model scale at breast cancer screening. *Hemşirelikte Araştırma Geliştirme Dergisi*, **1**, 2, 71-85.
- Gürsoy AA, et al (2009). A different approach to breast self-examination education: daughters educating mothers creates positive results in Turkey. *Cancer Nurs*, **32**, 127-34.
- İğci A, Asoğlu O (2003). Meme Kanserinin Erken Tanısında Tarama Yöntemleri. Eds: Topuz E, Aydın A, Dincer M. *Meme Kanseri. İstanbul: Nobel Tıp Kitabevleri*, 113-23.
- Jane Lu ZY (1995). Variables associated with breast self-examination among chinese women. *Cancer Nurs*, **18**, 29-34.
- Karayurt Ö, Dramalı A (2007). Adaptation of champion's health belief model scale for Turkish women and evaluation of the selected variables associated with breast self-examination. *Cancer Nurs*, **30**, 69-77.
- Kılıç S, et al (2006). Determination of the knowledge and practice level of breast self-examination among the nurses of the GATA hospital, the students of the GATA nursing school and some female patients. *Gülhane Med J*, **48**, 200-4.
- Maurer F (1997). A peer education model for teaching breast self-examination to undergraduate college women. *Cancer Nurs*, **20**, 49-61.
- McCready T, Littlewood D, Jenkinson J (2005). Breast self-examination and breast awareness: a literature review. *J Clin Nurs*, **14**, 570-8.
- Memis S, Balkaya NA, Demirkiran F (2009). Knowledge, attitudes, and behaviors of nursing and midwifery students regarding breast self-examination in Turkey. *Oncol Nurs Forum*, **36**, 39-46.
- Ministry of Health in Turkey (2006). Control programme of breast cancer: Education on breast self examination. Available from: <http://www.ukdk.org/pdf/kitap/27.pdf>,
- Ozmen V (2008). Breast cancer in the world and Turkey. *Meme Sağlığı Dergisi*, **4**, 7-12.
- Plesnicar A (2004). Golicnic M. Kralj B. Midwifery students and breast self examination. *Breast J*, **10**, 560.
- Rajaram SS, Rashidi A (1999). Asian-Islamic women and breast cancer screening: a socio-cultural analysis. *Women Hlth*, **28**, 45-58.
- Seğginli S, Nahcivan N (2004). Reliability and validity of the breast cancer screening belief scale among Turkish women. *Cancer Nurs*, **27**, 1-8.
- Semiglazov VF, et al (1999). Role of breast self examination in early detection of breast cancer: Russia/WHO prospective

- randomized trial in St. Petersburg. *Cancer Strategy*, **1**, 145-51.
- Smith RA, et al (2003). American cancer society guidelines for breast cancer screening: update 2003. *CA Cancer J Clin*, **53**, 141-69.
- Thomas B, et al (2002). Breast health educational interventions. Changes in beliefs and practices of working women. *AAOHN J*, **50**, 460-7.
- Tuna Malak A, Dicle A (2007). Assessing the efficacy of a peer education model in teaching breast self examination to universty students. *Asian Pac J Cancer Prev*, **8**, 481-4.
- Uzun Ö, Karabulut N, Karaman Z (2004). Knowledge and practices of nursing students about breast self-examination. *Atatürk Üniversitesi Hemşirelik Yüksekokulu Dergisi*, **7**, 1, 10-8.
- Wu TY, Yu MY (2003). Reliability and validity of the mammography screening beliefs questionnaire among Chinese American women. *Cancer Nurs*, **26**, 131-42.