

RESEARCH COMMUNICATION

Dietary and Lifestyle Characteristics of Colorectal Cancer in Jordan: a Case-control Study

Mostafa A Arafa^{1*}, Mostafa I Waly², Sahar Jriesat³, Ahmed Al Khafajei⁴, Sunny Sallam⁵

Abstract

Objectives: We evaluated dietary pattern and lifestyle characteristics of patients with colorectal cancer (CRC) in Jordan. **Design:** The case-control study included 220 recently diagnosed CRC cases and 220 age and gender matched healthy subjects as a control group. **Results:** The participating CRC cases had lower dietary intake of fibre, folate, vitamin B12, β -carotene, vitamin C and selenium as compared to controls ($P < 0.05$). The frequency of consumption of fruits and vegetables was also lower among CRC cases, while the frequency of consumption of red meat and saturated fat was higher and positively associated with CRC risk. Furthermore, family history for CRC played a positive role and the majority of CRC cases and controls had a low physical activity level. **Conclusions:** A sedentary lifestyle and a diet low in fruits and vegetables, and high in animal red meat and saturated fat, appeared associated with CRC among the studied Jordanian subjects. This is consistent with the reported CRC studies in developed nations indicating global causal effects for this tumour type.

Keywords: Colorectal cancer - diet - lifestyle - case-control - Jordan

Asian Pacific J Cancer Prev, 12, 1931-1936

Introduction

Cancer is considered a leading cause of death worldwide, accounting for 7.6 million of all deaths from a total of 58 million deaths reported in year 2005 and the world health organization (WHO) estimates that there will be 20 million new cases and 12 million deaths from cancer alone by the year 2020 (WHO, 2006; Stewart & Kleihues, 2003). The prevalence of cancer is higher in developed nations, yet 70% of all cancer deaths in 2005 occurred in developing countries (WHO, 2005). Recent reports indicating that migrant populations moved from countries of low cancer incidence to the west adopted the high cancer incidence trend of the new country (Hardling et al., 2009; Ronellenfistch et al., 2009; Sung et al., 2005). The international experts from the recent World Cancer Research Fund Report, 2007, concluded that CRC is a major public health problem worldwide and modernization in food choices, sedentary life style and increased life expectancy are the major factors that might synergize with genetics for the epidemic of CRC worldwide (WCRF, 2008).

There are several established dietary risk factors for CRC, largely related to folate deficiency (Choi and Mason, 2000), diets from animal sources that are low in

fibre and high in fat (Willett et al., 1990; Slattery et al., 1997). In addition experimental evidence from animal studies suggests that reactive oxygen species production alter DNA and lipid membrane structures in proliferated cells, suggesting a protective effect of antioxidants nutrients against different types of cancer, including CRC (Greenberg et al., 1994; La Vecchia et al., 2001).

Non dietary risk factors for CRC include, decreased physical activity (Friedenreich and Orenstein, 2002), obesity, a body mass index (BMI) of greater than or equal to 30 kg/m² (Calle et al., 2003; Jass, 2005), type 2 diabetes mellitus (Campbell et al., 2010), genetic susceptibility (Heavy et al., 2004; Frezza et al., 2006), alcohol drinking (Wu et al., 1987; Shimizu et al., 2003), smoking (Uldrich et al., 2001) and exposure to environmental carcinogens that may lead to proliferation and malignant transformation of colorectal cells (IARC 1993; Irigaray et al., 2007).

In spite of considerable public health significance of CRC, a recent study in Jordan revealed that most study participants are not well informed about CRC and early screening (Omran and Ismail, 2010). Dietary studies, pertinent to colorectal cancer, in our region are limited, where life style related factors, mainly the diet and physical activity, are much different than developed countries. In Jordan there was no reported

¹Princess Aljohara Alibrahim Center for Cancer Research, King Saud University, Riyadh, Saudi Arabia, ²Food Science and Nutrition Department, College of Agricultural and Marine Sciences, Sultan Qaboos University, Muscat, Sultanate of Oman, ³Ministry of Health, Data Management Department, Jordan Cancer Registry, ⁴Public Health Department, Faculty of Medicine, Jordan University for Science & Technology, Amman, Jordan, ⁵Epidemiology Department, High Institute of Public Health, Alexandria University, Egypt
*For correspondence: mostafaarafa@hotmail.com

studies conducted to investigate the association between diet, lifestyle factors and CRC risk. Therefore this case-control study was planned to evaluate the dietary pattern, sociodemographic and other lifestyle risk factors of CRC patients as compared to control subjects in Jordan.

Materials and Methods

Study participants

A case-control approach was used to conduct this study between the period of February 2008 and through January 2009 at Al-Bashir Hospital, a national referral hospital and the principal governmental center for CRC registry and therapy in Jordan. This study included all recently diagnosed and histologically confirmed CRC cases within the specified period of the study and they were 118 males and 102 females. The controls were selected from those attending the outpatient departments at the same hospital and they were a total of 600 controls sampled, 417 of whom were eligible to participate based on the study inclusion criteria; being free of gastrointestinal diseases and have no previous diagnosis of CRC or other types of cancers. Of the 417 eligible controls, 220 were interviewed based on age and gender matching criteria to CRC cases. All female cases and controls were not currently pregnant or lactating.

Ethical approval

This study was approved by the Institution Research Board (IRB) of the High Institute of Public Health, Alexandria University, Egypt and Research Ethics Committee of the Ministry of Health in Jordan. All patients and controls gave informed, written consent for enrollment in the study.

Data collection

In-person interviews were scheduled for study participants (cases and controls). They were asked to complete the study questionnaire included questions related to: (I) Sociodemographic data, smoking, alcohol drinking, family, history of CRC, vitamins supplement use, monthly income and physical activity (II) Dietary intake using a food frequency questionnaire (FFQ). Appetite problems (poor, good or excellent) were also reported. (III) Anthropometric measurements of cases and controls.

Anthropometric measurements

For all participants; height was measured to the nearest 0.5 cm and weight was recorded to the nearest 0.5 kg using a precision scale. Body mass index (BMI) for every subject was calculated as weight in kilograms divided by height in meters squared (kg/m^2) and categorized as normal ($\text{BMI}=18-25$), overweight and obese ($\text{BMI}>25$).

Clinical presentation for cases with colorectal cancer

Cases were inquired about date of first complaint and the interval of time between the appearance of first sign and/or symptom and visit to health provider confirming diagnosis. Basis of confirmed diagnosis, staging and diagnostic facility information (hospitals and laboratories)

were collected from medical records. Clinical data were collected and coded by following of the International Classification of Diseases for Oncology (ICD-O-3) as a dual classification with coding system for both topography and morphology.

Dietary assessment

To overcome the problem of reverse causality, that cancer patients might change their dietary habits after the diagnosis, we included only newly diagnosed cases in which diagnosis was confirmed 10-14 days prior to dietary intake questionnaire. The retrospective dietary intake of the study participants was estimated using a semi-quantitative FFQ (Block et al., 1990) where, cases and controls were asked to report the frequency (how often) and portion size for each food item consumed during a period of 12-months prior CRC diagnosis for cases and prior being interviewed for controls, this period was chosen to take into account seasonal variation in food consumption. Also, all study subjects were inquired if they have changed their diet than the usual routine in the last 12 months. The FFQ was adapted according to portion sizes based on commonly used household serving units/utensils in Jordan, and was tested for its validity, reliability and reproducibility before conducting the study (Pietinen et al., 1988).

The different food groups included in the questionnaire were 9 groups as follow: Breads/cereals, vegetables, fruits, meat/meat substitutes, milk/dairy products, deserts, beverages, sandwiches, and traditional Jordanian dishes. The collected dietary data were categorized into: (a) Dietary habits of CRC Jordanians as compared to their matched controls, (b) Food groups' analysis: The numbers of daily servings of food groups, based on the frequency of consumption, for all respondents were subsequently grouped according to the Food Guide Pyramid from US Department of Agriculture (USDA, 2005). (c) Daily intake of nutrients: The Food Processor software version 10.2 (ESHA Research, Salem, OR, USA) was used to calculate means of daily nutrient intakes of macronutrients, fiber, and specific micronutrients as estimated from the frequency of consumption, reported portion size, and nutrient content for all foods reported by each participant. (d) The percentage of kilocalories (Kcal) contribution to the total energy intake from protein, carbohydrate and fat were calculated for each participant.

Physical activity assessment

Physical activity was expressed as either low, moderate or high and calculated according to World Health Organization (Global Physical Activity Questionnaire) (WHO, Physical activity/GPAQ v2.0) as detailed below: Total Physical Activity = $[(P2 * P3 * 8) + (P5 * P6 * 4) + (P8 * P9 * 4) + (P11 * P12 * 8) + (P14 * P15 * 4)]$.

In the equation; constant 8: means vigorous metabolic equivalents (MET) in both domains (work and recreational). Constant 4: means moderate MET in both domains (work and recreational). One MET is defined as 1 kcal/kg/hour and is equivalent to the energy cost of sitting quietly.

The three physical activities suggested for classifying populations were low, moderate, and high. The criteria for

these are shown below:

High: If $(P2 + P11) \geq 3$ days & total physical activity MET minutes per week is ≥ 1500 . OR: IF: $(P2 + P5 + P8 + P11 + P14) \geq 7$ days & total physical activity MET minutes per week is ≥ 3000 .

Moderate: If: $(P2 + P11) \geq 3$ days & $((P2 * P3) + (P11 * P12)) \geq 60$ minutes. OR IF: $(P5 + P8 + P14) \geq 5$ days & $((P5 * P6) + (P8 * P9) + (P14 * P15)) \geq 150$ minutes OR • IF: $(P2 + P5 + P8 + P11 + P14) \geq 5$ days & total physical activity MET minutes per week ≥ 600 .

Low: If the value does not reach the criteria for either high or moderate levels of physical activity.

Statistical analysis

Data analysis was done using The Statistical Package for Social Sciences, for personal computer, (SPSS-PC) version 17. The chi-squared test was used to analyze categorical variables and the unpaired Student’s t-test for analyzing continuous variables. P-value <0.05 was considered statistically significant. The Logistic Regression Analysis, stepwise method was used as a multivariate analysis, where the dependent variable was being either cases or controls. The procedure began by selecting a group of variables that were associated with the disease status. First variable was selected for regression model, and then coupled with the next variable and so on. The procedure continued until all variables were selected or no additional variables provided an improvement in the model.

Results

There were 220 (118 males and 102 females) CRC cases and 220 matched controls for age and gender. The CRC cases and controls were similar in age (56.3 ± 12.3 years for males and 53.7 ± 6.8 years for females) and gender (53.6 % males & 46.4% females). As presented on table 1; the CRC female cases and controls had no significant difference with regards to BMI ($t=1.33$, $P>0.05$), mean while, BMI for males of controls group was higher than that of CRC cases, $t=2.78$, $P<0.05$.

CRC cases and controls had comparable monthly income, educational level, marital status, and residence status. None of the study participants consumed either multi-vitamin supplements or alcoholic beverages and the majority (67.7% for CRC cases and 71.8% for controls) never smoke, table 1. The assessment of physical activity levels (low, moderate and high) and routine exercise practice were comparable among CRC cases and controls group, table 1.

Table 2 shows the common dietary habits of the study participants, the majority of the study sample consumed fresh green salad on daily basis 80% and 85% for CRC cases and controls group respectively compared to 20% of cases and 15% of control group who consumed fresh green salads 1-2 times/week, the difference was not significantly different, $P>0.05$. CRC cases and controls had a similar dietary pattern with regard; adding olive oil to foods, adding lemon to beans and salads, type of bread eaten, and citrus fruits consumption, Table 2. All study participants reported a good appetite for eating and

Table 1. Lifestyle Characteristics of CRC Cases and Controls

Characteristic	Cases**		Controls**		Test of significance
	No.	%	No.	%	
Physical Activity Level					$\chi^2= 1.98$, P=0.372
Low	170	77.3	166	75.5	
Moderate	40	18.2	37	16.8	
High	10	4.5	17	7.7	
Routine Exercise Practice					$\chi^2= 21.74$, P=0.000*
Yes	40	18.2	84	38.2	
No 1	80	81.8	136	61.8	
Smoking History					$\chi^2= 19.24$, P=0.000*
Current Smoker	48	21.8	60	27.3	
Ex-Smoker	23	10.5	2	0.9	
Never Smoke	149	67.7	158	71.8	
First Degree Family History of CRC					COR=2.34 1.04 -5.27
Yes	20	9.1	9	4.1	
No	200	90.9	211	95.9	
Body mass index (kg/m ²) Mean \pm S.D					
Male	23.9 \pm 3.6		25.3 \pm 4.1		P=0.006*
Female	27.3 \pm 5.8		28.5 \pm 7.05		P=0.185

*Significant (P<0.05); **n=220; COR, Crude Odds Ratio

Table 2. Dietary Habits of CRC Cases and Controls

Characteristic	Cases**		Controls**		Test of significance
	No.	%	No.	%	
Drinking Tea					$\chi^2=3.96$, P=0.046*
After meal	209	95.0	198	90.0	
Not drinking	11	5.0	22	10.0	
Adding lemon to beans and salads					$\chi^2=1.13$, P=0.288
Yes	132	60.0	121	55.0	
No	88	40.0	99	45.0	
Fresh green vegetables consumption					$\chi^2=1.90$, P=0.17
Daily	176	80.0	187	85.0	
1-2 times/week	44	20.0	33	15.0	
Adding olive oil to foods					$\chi^2=0.18$, P=0.67
Yes	209	95.0	207	94.0	
No	11	5.0	13	6.0	
Type of bread eaten					$\chi^2=0.87$, P=0.35
White bread	202	92.0	207	94.0	
With Zatar	18	8.0	13	6.0	
Citrus fruits					$\chi^2= 0.48$, P= 0.49
Daily	143	65.0	136	62.0	
1-2 times/week	77	35.0	84	38.0	

*Significant (P<0.05); **n=220

consuming foods without any preference.

Foods group analysis (number of daily servings as compared to USDA Food Guide Pyramid) is represented in Table 3. It shows that the daily intake of servings from various food groups based on the USDA Food Guide Pyramid. Highly significant differences were observed from the vegetables, fruits and meat groups between CRC cases and controls (P<0.05), where for CRC cases, 30 % consumed than 3 servings vegetables/day, 43% consumed less than 2 fruits servings/day, and 43.6% consumed more than 3 servings of meat/day. On the other hand, no significant differences were observed in the daily intake of servings from the cereals and milk groups (P>0.05).

The percentage of energy from different macronutrients analysis revealed that for CRC cases, approximately 30% of dietary calories in the standard diet consists of fat, that is statistically higher than controls (P<0.05). The

Table 3. Frequency of Food Consumption of CRC Cases as Compared to Controls

Food group ¹	Cases**		Controls**		Test of significance
	No.	%	No.	%	
Bread, Cereal, Rice& Pasta Group					
<6	1	0.45	2	0.91	$\chi^2=1.66$ P=0.44
6-11	217	98.6	213	96.8	
≥ 11	2	0.91	5	2.27	
Vegetables Group					
<3	65	29.6	10	4.54	$\chi^2=48.63$ P=0.000*
3-5	81	36.8	109	49.5	
≥ 5	74	33.6	101	45.9	
Fruits Group					
<2	94	42.7	36	16.4	$\chi^2=37.09$ P=0.000*
2-4	97	44.1	136	61.8	
>4	29	13.2	48	21.8	
Milk, Yogurt & Cheese Group					
<2	17	7.7	26	11.8	$\chi^2=2.34$ P=0.31
2-3	196	89.1	189	85.9	
≥ 3	7	3.18	5	2.27	
Meat, Poultry, Fish, Dry Beans, Egg & Nuts Group					
<2	17	7.73	40	18.2	$\chi^2=33.18$ P=0.000*
2-3	107	48.6	137	62.3	
≥ 3	96	43.6	43	19.6	

¹number of servings/day; *Significant (P<0.05); **n=220

Table 4. Energy from Different Macronutrients and Mean Daily Intake of Different Fat Types

Variable	Cases**	Controls**	Statistical Analysis	
Percentage of energy				
% protein	23.7±0.16	22.3±0.15	t=94.0,	P=0.000*
% fat	26.9±0.52	24.4±0.72	t= 40.8,	P=0.000*
% carbohydrate	49.4±2.15	53.2±1.61	t=21.2,	P=0.000*
Fat Types (g/day or mg)				
Saturated	18.0± 9.80	13.3±7.28	t-test=5.67,	P=0.000*
Monounsaturated	16.5±12.3	13.9±9.83	t-test=2.46,	P=0.014*
Polyunsaturated	8.5±6.2	5.8±5.1	t-test=4.98,	P=0.000*
Cholesterol	159±153	111±170	t-test=3.07,	P=0.002*

Data presented as Mean ± S.D, *Significant (P<0.05); **n=220

same pattern was observed for the percentage of energy derived from protein and carbohydrate among CRC cases as compared to controls, P<0.05, Table 4. The different types of fats consumed daily and contributing to total daily fat amount is also presented in Table 4, and it is observed that the daily intake from saturated, mono and polyunsaturated fats and cholesterol is significantly higher among CRC cases as compared to controls, (P<0.05).

The mean daily intake of fibre for CRC patients (4.63 ± 5.46 g/day) was significantly lower than the control (18.8 ± 3.1 g/day), t= 33.47, (P<0.05), and lower than the RDA for fibre (25 g/day). The mean daily intake of vitamins and minerals for CRC patients was comparable to controls and within RDA values with the exception to the folate and vitamin B12 for CRC patients (189.2 ± 11.7 µg/day and 1.09 ± 1.1 µg/day respectively) was significantly lower than the controls group (212.5 ± 13.1 µg/day and 1.49 ± 1.3 µg/day respectively), (t= 19.68 and t = 3.48 respectively), (P<0.05). Furthermore, β-carotenes (1946.96 ± 1485.18 mg/day for CRC cases and 2257.23 ± 1561.90 mg/day for controls), vitamin C (45.48 ± 16.81 µg/day for CRC cases and 64.78 ± 13.74 µg/day for controls), and selenium (38.75 ± 11.42 µg/day for CRC cases and 59.26 ± 8.91

µg/day for controls), were significantly lower, (t= 2.13, t= 13.19 and t = 21.00 respectively), (P<0.05).

The association between nutrients intake as independent predictors for CRC risk was determined using the multivariate analysis, stepwise method for logistic regression analysis, where the dependent variables were either CRC cases or controls. The model showed that: total fat: OR=1.46 (1.15– 1.84), potato: OR=1.77 (1.18 – 2.66), sweets and cakes: OR= 2.33 (1.67– 3.25), fruits: OR=1.80 (1.26 – 2.58), red meat: OR=2.66 (1.83 – 3.88), saturated fat: OR=1.03 (1.01– 1.05), were the higher independent predictors for CRC among the interviewed groups. Meanwhile, calcium: OR = 0.99 (0.99 – 1.00), green leafy vegetables: OR = 0.99 (0.99 – 1.00), vegetables: OR = 0.99 (0.99 – 0.99) and eggs: OR =0.89 (0.84 – 0.94), were marginally independent predictors for CRC among interviewed groups.

Discussion

Diet represents the most significant environmental factor to which humans are exposed on a regular basis over prolonged periods of time and its impact in the etiology or primary prevention of CRC had been extensively studied in developed nations. Few dietary population-based studies were conducted at the Middle East, yet reported high consumption of red meat, fast foods and saturated dietary fats and a decreased consumption of fresh vegetables and fruits, a typical high risk dietary profile for CRC (Musaiger and Miladi, 2001; Galal, 2003). In Jordan, CRC cases accounted for 12.1% of all newly diagnosed cancer cases in 2007, and ranked the first among all new cancers among males and second among females (Tarawneh and Nimri, 2007). The results of our study suggested three associations regarding the modifiable risk factors associated with CRC risk in Jordan:

A family history of CRC is recognized as an important risk factor for the CRC incidence (Fuchs et al., 1994) and our data showed a significant high risk for CRC among those of first degree of family history for CRC. In previous reports physical activity have been related to risk of CRC (Byers et al., 2002, Giovannucci et al., 1997), our data support this link where it was observed a trend of decreased physical activity among CRC cases, that is similar to the recent published meta-analysis report (Harriss et al., 2009). Unlike the published results at developed nations with regard to positive association between obesity and CRC risk (Calle et al., 2003, Adami and Trichopoulos, 2003), obesity was not prevalent among the study male groups for both CRC cases and controls, meanwhile female subjects for both groups had higher mean BMI, < 30, with no statistical differences.

Fruits and vegetables are good sources for different mineral, vitamins and dietary fiber. We observed that the daily fiber intake of the studied subjects was lower than the recommended dietary allowances (RDA) of fiber (25 g/day). This is in accordance with numerous epidemiological studies suggested that populations with diets low in fiber have higher CRC rates (Sheila et al., 2003; Timothy et al, 2009). On the other hand, the recent WCRF report did not support this relation (WCRF, 2008), in addition to

two large prospective studies, the Iwo Women's Health Study (Negri et al., 1998), and the Health Professionals Follow-Up Study (Steinmetz et al., 1994) have shown no significant protective effect of dietary fiber against CRC. In addition controlled clinical studies had revealed that high consumption of dietary fiber did not reduce polyp recurrence among patients with recently detected colorectal adenomas (Giovannucci et al., 1994).

The CRC patients had lower daily intake than the RDA for dietary antioxidants; vitamin C (60 mg/day) and selenium (55-70 μ g/day). This increases the risk of oxidative stress and subsequent cellular damage by reactive oxygen species; this has been involved in the pathology of many non communicable diseases including cancer (Baron, 2005; Schrauzer et al., 1977; Johnson, 2001). Dietary folic acid and vitamin B12 levels for CRC cases were lower than the RDA (400 and 2.4 μ g/day respectively) this increase the risk of CRC as reported by a recent meta-analysis of prospective cohort studies indicated a significant reduction in CRC risk for subjects in the highest quintile of dietary folate and vitamin B12 intake compared with the lowest (Sanjoaquin et al., 2005).

Few prospective studies showed a positive association between animal protein intake and colorectal cancer risk (Cross et al., 2010; Alexander et al., 2009). Likewise our study shows that the consumption of red meat among CRC patients was higher than controls and associated with high risk for CRC. It is quiet interested that the methods of cooking of meat in Jordan is mainly charring and boiling and it is the same for CRC patients and controls with no significant differences, attributed that cooking methods are not associated with CRC incidence unlike some Western studies that reported a positive association (Gunter et al., 2005).

Among the studied subjects, contribution of saturated fat to the daily intake and total energy intake was higher in CRC patients as compared to controls and increases their total energy intake which is positively correlated with CRC incidence as indicated by previous reports (Schatzkin et al., 2000; Giovannucci et al., 1994).

In conclusion, our results indicated that the CRC cases enrolled in this study, were adopting a sedentary lifestyle and their dietary pattern was high frequency of red meat and saturated fat consumption and low consumption of fruits and vegetables. In addition they had inadequate daily intake of four key nutrients that protect against CRC risk, vitamin C, selenium folate and B12. These findings are similar to risk factors associated with increased CRC incidence at developed nations. Further prospective population-based studies are needed to establish a base line dietary intake data for the Jordanian population.

Acknowledgements

The authors deeply extend their appreciated acknowledgment to all the staff of all Jordanian Health facilities, the Director of the Jordan Cancer Registry and the Middle East cancer Consortium for their dedicated effort to complete this study. The authors indicated no potential or actual conflict of interest pertaining to this submission. All authors read and approved the

final manuscript. Prof. Mostafa Arafa and Dr. Mostafa Waly have made equal contribution to study design, interpretation of data, and drafting the manuscript. Dr. Sahar Jriesat, has made full contribution to data collection, acquisition of data and statistical analysis. Prof. Sunny Sallam and Prof. Ahmed Al-Khafajei conceived the study and participated in its design and work coordination. This study has some limitations and subjected to bias as being a hospital based- case control study, small sample size and memory issues related to recall of foods reported in FFQ.

References

- Adami HO, Trichopoulos D (2003). Obesity and mortality from cancer. *N Engl J Med*, **348**, 1623-4.
- Alexander DD, Cushing CA, Lowe KA, et al (2009). Meta-analysis of animal fat or animal protein intake and colorectal cancer. *Am J Clin Nutr*, **89**, 1402-9.
- Baron JA (2005). Dietary fiber and colorectal cancer: a meta analysis study, an ongoing saga. *J Am Med Ass*, **294**, 2904-6.
- Block G, Hartman AM, Naughton D (1990). A reduced dietary questionnaire: development and validation. *Epidemiology*, **1**, 58-64.
- Byers T, Nestle M, McTiernan A, et al (2002). American Cancer Society and Nutrition and Physical Activity Guidelines Advisory Committee. American Cancer Society guidelines on nutrition and physical activity for cancer prevention: Reducing the risk of cancer with healthy food choices and physical activity. *CA Cancer J Clin*, **52**, 92-119.
- Calle EE, Rodriguez C, Walker-Thrumind K, et al (2003). Overweight, obesity, and mortality from cancer in a prospectively studied cohort of US adults. *N Engl J Med*, **348**, 1625-38.
- Calle EE, Rodriguez C, Walker-Thrumind K, et al (2003). Overweight, obesity, and mortality from cancer in a prospectively studied cohort of US adults. *N Engl J Med*, **348**, 1625-38.
- Choi S, Mason JB (2000). Folate and carcinogenesis: an integrated scheme. *J Nutrition*, **130**, 129-32.
- Campbell PT, Deka A, Jacobs EJ, et al (2010). Prospective study reveals associations between colorectal cancer and type 2 diabetes mellitus or insulin use in men. *Gastroenterology*, **139**, 1138-46.
- Cross AJ, Ferrucci LM, Risch A, et al (2010). A large prospective study of meat consumption and colorectal cancer risk: an investigation of potential mechanisms underlying this association. *Cancer Res*, **70**, 2406-14.
- Friedenreich CM, Orenstein MR (2002). Physical activity and cancer prevention: etiologic evidence and biological mechanisms. *J Nutrition*, **132**, S3456-64.
- Frezza EE, Wachtel MS, Chiriva-Internati M (2006). Influence of obesity on the risk of developing colon cancer. *Gut*, **55**, 285-91.
- Fuchs CS, Giovannucci EL, Colditz GA, et al (1994). A prospective study of family history and the risk of colorectal cancer. *N Engl J Med*, **331**, 1669-74.
- Galal O (2003). Nutrition related health patterns in the Middle East. *Asia Pacific J Clin Nutr*, **12**, 337-43.
- Giovannucci E, Rimm EB, Stampfer MI, et al (1994). Intake of fat, meat, and fiber in relation to risk of colon cancer in men. *Cancer Res*, **54**, 2390-7.
- Giovannucci E, Colditz GA, Stampfer MJ, et al (1997). Physical activity, obesity and risk of colorectal adenoma in women (United States). *Cancer Causes Control* **7**, 253-63.
- Greenberg ER, Baron JA, Tosteson TD, et al (1994). A clinical

- trial of antioxidant vitamins to prevent colorectal adenoma. Polyp Prevention Study Group. *N Engl J Med*, **21**, 141-7.
- Gunter MJ, Probst-Hensch NM, Cortessis VK, et al (2005). Meat intake, cooking-related mutagens and risk of colorectal adenoma in a sigmoidoscopy-based case-control study. *Carcinogenesis*, **26**, 637-42.
- Harding S, Rosato M, Teyhan A (2009). Trends in cancer mortality among migrants in England and Wales, 1979-2003. *Eur J Cancer*, **45**, 2168-79.
- Harriss DJ, Atkinson G, Batterham A, et al (2009). Lifestyle factors and colorectal cancer risk (2): a systematic review and meta-analysis of associations with leisure-time physical activity. *Colorectal Dis*, **11**, 689-701.
- Heavy PM, McKenna D & Rowland IR (2004). Colorectal cancer and the relationship between genes and the environment. *Nutr Cancer*, **48**, 124-41.
- Irigaray P, Newby JA, Clapp R, et al (2007). Lifestyle-related factors and environmental agents causing cancer: an overview. *Biomed Pharmacother*, **61**, 640-58.
- IARC Working group on the Evaluation of Carcinogenic Risks to Humans (1993). Some naturally occurring substances: food items and constituents, heterocyclic aromatic amines and mycotoxins. Vol. 56. IARC monograph on the evaluation of carcinogenic risks to humans.
- Jass JR (2005). What is new in hereditary colorectal cancer? *Arch Pathol Lab Med*, **129**, 1380-4.
- Johnson IT (2001). Mechanisms and anticarcinogenic effects of diet-related apoptosis in the intestinal mucosa. *Nutr Res Reviews*, **14**, 229-56.
- La Vecchia C, Altieri A, Tavani A (2001). Vegetables, fruit, antioxidants and cancer: a review of Italian studies. *Eur J Nutr*, **40**, 261-7.
- Musaiger AO, Miladi (2001). Food consumption patterns and dietary habits in the Arab countries of Gulf. FAO/RNE, UAE University, Al-Ain UAE.
- Negri E, Franceschi S, Parpinel M, et al (1998). Fiber intake and risk of colorectal cancer. *Cancer Epidemiol Biomarkers Prev*, **7**, 667-71.
- Omran S, Ismail AA (2010). Knowledge and beliefs of Jordanians toward colorectal cancer screening. *Cancer Nursing*, **33**, 141-8.
- Pietinen P, Hartman AM, Haapa E, et al (1988). A self-administered food use questionnaire with a portion size picture booklet. *Am J Epidemiol*, **128**, 655-66.
- Physical Activity: Department of Chronic Diseases and Health Promotion Surveillance and Population-Based Prevention World Health Organization 20 Avenue Appia, 1211 Geneva 27, Switzerland For further information: www.who.int/chp/steps GPAQ v2.0.
- Ronellenfitsch U, Kyobutungi C, Ott JJ, et al (2009). Stomach cancer mortality in two large cohorts of migrants from the former Soviet Union to Israel and Germany: are there implications for prevention? *Eur J Gastroenterol Hepatol*, **21**, 409-16.
- Shimizu N, Nagata C, Shimizu H, et al (2003). Height, weight, and alcohol consumption in relation to the risk of colorectal cancer in Japan: a prospective study. *Br J Cancer*, **7**, 1038-43.
- Sanjoaquin MA, Allen N, Couto E, et al (2005). Folate intake and colorectal cancer risk: a meta-analytical approach. *Int J Cancer*, **113**, 8258.
- Schrauzer GN, White DA & Schneider CJ (1977). Cancer mortality correlation studies. III: Statistical associations with dietary selenium intakes. *Bioinorganic Chemistry*, **7**, 33-9.
- Schatzkin A, Lanza E, Corle D, et al (2000) Lack of effect of a low-fat diet on the recurrence of colorectal adenomas. *N Engl J Med*, **342**, 1149-55.
- Sheila AB, Nicholas ED, Robert L, et al (2003). Dietary fibre in food and protection against colorectal cancer in the European prospective investigation into cancer and nutrition (EPIC): an observational study. *Lancet*, **361**, 1496-501.
- Stewart BW & Kleihues P (2003). World cancer report. Geneva, World Health Organization/Lyon, International Agency for Research on Cancer.
- Slattery ML, Berry TD, Potter J, et al (1997). Diet diversity, diet composition and risk of colon cancer (United States). *Cancer Causes and Control*, **8**, 872-82.
- Steinmetz KA, Kushi LH, Bostick RM, et al (1994). Vegetables, fruit and colon cancer in the Iwo Women's Health Study. *Am J Epidemiol*, **139**, 1-15.
- Sung JJ, Lau JY, Goh KL, et al (2005). Increasing incidence of colorectal cancer in Asia: implications for screening. *Lancet Oncol*, **6**, 871-6.
- Tarawneh M, Nimri O (2007). Cancer incidence in Jordan. Jordan National Cancer Registry.
- Timothy JK, Paul NA, Elizabeth AS, et al (2009) Cancer incidence in vegetarians: results from the European prospective investigation into cancer and nutrition (EPIC-Oxford). *Am J Clin Nutr*, **89**, S1620-6.
- Ulrich CM, Bigler J, Whitton JA, et al (2001). Epoxide hydrolase Tyr113His polymorphism is associated with elevated risk of colorectal polyps in the presence of smoking and high meat intake. *Cancer Epidemiol Biomarkers Prev*, **10**, 875-82.
- US department of Agriculture, US Department of Health and Human Services (2005). Nutrition and your Health: Dietary Guidelines for Americans. 6th ed. Home and Garden Bulletin no.232. Washington, DC: US government Printing Office.
- WCRF, World Cancer Research Fund/American Institute for Cancer Research (AICR) (2008). Food, Nutrition and the Prevention of Cancer; a Global Perspective. Washington, DC: American Institute for Cancer Research.
- Willett WC, Stampfer MJ, Colditz GA, et al (1990). Relation of meat, fat and fiber intake to the risk of colon cancer in a prospective study among women. *N Engl J Med*, **13**, 1664-72.
- WHO, World Health Organization (2006). Cancer, Geneva (Fact Sheet No. 297).
- WHO, World Health Report (2005). Make every mother and child count. Geneva, World Health Organization.
- Wu AH, Paganin-Hill A, Ross RK, et al (1987). Alcohol, physical activity, and other risk factors for colorectal cancer: a prospective study. *Br J Cancer*, **55**, 687-94.