

RESEARCH ARTICLE

Cytotoxicity, Toxicity, and Anticancer Activity of *Zingiber Officinale* Roscoe Against Cholangiocarcinoma

Tullayakorn Plengsuriyakarn¹, Vithoon Viyanant¹, Veerachai Eursitthichai¹, Smarn Tesana², Wanna Chaijaroenkul¹, Arunporn Itharat³, Kesara Na-Bangchang^{1*}

Abstract

Cholangiocarcinoma (CCA) is an uncommon adenocarcinoma which arises from the epithelial cells of the bile ducts. The aim of the study was to investigate the cytotoxicity, toxicity, and anticancer activity of a crude ethanolic extract of ginger (*Zingiber officinale* Roscoe) against CCA. Cytotoxic activity against a CCA cell line (CL-6) was assessed by calcein-AM and Hoechst 33342 assays and anti-oxidant activity was evaluated using the DPPH assay. Investigation of apoptotic activity was performed by DNA fragmentation assay and induction of genes that may be involved in the resistance of CCA to anticancer drugs (MDR1, MRP1, MRP2, and MRP3) was examined by real-time PCR. To investigate anti-CCA activity *in vivo*, a total of 80 OV and nitrosamine (OV/DMN)-induced CCA hamsters were fed with the ginger extract at doses of 1000, 3000, and 5000 mg/kg body weight daily or every alternate day for 30 days. Control groups consisting of 10 hamsters for each group were fed with 5-fluorouracil (positive control) or distilled water (untreated control). Median IC₅₀ (concentration that inhibits cell growth by 50%) values for cytotoxicity and anti-oxidant activities of the crude ethanolic extract of ginger were 10.95, 53.15, and 27.86 µg/ml, respectively. More than ten DNA fragments were visualized and up to 7-9 fold up-regulation of MDR1 and MRP3 genes was observed following exposure to the ethanolic extract of ginger. Acute and subacute toxicity tests indicated absence of any significant toxicity at the maximum dose of 5,000 mg/kg body weight given by intragastric gavage. The survival time and survival rate of the CCA-bearing hamsters were significantly prolonged compared to the control group (median of 54 vs 17 weeks). Results from these *in vitro* and *in vivo* studies thus indicate promising anticancer activity of the crude ethanolic extract of ginger against CCA with the absence of any significant toxicity. Moreover, MDR1 and MRP3 may be involved in conferring resistance of CCA to the ginger extract.

Keywords: Cholangiocarcinoma - cytotoxicity - ginger - *Zingiber Officinale* roscoe

Asian Pacific J Cancer Prev, 13 (9), 4597-4606

Introduction

Cholangiocarcinoma (CCA) is a devastating cancer with increasing worldwide incidence and mortality rates. It is an uncommon adenocarcinoma which arises from the epithelial cells of the bile ducts anywhere along the intrahepatic and extrahepatic biliary tree excluding the papilla of Vater and the gall bladder (Mosconia et al., 2009). *Opisthorchis viverrini* (OV) infection is a high risk factor of CCA (Haswell-Elkins et al., 1994). The highest incidence rate in the world is observed in the northeast region of Thailand, where the prevalence of infection with OV is also highest (Haswell-Elkins et al., 1994; Sriamporn et al., 2004). The challenges posed by this often lethal biliary tract cancer are daunting, with conventional treatment options being limited and the only hope for long-term survival being that of complete surgical resection of the tumor. Chemotherapeutics for CCA is largely

ineffective and clinical efficacy of the standard treatment with 5-fluorouracil (5-FU) is low. Furthermore, resistance of this type of cancer to chemotherapy and radiotherapy is a major problem (Hejna et al., 1998).

Chemotherapy with plant-derived compounds or dietary phytochemicals has emerged as an accessible and promising approach to cancer control and management (Surh, 2003). A growing trend among some cancer patients is to combine conventional therapy with some form of complementary therapy (Vapiwala et al., 2006). Ginger is a food plant known worldwide and is equally reputed for its medicinal properties (Shukla and Singh, 2007). It is a herbaceous, rhizomatous perennial plant widely distributed throughout the tropical and subtropical regions. The rhizome of *Zingiber officinale* Roscoe is widely used as a dietary condiment throughout the world. Besides its extensive utilization as a spice, ginger has also been used in traditional oriental medicine to

¹Thailand Excellence Center in Drug Discovery and Development, ³Applied Thai Traditional Medicine Center, Faculty of Medicine, Thammasat University, Pathumthani, ²Department of Parasitology, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand
*For correspondence: kesaratmu@yahoo.com

ameliorate such symptoms as inflammation, rheumatic disorders, gastrointestinal discomforts, loss of appetite, travel sickness, hypercholesterolemia, and high level of triglyceride (Phuphathanaphong, 1979; Chrubasik et al., 2005). These diverse pharmacological activities of their major principles have already been confirmed (Surh et al., 2002). Several lines of evidence suggest that 6-gingerol is effective in the suppression of the transformation, hyperproliferation, and inflammatory processes that initiate and promote carcinogenesis, as well as the later steps of carcinogenesis, the angiogenesis and metastasis (Suzuki et al., 1997; Bode et al., 2001; Kim et al., 2005a; 2005b; Lee et al., 2008). It is regarded as a promising chemopreventive dietary agent exhibiting inhibition of cyclooxygenase and lipoxygenase activities (Kiuchi et al., 1982; 1992; Huang et al., 1991), apoptosis induction (Lee et al., 1998; Chauhan, 2002), and anti-tumorigenic effects (Park et al., 1998; Surh et al., 1999). The pungent vallinoids of ginger, [6]-gingerol and [6]-paradol, exhibit antiproliferation activity in liver, pancreatic, prostate, gastric, and leukemia cancer cells (Lee et al., 1998; Chen et al., 2007; Shukla and Singh, 2007). Furthermore, [6]-shogaol has also been shown to exhibit anticancer activities against breast cancer through the inhibition of cell invasion reduction of matrix metalloproteinase-9 expression (via blockade of nuclear factor activation) (Ling et al., 2010), anti-proliferation activity (through disruption of microtubule network of non-small lung epithelium cancer) (Choudhury et al., 2010), and anti-invasion on human hepatocellular cell (Weng et al. 2010). To our knowledge, there has been no report on the anticancer activity of ginger against CCA. Our previous study has demonstrated a promising cytotoxic activity of the ethanolic extract of ginger against CL-6 (CCA cell line obtained from human), HepG2 (hepatocarcinoma), and Hep-2 (laryngeal carcinoma) cell lines *in vitro* with IC₅₀ (concentration that inhibits cell growth by 50%) of less than 50 µg/ml (Mahavorasirikul et al., 2010).

The aim of the present study was to further investigate the cytotoxic activity of the crude ethanolic extract of ginger in other *in vitro* models (calcein-AM release and Hoechst 33342 assays), as well as its anti-oxidant activity, apoptotic activity, and activity on inducing the expression of multidrug resistance genes. Finally, the *in vivo* anticancer activity and toxicity of the crude extract was evaluated in OV/dimethylnitrosamine (DMN)-induced CCA in a hamster model.

Materials and Methods

Chemicals and reagents

Commercial grade ethanol was purchased from Labscan Co. Ltd. The cell culture medium Ham-F12, fetal bovine serum (FBS), L-glutamine, dimethylsulfoxide (DMSO), and the antibiotics streptomycin and penicillin were purchased from Gibco BRL Life Technologies (Grand Island, NY, USA). Renal epithelium cell growth medium and SupplementPack were purchased from Promocell Co. Ltd. (Germany). 5-fluorouracil (5-FU), DPPH (2,2-diphenyl-2-picrylhydrazyl), L-ascorbic acid (vitamin C), dimethylnitrosamine (DMN), and Tween-80

were purchased from Sigma-Aldrich Inc. (St. Louis, MO, USA).

Preparation of plant extract

Rhizomes of *Zingiber officinale* Roscoe (voucher number SKP206261501) were obtained from the Applied Thai Traditional Medicine Center, Faculty of Medicine, Thammasat University, Thailand. Preparation of the ethanolic extract of ginger was performed according to the previously described method (Mahavorasirikul et al., 2010). The extract was standardized using high-performance liquid chromatography to examine the amount of 6-gingerol which was 243±1 mg/g of dried rhizome. Chromatographic separation condition used was as follows: Phenomenex™ Luna 5 µm C18 column; mobile phase: a mixture of water and acetonitrile with gradient elution 0 min (55:45), 8 min (50:50), 17 min (35:65), 32 min (0: 100), and 43 min (55:45) at follow rate of 1 ml/min (Figure 1).

Cell lines and culture

The CCA cell line CL-6, was used for the *in vitro* assessment of cytotoxicity (calcein-AM release and Hoechst 33342 assays), anti-oxidant, apoptotic activity, and inducing effect on resistance gene expression of the ethanolic extract of ginger. CL-6 cell line was established and kindly provided by Associate Professor Dr. Adisak Wongkajornsilp of the Department of Pharmacology, Faculty of Medicine (Siriraj Hospital), Mahidol University, and were cultured in Ham-F12 medium supplemented with 10% heated fetal bovine serum and 100 IU/ml of anti-anti. Assessment of the cytotoxicity of ginger extract against CL-6 cell line was performed in comparison with HepG2 (hepatocarcinoma) and HRE (normal human renal epithelium) cell lines. HepG2 cell line was purchased from the Cell Line Service Co. Ltd. (Germany) and was cultured in a complete RPMI medium supplemented with 10% fetal bovine serum and 100 IU/ml pen-strep. HRE cell line was purchased from Promocell Co. Ltd. (Germany) and was cultured in renal epithelial cell growth medium 2 with SupplementPack. All cells were maintained at 37°C in a 5% CO₂ atmosphere with 95% humidity.

In vitro models for assessing cytotoxic, anti-oxidant, and apoptotic activities

Calcein-AM release assay: CL-6, HepG2, and HRE cells were plated in 96-well culture plates (1 ×10⁴ cells/well). After 24 hours of incubation, cells were incubated with various concentrations of ethanolic extract of ginger (1.95, 3.90, 7.81, 15.62, 31.25, 62.5, 125, and 250 µg/ml) at 37°C for 24 hours. 5-FU (at concentrations of 3.90, 7.81, 15.62, 31.25, 62.5, 125, 250, and 500 µg/ml) was used as positive control drug. Cells were then resuspended in complete medium and washed three times with PBS and incubated with 15 µM calcein-AM at 37°C for 30 min with occasional shaking. Calcein cellular fluorescence was read directly using a plate reader machine at excitation and emission wavelengths of 490 and 530 nm, respectively.

Hoechst 33342 assay: Inhibition of proliferation of CL-6, HepG2, and HRE cells by the ethanolic extract of ginger was measured by Hoechst 33342 assay (Schoonen

et al., 2005) using the same seeding cells and concentration ranges as that used in calcein-AM release assay. Wells of H342 plates were washed twice with 250 μ l PBS and tapped dry on a tissue paper. Plates were wrapped up and stored at -20 °C for DNA quantification with Hoechst 33342. An amount of 100 μ l of 0.01% SDS solution was added to each well. The plates were shaken for 30 min and then refrozen at -70 °C. The plates were thawed and 100 μ l of a H342 solution (2 μ g/ml Hoechst 33342, 10 mM Tris-HCl pH 7.4, 1 mM EDTA, 1 M NaCl) were added and plates were shaken at 37°C for 1 hour (in the dark). The fluorescence intensity was monitored in a plate reader machine at excitation and emission wavelengths of 355 and 460 nm, respectively.

Anti-oxidant activity: The anti-oxidant activity of the ethanolic extract of ginger was determined by measuring radical-scavenging activity of DPPH (2,2-diphenyl-2-picrylhydrazyl) (Szabo et al., 2007). Vitamin C (ascorbic acid) was used as a positive control reagent. The extract and vitamin C (at concentrations 1.95, 3.90, 7.81, 15.62, 31.25, 62.5, 125, and 250 μ g/ml) were added into each well of a 96-wells plate (100 μ l each). DPPH solution (0.1 ml DPPH with 3.9 ml methanol) at final concentration of 6×10^{-5} M was then added. The decrease in absorbance compared with the control well was determined at a UV wavelength of 515 nm.

For all of the above mentioned assays, results were generated from three independent experiments, triplicate each. Percentage of inhibition of the activity (cytotoxic, anti-proliferation, and anti-oxidant) was calculated as follows: % Inhibition = $[(\text{Absorbance}_{\text{control}} - \text{Absorbance}_{\text{test}}) / \text{Absorbance}_{\text{control}}] \times 100$

The IC₅₀ (concentration that inhibits the activity by 50%) values were calculated using CalcuSyn™ software (Biosoft, UK).

Apoptosis assay

The apoptotic activity of the crude ethanolic extract of ginger was determined by electrophoresis of DNA (Hsu et al., 2008). CL-6 cells (1×10^6) were treated with various concentrations of the ethanolic extract of ginger (6.25, 12.5, 25, 50, and 100 μ g/ml) at 37°C for 48 hours and collected by trypsinization and centrifugation. Pellets were washed twice with PBS and lysed by DNA lysis buffer (Tris EDTA, pH 8, 5 M NaCl, and 0.5 M EDTA). Following centrifugation, supernatant was incubated overnight with proteinase K (0.1 mg/ml) and RNase (0.2 mg/ml) at 37°C for 2 hours. DNA was collected by precipitation with two volumes of isopropanol in the presence of 3 M sodium acetate. After centrifugation, the DNA pellets were washed overnight with 70% ethanol and air-dried. DNA was separated on 1.8% agarose gel containing 1 mg/ml ethidium bromide and DNA fragmentation was visualized under a UV lamp. Negative control wells consisted of untreated and cells similarly treated with 50% ethanol. 100 bp plus DNA ladder was used as a marker.

Induction of the expression of multidrug resistant genes

CL-6, HepG2, and HRE cells (1×10^6) were treated with the ethanolic extract of ginger at concentrations of

25, 50, and 100 μ g/ μ l for 24 and 48 hours. Total RNA was extracted by RNeasy Mini Kit (Qiagen, Hilden, Germany) and quantified spectrophotometrically by Nanodrop machine (Thermo Scientific, Wilmington, USA). First-strand cDNA was synthesized from 100 ng total RNA by reverse transcription using oligo-dT primers and reverse transcriptase (Superscript III; Invitrogen, USA) according to the manufacturer's instructions. Relative quantitation of gene expression was measured by real-time PCR. Five sets of primers were used in all reactions to obtain amplification of housekeeping genes control, GAPDH, and a specific target gene of interest, multidrug resistance 1 (MDR1), multidrug resistance protein 1 (MRP1), multidrug resistance protein 2 (MRP2), and multidrug resistance protein 3 (MRP3) genes. Following an initial denaturation step at 94°C for 5 minutes, 35 cycles of PCR amplification were performed, with each consisting of a denaturation step at 94°C for 30 seconds, annealing at 62°C for 45 seconds, and extension at 72°C for 1 min. At the end of the 35 cycles, a 5-minute extension phase at 72°C was included to provide complete synthesis. Primers were obtained from Eurofins MWG Operon (Huntsville, ALB, USA). The PCR primers used to amplify MDR1, MRP1, MRP2, and MRP3 genes were 20/20, 23/22, 19/20, and 21/22 nucleotide long oligonucleotides, respectively. The MDR1 sequences of sense and antisense strand primers were 5'GTCTTTGGTGCCATGGCCGT and 5'ATGTCCGGTCGGGTGGGATA, respectively. The MRP1 sequences of sense and antisense strand primers were 5'CTGACAAGCTAGACCATGAATGT and 5'CCTTTGTCCAAGACGATCACCC, respectively. For MRP2 and MRP3, sequences of sense vs. antisense strand primers were 5'GCCAGATTGGCCAGCAA vs. 5'AATCTGACCACCGGCAGCCT and 5'GGGACCCTGCGCATGAACCTG vs. 5'TAGGCAAGTCCAGCATCTCTGG, respectively. GAPDH, 22/21 long oligonucleotides, was an internal standard for mRNA expression. The sequences of sense and antisense strand primers were 5'CAACAGCCTCAAGATCATCAGC and 5'TTCTAGACGGCAGGTCAGGTC, respectively. The fluorescence threshold (Ct) was calculated from $2^{-\Delta\Delta C_T}$. The absence of non-specific products was confirmed by the analysis of the melting-point curves.

In vivo model for evaluation of toxicity and anticancer activity

Animals: Syrian golden hamsters, 6-8 weeks of age, weighting 105-120 g used in all experiments were purchased from The National Laboratory Animal Centre of Thailand (Thamavit et al., 1978). They were housed under standard conditions and fed with a stock diet and water *ad libitum*. Approval of the study protocol was obtained from the Ethics Committee for Research in Animals, Thammasat University, Thailand.

Acute and subacute toxicity study: Acute and subacute toxicity tests were performed according to the OECD guideline for chemicals (OECD, 2004). Sixty hamsters (5 males and 5 females for each group) were fed (via gastric gavage) with three dose levels of ethanolic extract of ginger (resuspended in a mixture of distilled water

and Tween-80, 4:1, v:v), i.e., 1000, 3000, and 5000 mg/kg body weight. The control hamsters were fed with the mixture of distilled water and Tween-80. Animals were closely observed for awareness, status of mood, motor activity, CNS excitation, posture, muscle tone, reflexes, and autonomic signs during the first 30 minutes, periodically during the first 24 hours, and then daily for 14 days (acute toxicity) or 30 days (subacute toxicity) (OECD 2004). For subacute toxicity test, body weight, and food and water consumption were recorded daily for 30 days. At the end of the observational period, all animals were sacrificed under ether anesthesia and vital organs (brain, heart, kidneys, liver, spleen, stomach, large and small intestine, and lungs) were removed from all animals.

OV-DMN induced CCA in hamster model: Assessment of anticancer activity of the crude ethanolic extract of ginger against CCA was performed in ninety hamsters (45 males and 45 females) (Somchai et al., 2004). The metacercariae of *O. viverrini* were collected as the naturally infected cyprinoids fish captured from an endemic area of Khon Kaen, northeast Thailand. The parasite species were confirmed under light microscope (Boonmars et al., 2009) and were minced and digested with pepsin-HCl, then filtrated and washed with normal saline.

Animals were divided into nine groups (5 males and 5 females each). The first eight groups were treated as OV-infected groups. Development of CCA was induced by initial feeding of all animals (by gastric gavage) with 50 metacercariae of OV, followed four weeks later by drinking water containing 12.5 ppm of dimethylnitrosamine (DMN) for eight weeks (Tesana et al., 2007). 5-FU was used as a positive control treatment (group 1) and the groups consisting of healthy hamsters (group 9) and OV/DMN-induced hamsters without any treatment (group 2) were served as normal control and untreated control groups, respectively. The occurrence and development of CCA was detected and confirmed by ultrasonography throughout the investigation period and finally by histopathology at autopsy. **Group 1 (5-FU treated, positive control):** OV/DMN induced CCA hamsters, treated with 5-FU (40 µg/kg body weight, single intravenous injection). **Group 2 (negative control):** OV/DMN-induced CCA hamsters treated with vehicle (a mixture of distilled water and Tween-80) daily for 30 days. **Group 3 (high dose-1):** OV/DMN-induced CCA hamsters treated with 5,000 mg/kg body weight crude ethanolic extract of ginger daily for 30 days. **Group 4 (high dose-2):** OV/DMN-induced CCA hamsters, treated with 5,000 mg/kg body weight crude ethanolic extract of ginger every alternate day for 30 days. **Group 5 (medium dose-1):** OV/DMN-induced CCA hamsters, treated with 3,000 mg/kg body weight crude ethanolic extract of ginger daily for 30 days. **Group 6 (medium dose-2):** OV/DMN-induced CCA hamsters, treated with 3,000 mg/kg body weight crude ethanolic extract of ginger every alternate day for 30 days. **Group 7 (low dose-1):** OV/DMN-induced CCA hamsters, treated with 1,000 mg/kg body weight crude ethanolic extract of ginger daily for 30 days. **Group 8 (low dose-2):** OV/DMN-induced CCA hamsters, treated with 1,000 mg/kg body weight crude ethanolic extract of




Figure 1. Standardization using High-performance Liquid Chromatography of 6-gingerol. Chromatographic separation condition used was as follows: Phenomenex™ Luna 5 µm C18 column; mobile phase: a mixture of water and acetonitrile with gradient elution 0 min (55:45), 8 min (50:50), 17 min (35:65), 32 min (0: 100), and 43 min (55:45) at follow rate of 1 ml/min.




Figure 2. Schematic Protocol for the Evaluation of Anticancer Activity of the Ethanolic extract of Ginger against CCA in an OV/DMN induced CCA Hamster Model. OV, infection with 50 *Opisthorchis viverrini* metacercariae; DMN start, administration of carcinogen; DMN stop, withdrawal; Treated, administration of treatment (ethanolic extract of ginger or 5-FU), Died, time of start of death of hamsters

ginger every alternate day for 30 days **Group 9 (normal control):** healthy (non-CCA induced) hamsters treated with vehicle (a mixture of distilled water and Tween-80) daily for 30 days.

In groups 3-8, the crude extract of ginger was fed to the animals at 12 weeks after induction with OV metacercariae. Schematic diagram summarizing the experimental design is shown in Figure 2. Body weight and food and water consumption were recorded daily for 30 days. At autopsy, livers and bile ducts were removed from all animals.

Survival time and survival rate were the primary endpoint parameters for the evaluation of the anticancer activity of the crude ethanolic extract of ginger against CCA.

Autopsy and histopathology: For both toxicity and anticancer activity evaluation, all organs were removed at autopsy and observed macroscopically. Samples were




Figure 3. DNA Fragmentation of the Ethanolic Extract of Ginger-treated CL-6 cells. DNA was extracted from untreated control or after treatment with 50% ethanol (negative control) or various concentration of the ethanolic extract of ginger (100, 50, and 25 µg/ml) for 48 h DNA was electrophoresed in 1.8% agarose and then stained with ethidium bromide. Marker represents the 100 bp plus DNA ladder.




Figure 4. Expression (gene copy number compared with baseline) of Multidrug Resistance Genes MDR1, MRP1, MRP2, and MRP3 Following 24 and 48 Hours Exposure of (a) CL-6, (b) Hep-G2, and (c) HRE cell lines, to the Ethanolic Extract of Ginger at Concentrations of 20, 50, and 100 µg/ml

fixed with 10% formalin solution. Specimens were washed in phosphate buffer three times, then dehydrated in an ascending series of ethanol for 15 min each and embedded in paraffin, followed by sectioning and staining with hematoxylin and eosin (Chaimuangraj et al., 2003).

Statistical analysis

Data are expressed as median (range) values. Significant difference between quantitative data of more than two data sets was performed by Kruskal-Wallis test. Significant difference between two quantitative data sets was performed by Mann-Whitney test. Statistical significance level was set at $\alpha=0.05$ for all tests.

Results

In vitro models for assessing cytotoxic, anti-oxidant, and apoptotic activities

Cytotoxic activity: The cytotoxic activity of the ethanolic extract of ginger was investigated against the human CCA cell line CL-6 based on calcein-AM and Hoechst 33342 assays, in comparison with HepG2 and HRE cell lines. In both assays, the extract was found to inhibit cell viability in a dose-dependent manner following 48 hours of exposure. The IC_{50} [median (range)] values of the extract in CL-6, HepG2, and HRE cell lines based on the calcein-AM and Hoechst assays are presented in Table 1. The extract exhibited about 1.5-6 times as potent as the positive control drug 5-FU against CL-6 cells in both assays. The cytotoxic activity against CL-6 cells was more specific when compared with HepG2 and HRE cells. The median values of selectivity index (SI) of the extract against CL-6, HepG2, and HRE cells were 18.09, 2.76, and 1 in the calcein-AM release assay, respectively. The corresponding SI values in the Hoechst 33342 assay were 4.63, 2.68, and 1, respectively.

Anti-oxidant activity: The *in vitro* DPPH assay demonstrated comparable anti-oxidant activity of the crude ethanolic extracts of ginger with ascorbic acid (positive control), with IC_{50} [median (range)] values of 27.86 (27.05-28.03) and 21.38 (21.25-21.49) µg/ml, respectively.

Apoptotic activity: The effect of the crude extract of ginger on the induction of apoptosis in CL-6 cells was assessed by DNA fragmentation assay. Agarose gel electrophoresis showed that treatment with ginger extract resulted in the formation of DNA fragments in CL-6 cells (Figure 3). More than ten DNA fragments were visualized and the activity was concentration- dependent as the proportion of apoptotic cells increased with higher

concentrations of the extract.

Induction of the expression of multidrug resistant genes

The inducing effect of different concentrations (20, 50, and 100 µg/ml) of crude ethanolic extract of ginger on the mRNA expression of different multidrug resistance genes (MDR1, MRP1, MRP2, and MRP3) following exposure to CL-6, HepG2, and HRE cells for 24 and 48 hours was investigated using real-time PCR. The expression of MDR1 gene following 24 and 48 hours exposure to the ethanolic extract clearly exhibited both time- and concentration-dependency. A quantitative analysis showed that the copy number of MDR1 and MRP3 genes were found to be 7-9 folds up-regulated following exposure to the extract compared with the non-exposed CL-6 cell at time zero (Figure 4). Compared with the non-exposed cells, the expression of MDR1 and MRP3 in HepG2 cells was 5-8 fold up-regulated following exposure to the extract, whereas for HRE cells, only MRP3 was found to be about 5 fold up-regulated

In vivo model for evaluation of toxicity and anticancer activity

Toxicity test: For the acute and subacute toxicity studies, single oral doses of crude ethanolic extract of ginger at all of the three levels (1000, 3000, and 5000 mg/kg body weight) did not cause mortality in any animal (0% mortality) during the investigation period. Only stomach

Table 1. *In vitro* Cytotoxic Activity [median (range) values] and Selectivity Index (SI) of the crude ethanolic extract compared with 5-FU

Cell line	Ethanolic extract of ginger	5-FU
CL-6;		
Calcein-AM:		
IC_{50} (µg/ml)	10.95 (10.87-11.12)	89.87 (89.57-90.84)
SI	18.09	3.19
Hoechst33342:		
IC_{50} (µg/ml)	53.13 (48.25-55.13)	95.29 (92.84-98.24)
SI	4.63	3.12
HepG2;		
Calcein-AM:		
IC_{50} (µg/ml)	71.89 (69.88-73.14)	74.86 (73.42-77.96)
SI	2.76	3.83
Hoechst33342:		
IC_{50} (µg/ml)	92.88 (87.15-94.26)	118.60(115.67-120.19)
SI	2.68	2.51
HRE;		
Calcein-AM:		
IC_{50} (µg/ml)	198.15 (196.99-205.67)	286.74(275.78-286.74)
SI	1	1
Hoechst33342:		
IC_{50} (µg/ml)	245.91 (234.87-250.17)	297.39(289.57-311.87)
SI	1	1




Figure 5. Liver Morphology and Histopathological Examination under Light Micrograph of Small Foci Cancer Stained with Hematoxylin and Eosin Staining of the Normal Control (5a and 5e), 5-FU Treated Hamster (5b and 5f), Untreated Hamster (5c and 5g), and High Dose Treated Hamster (5d and 5h).

Table 2. Survival Time [median (range) values] of 5-FU Treated OV/DMN-induced CCA Hamsters (group 1) and Treated OV/DMN Induced CCA Hamsters (groups 2-8)

Control hamsters		Treated hamsters					
		Daily regimen			Every alternate regimen		
5-FU treated ^a (group 1)	Untreated control ^b (group 2)	High dose (group 3)	Medium dose (group 5)	Low dose (group 7)	High dose (group 4)	Medium dose (group 6)	Low dose (group 8)
25.5 (22-28)	17 (16-20)	54 (52-57)	51.5 (49-53)	47 (44-49)	55 (53-58)	51 (50-55)	47 (44-49)

^aStatistically significant difference with Group 2 ($p < 0.001$), Group 3 ($p < 0.001$), Group 4 ($p < 0.001$), Group 5 ($p < 0.001$), Group 7 ($p < 0.001$), Group 2 ($p < 0.001$), Group 4 ($p < 0.001$), and Group 6 ($p < 0.001$). ^bStatistically significant difference with Group 3 ($p < 0.001$), Group 5 ($p < 0.001$), Group 7 ($p < 0.001$), Group 2 ($p < 0.001$), Group 4 ($p < 0.001$), and Group 6 ($p < 0.001$).

irritation was observed in all animals immediately after feeding them with the extract. The animals however, recovered from the symptom within one hour of dosing. The average daily intake of water and food, including the average body weight of animals were comparable in all groups. No abnormal histopathology was observed in any vital organ at autopsy.

Development of CCA: Compared with the normal control (group 9), the average body weights of the animals in groups 1-8 were significantly reduced after DMN withdrawal. Morphology and histology of normal and cancerous hamster are shown in Figure 5. Histopathological examination of all OV/DMN induced hamsters (groups 1-8), but not in the control group (group 9), confirmed adenocarcinoma and cholangiofibrosis.

Anticancer activity against CCA: Significant prolongation of survival time was observed in OV/DMN induced CCA hamsters which were treated with the ethanolic extract of ginger in all treatment regimens compared with both the 5-FU treated (group 1) and untreated control (group 2) group (Table 2). Animals started to die as early as 17 weeks in the untreated control group. Median (range) survival time of CCA-bearing hamsters following feeding with all the six treatment regimens was about 3- and 2- times of the untreated and 5-FU treated (positive control) groups, respectively. The extract when given at alternate days for 30 days was found to significantly prolong the survival time of animals compared with the daily dose regimens (Table 2). Survival rate at week 58 was 0% in all groups, except the group treated with the highest dose level of 5,000 mg/kg every alternate day for 30 days (group 6), of which 2/10 (20%) of animals still survived (20% survival rate at week 58).

Discussion

Plant-derived compounds are gaining increasing

interest as potential cancer therapeutics including for the treatment of refractory cancers such as CCA. The present study demonstrated the promising cytotoxic and anticancer activities of the crude ethanolic extract of ginger (rhizome of *Zingiber officinale* Roscoe) against CCA at tolerated dose levels. Results of toxicity tests suggest that the extract was well-tolerated when given by oral route even at a high dose of 5,000 mg/kg body weight to a total of sixty male and female hamsters. All animals survived during the investigation period with no overt sign of morbidity or abnormal locomotor activity. Only reversible gastrointestinal irritation occurred and hamsters recovered within one hour of drug administration.

Different *in vitro* cytotoxicity assays with different endpoints have been employed for screening of potential compounds or plant-derived extract preparation for their anticancer activities. The most commonly used assays involve the use of dye stains that include MTT, calcein-AM, and Hoechst 33342 assays. These dyes have some indicator properties allowing them to reveal ongoing cellular processes, providing indirect measure of mitochondria function (MTT), esterase activity (calcein-AM), and DNA binding (Hoechst 33342). In our previous study (Mahavorasirikul et al. 2010), the crude ethanolic extract of ginger was shown by MTT assay to exhibit cytotoxic activity against human CCA CL-6 cell line with IC_{50} of 34.26 μ g/ml. This activity was more selective to CL-6 cell compared with normal cell (HRE) with a selectivity index of 3.5. In the present study, we applied two cytotoxicity assays, i.e., calcein-AM release and Hoechst 33342 to further obtain additional support for the cytotoxicity of the ethanolic extract of ginger. The median IC_{50} values of the crude extract based on calcein-AM release and Hoechst 33342 assays were 10.95 and 53.15 μ g/ml, respectively. This promising cytotoxic activity of the crude ethanolic extract of ginger was also reported with HepG2 (IC_{50} 9.67 μ g/ml) and Hep2 (32.40 μ g/ml)

cell lines (Mahavorasirikul et al. 2010). The selectivity index (SI) of the extract for CL-6 cells in both assays were relatively high (18.09 and 4.63) compared with 5-FU (3.19 and 3.12). In addition, the extract was more selective to CL-6 cells compared with HepG2 cells (SI 2.76 and 2.68). In support of cytotoxic activity, our results also showed the apoptotic activity of the crude extract, a mechanism by which it induced cytotoxicity. A dose-dependent increase of DNA fragmentation (by DNA gel-electrophoresis) was observed in CL-6 cell line after exposure to the extract compared with control cells. DNA fragmentation and disintegrating apoptotic cells could be observed within 48 hours of exposure to 12.5 µg/ml of the extract. It is interesting to further elucidate the mechanisms of induced apoptosis by major components of ginger, e.g., the possible involvement of caspase enzymes.

Evidence is accumulating which indicates that many chemotherapeutic agents may be selectively toxic to tumor cells because they increase oxidant stress and enhance these already stressed cells beyond their limit (Moungjaroen et al., 2006). A potent scavenger of these free radical species may therefore serve as a possible prevention intervention for free radical mediated cancer (Ames et al., 1995). The efficacies of various anti-oxidants have been associated with their ability to scavenge free radicals (Wang et al., 1999). In the present study, the crude ethanolic extract of ginger was found to exhibit radical scavenging activity with potency comparable to that of the standard compound ascorbic acid with median IC₅₀ of 27.86 and 21.38 µg/ml, respectively.

Resistance of cancerous cells to chemotherapeutic drugs (multidrug resistance) is a major cause of the failure of cancer chemotherapy (Lonning, 2003). The drug resistant phenomenon seems particularly more obvious with CCA (Liu et al., 2010). Multiple mechanisms have been hypothesized to play a role in chemotherapeutic drug resistance in cancers and the most important ones are associated with the over-expression of various members of ATP-binding cassette (ABC) MDR1 and MRPs (Gottesman et al., 2002), an increase in detoxification of chemotherapeutic drugs (e.g. glutathione S-transferases: GST) and dihydropyrimidine dehydrogenase: DPD) (Tew, 1994; Nita et al., 1998), as well as an alteration of drug targets and suppression of drug-induced apoptosis (Stavrovskaya, 2000). Several lines of evidence suggest that both MDR1 and MRP1 are the major contributors of the multidrug resistance phenotypes observed in a number of tumor cells (Ambudkar et al., 2003; Larkin et al., 2004). Many compounds have been investigated for their ability to inhibit the efflux protein function, thus leading to the development of several generations of MDR modulators (Lee, 2004). To assess the mechanism by which the crude ethanolic extract of ginger may modulate cellular efflux in CL-6 cells, we evaluated the expression of the efflux transporters MDR1 and MRPs (MRP1, MRP2, and MRP3), which are the membrane pump proteins that were shown to be expressed in cholangiocytes (Cao et al., 1998; Courtois et al., 1999; Gigliozzi et al., 2000; Rost et al., 2001). The extract was found to induce mRNA expression of these genes at varying potencies and patterns of time- and concentration-dependency. The expression level was

found relatively higher with MDR1 and MRP3 and tended to be time- and concentration-dependent. The copy number of cDNA was increased with prolonged exposure time and higher concentrations of the extract. Expression of MRP1 was unchanged following exposure of the cell for 24 hours to the extract at concentrations of 20 and 50 µg/ml, but the induction of expression occurred at 50 µg/ml when the exposure time was increased to 48 hours. On the other hand, the mRNA expression of MRP2 was induced with 3 and 5 copies of genes expression when CL-6 cells were exposed to the extract at concentrations of 50 and 100 µg/ml. Interestingly, the expression was more or less stable when the exposure time was prolonged to 48 hours. MRP2 is one of the expressed efflux protein on the apical membrane of hepatocytes and cholangiocytes. It plays an important role in the biliary clearance of endogenous and exogenous toxic compounds. Association between a common MRP2 variant and CCA risk was reported (Hoblinger et al., 2009). In a previous study, MRP1 and MRP3 have been reported to be highly expressed in the five human CCA cell lines, while MRP2 was only moderately expressed and MDR1 expression was detected only in one cell line. A strong correlation was also found between the level of MRP3 expression and the IC₅₀ values of the anticancer drugs etoposide, doxorubicin, and pirarubicin (Tepsiri et al., 2005). Consistent with this finding, a strong association between MRP3 mRNA level and response to doxorubicin has been reported in lung cancer (Young et al., 1999). The role of MDR1 and MRP3 in chemotherapeutic resistance in CCA patients needs to be further investigated. Our data provide evidence that alteration in the expression or function of drug efflux pathways as a mechanism by which the ethanolic extract of ginger, may modulate the response to chemotherapy. Increased expression of these membrane pumps enhances drug efflux and may be associated with chemo-resistance, and subsequently, with poor clinical response. While recognizing the existence of numerous constituents of a ginger extract, further investigation on the MDR1 and MRP3 gene inducing potential of [6]-gingerol, [6]-shogaol, and [6]-paradol is required.

Several *in vivo* models have been described for assessing the anticancer activity of candidate compounds or extracts from natural products against CCA. These include subcutaneous xenograft model (Mareinfeld et al., 2003; Fava et al., 2005; Jimeno et al., 2005) or hepatobiliary CCA model (Johnson et al., 2001; Voskoglou-Nomokos et al., 2003; Bibby, 2004; Sausville and Burger, 2006) in hamsters or rats after treatment with carcinogens [N-nitrosobis (2-oxopropyl) amine, methylazoxymethyl acetate, dimethylnitrosamine, furan, thioacetamide] or infection with *O. viverrini* (OV) (Maronpot et al., 1991; Imray et al., 1992; Thamavit et al., 1993; Iki et al., 1998; Jan et al., 2007; Tesana et al., 2007), and genetic CCA models (Lai et al., 2005; Xu et al., 2006; Sirica et al., 2008). In our study, OV/DMN induced CCA in hamster model was used. Metacercariae of OV was fed (gastric gavage) to animals four weeks before DMN. DMN was given in drinking water daily for eight weeks before the start of treatment with the ethanolic extract of ginger at different dose schedules (groups 3-8) and standard drug

5-FU (group 1). Significant weight loss was observed in all groups induced with OV/DMN compared with the normal control (group 9). The crude ethanolic extract of ginger at all dose regimens markedly prolonged survival time and survival rate of the CCA-bearing hamsters compared with the untreated control group or even with 5-FU treated (positive control) group. The extract when given at highest dose of 5,000 mg/kg body weight daily for 30 days resulted in prolongation of survival time (median of 54 weeks) which was about 2- and 3- times of 5-FU (median of 25.5 weeks) and untreated (median of 17 weeks) control groups, respectively. When the extract was given at the same dose levels (1000, 3000, and 5000 mg/kg) but at every alternate day, survival time appeared to be significantly more prolonged (about 2- and 3- times of the 5-FU treated and untreated control groups). A number of plant-derived compounds have been investigated for anticancer activity against CCA, notably triptolide from *Tripterygium wilfordii* (Tengchaisri et al., 1998) and the ubiquitous tannic acid (Naus et al., 2007). Anticancer activity of triptolide against CCA growth *in vitro* was reported with IC₅₀ as low as 0.05 µg/ml. In addition, the compound showed a significant reduction of tumor growth when given at a total dose of 1.2 mg. Tannic acid, a plant-derived polyphenol was also shown to exert anticancer activity against CCA cells *in vitro* and *in vivo*. These effects include growth inhibition by blocking cell-cycle progression *in vitro* and decreased growth of xenografts in nude athymic mice. With regards to the anticancer activity of gingerol, [6]-gingerol, reputedly the most active ginger constituent, has only been evaluated for its effect on various stages of carcinogenesis, whereas [6]-paradol has been demonstrated for antiproliferation activity in liver, pancreatic, prostate, gastric and leukemia cancer cells, and [6]-shogaol (dehydrated [6]-gingerol) for anticancer activity against breast cancer (Pereira et al., 2011).

In conclusion, results from the present study suggest that the crude ethanolic extract of ginger exhibited *in vitro* cytotoxic, anti-oxidant, and apoptotic activities. In addition, *in vivo* study showed promising anticancer activity of the extract against CCA. Its remarkable tumor inhibition effect *in vivo* offers an alternative treatment for CCA when either used alone or in conjunction with other anticancer or immune-modulating agents. Nevertheless, the inducing activity of the extract on the expression of drug resistant genes is of major concern. In future the molecular and cellular mechanisms of action and resistance of the active principle [6]-gingerol, including [6]-shogaol and [6]-paradol in CCA should be further investigated.

Acknowledgements

The study was supported by the Commission for Higher Education, Ministry of Education of Thailand (RG) and the National Research University Project of Thailand (NRU), Office of Higher Education Commission of Thailand. Tullayakorn Plengsuriyakarn receives financial support for the PhD program from the Commission for Higher Education, Ministry of Education of Thailand.

References

- Ambudkar SV, Kimchi-Sarfaty C, Sauna ZE, Gottesman MM (2003). P-glycoprotein: from genomics to mechanism. *Oncogene*, **22**, 7468-85.
- Ames BN, Gold LS, Willett WC (1995). The causes and prevention of cancer. *Proc Nat Acad Sci*, **92**, 5258-65.
- Bibby MC (2004). Orthotopic models of cancer for preclinical drug evaluation: advantages and disadvantages. *Eur J Cancer*, **40**, 852-7.
- Bode AM, Ma WY, Surh YJ, Dong Z (2001). Inhibition of epidermal growth factor-induced cell transformation and activator protein 1 activation by [6]-gingerol. *Cancer Res*, **61**, 850-3.
- Boonmars T, Wu Z, Boonjaruspinyo S, et al (2009). Alterations of gene expression of RB pathway in *Opisthorchis viverrini* infection-induced cholangiocarcinoma. *Parasitol Res*, **105**, 1273-81.
- Cao L, Duchrow M, Windhovel U, et al (1998). Expression of MDR1 mRNA and encoding P-glycoprotein in archival formalin-fixed paraffin-embedded gall bladder cancer tissues. *Eur J Cancer*, **34**, 1612-7.
- Chaimuangraj S, Thamavit W, Tsuda H, Moore MA (2003). Experimental investigation of *Opisthorchiasis*-associated cholangiocarcinoma induction in the Syrian hamster - pointers for control of the human disease. *Asian Pac J Cancer Prev*, **4**, 87-93.
- Chauhan DP (2002). Chemotherapeutic potential of curcumin for colorectal cancer. *Curr Pharm Res*, **8**, 1695-706.
- Chen CY, Liu TZ, Liu YW, et al (2007). 6-shogaol (alkanone from ginger) induces apoptotic cell death of human hepatoma p53 mutant Mahlavu subline via an oxidative stress-mediated caspase-dependent mechanism. *J Agri Food Chem*, **55**, 948-54.
- Choudhury D, Das A, Bhattacharya A, Chakrabarti G (2010). Aqueous extract of ginger shows antiproliferative activity through disruption of microtubule network of cancer cells. *Food Chem Toxicol*, **48**, 2872-80.
- Chrubasik S, Pittler MH, Roufogalis BD (2005). *Zingiberis rhizoma*: a comprehensive review on the ginger effect and efficacy profiles. *Phytomed*, **12**, 684-701.
- Courtois A, Payen L, Lagadic D, Guillouzo A, Fardel O (1999). Evidence for a multidrug resistance-associated protein 1 (MRP1)-related transport system in cultured rat liver biliary epithelial cells. *Life Sci*, **64**, 763-74.
- Fava G, Marucci L, Glaser S, et al (2005). Gamma-Aminobutyric acid inhibits cholangiocarcinoma growth by cyclic AMP-dependent regulation of the protein kinase A/extracellular signal-regulated kinase 1/2 pathway. *Cancer Res*, **65**, 11437-46.
- Gigliozzi A, Fraioli F, Sundaram P, et al (2000). Molecular identification and functional characterization of Mdr1a in rat cholangiocytes. *Gastroenterol*, **119**, 1113-22.
- Gottesman MM, Fojo T, Bates SE (2002). Multidrug resistance in cancer: role of ATP-dependent transporters. *Nat Rev Cancer*, **2**, 48-58.
- Haswell-Elkins MR, Mairiang E, Mairiang P, et al (1994). Cross sectional study of *Opisthorchis viverrini* infection and cholangiocarcinoma in communities within a high-risk area in northeast Thailand. *Int J Cancer*, **59**, 505-9.
- Hejna M, Pruckmayer M, Radrer M (1998). The role of chemotherapy and radiation in the management of biliary cancer: a review of the literature. *Eur J Cancer*, **34**, 977-86.
- Hoblinger A, Grunhage F, Sauerbruch T, Lammert F (2009). Association of the c.3972C>T variant of the multidrug resistance-associated protein 2 gene (MRP2/ABCC2) with susceptibility to bile duct cancer. *Digest*, **80**, 36-9.

- Hsu YL, Kuo PL, Tzeng TF, et al (2008). Huang-lian-jie-du-tang, a traditional Chinese medicine prescription, induces cell-cycle arrest and apoptosis in human liver cancer cells *in vitro* and *in vivo*. *J Gastroenterol Hepatol*, **23**, 290-9.
- Huang MT, Lysz T, Ferraro T, et al (1991). Inhibitory effects of curcumin on *in vitro* lipoxygenase and cyclooxygenase activities in mouse epidermis. *Cancer Res*, **51**, 813-9.
- Iki K, Tsujiuchi T, Majima T, et al (1998). Increased telomerase activity in intrahepatic cholangiocellular carcinomas induced by N-nitrosobis(2-oxopropyl)amine in hamsters. *Cancer Lett*, **131**, 185-90.
- Imray CH, Newbold KM, Davis A, Lavelle-Jones M, Neoptolemos JP (1992). Induction of cholangiocarcinoma in the Golden Syrian hamster using methylazoxymethyl acetate. *Eur J Sur Oncol*, **18**, 373-8.
- Jan YY, Yeh TS, Yeh JN, Yang HR, Chen MF (2007). Expression of epidermal growth factor receptor, apomucins, matrix metalloproteinases, and p53 in rat and human cholangiocarcinoma: appraisal of an animal model of cholangiocarcinoma. *Ann Sur*, **240**, 89-94.
- Jimeno A, Viqueira BR, ONE AUTHOR, et al (2005). Epidermal growth factor receptor dynamics influences response to epidermal growth factor receptor targeted agents. *Cancer Res*, **65**, 3003-10.
- Johnson JJ, Decker S, Zaharevitz D, et al (2001). Relationships between drug activity in NCI preclinical *in vitro* and *in vivo* models and early clinical trials. *Br J Cancer*, **84**, 1424-31.
- Kim EC, Min JK, Kim TY, et al (2005a). [6]-Gingerol, a pungent ingredient of ginger, inhibits angiogenesis *in vitro* and *in vivo*. *Biochem Biophys Res Com*, **335**, 300-8.
- Kim SO, Kundu JK, Shin YK, et al (2005b). [6]-Gingerol inhibits COX-2 expression by blocking the activation of p38 MAP kinase and NF-kappaB in phorbol ester-stimulated mouse skin. *Oncogene*, **224**, 2558-67.
- Kiuchi F, Shibuya M, Sankawa U (1982) Inhibitors of prostaglandin biosynthesis from ginger. *Chem Pharm Bull (Tokyo)*, **30**, 754-7.
- Kiuchi F, Iwakami S, Shibuya M, Hanaoka F, Sankawa U (1992). Inhibition of prostaglandin and leukotriene biosynthesis by gingerols and diarylheptanoids. *Chem Pharm Bull*, **40**, 387-91.
- Lai GH, Zhang Z, Shen XN, et al (2005). ErbB-2/neu transformed rat cholangiocytes recapitulate key cellular and molecular features of human bile duct cancer. *Gastroenterol*, **129**, 2047-57.
- Larkin A, O'Driscoll L, Kennedy S, et al (2004). Investigation of MRP-1 protein and MDR-1 P-glycoprotein expression in invasive breast cancer: a prognostic study. *Int J Cancer*, **112**, 286-94.
- Lee CH (2004). Reversing agents for ATP-binding cassette (ABC) transporters: application in modulating multidrug resistance (MDR). *Curr Med Chem Anticancer Agents*, **4**, 43-52.
- Lee E, Park KK, Lee JM, et al (1998). Suppression of mouse skin tumor promotion and induction of apoptosis in HL-60 cells by *Alpinia oxyphylla* Miquel (Zingiberaceae). *Carcinogen*, **19**, 1377-81.
- Lee HS, Seo EY, Kang NE, Kim WK (2008). [6]-Gingerol inhibits metastasis of MDA-MB-231 human breast cancer cells. *J Nutr Biochem*, **19**, 313-9.
- Ling H, Yang H, Tan SH, Chui WK, Chew EH (2010). 6-Shogaol, an active constituent of ginger, inhibits breast cancer cell invasion by reducing matrix metalloproteinase-9 expression via blockade of nuclear factor- κ B activation. *Br J Pharm*, **161**, 1763-77.
- Liu ZH, He YP, Zhou Y, Zhang P, Qin H (2010). Establishment and identification of the human multi-drug resistant cholangiocarcinoma cell line QBC939/ADM. *Mol Biol Rep*, **5**, 121-30.
- Lonning PE (2003). Study of suboptimum treatment response: lessons from breast cancer. *Lancet Oncol*, **4**, 177-85.
- Mahavorasirikul W, Viyanant V, Chaijaroenkul W, Itharat A, Na-Bangchang K (2010). Cytotoxic activity of Thai medicinal plants against human cholangiocarcinoma, laryngeal and hepatocarcinoma cells *in vitro*. *BMC Com Altern Med*, **10**, 55-62.
- Maronpot RR, Giles HD, Dykes DJ, Irwin RD (1991). Furan-induced hepatic cholangiocarcinomas in Fischer 344 rats. *Toxicol Pathol*, **19**, 561-70.
- Mosconia S, Beretta GD, Labiancaa R, et al (2009). Cholangiocarcinoma. *Crit Rev Oncol Hematol*, **69**, 259-70.
- Moungjaroen J, Nimmannit U, Callery PS, et al (2006). Reactive oxygen species mediate caspase activation and apoptosis induced by lipoic acid in human lung epithelial cancer cells through Bcl-2 down-regulation. *J Pharmacol Exp Ther*, **319**, 1062-9.
- Naus PJ, Henson R, Bleeker G, et al (2007). Tannic acid synergizes the cytotoxicity of chemotherapeutic drugs in human cholangiocarcinoma by modulating drug efflux pathways. *J Hepatol*, **46**, 222-9.
- Nita ME, Tominaga O, Nagawa H, Tsuruo T, Muto T (1998). Dihydropyrimidine dehydrogenase but not thymidylate synthase expression is associated with resistance to 5-fluorouracil in colorectal cancer. *Hepatogastroenterol*, **45**, 2117-22.
- OECD (2004.) Acute oral toxicity, guideline 420, The OECD Guideline for Testing of Chemical, 1-14.
- Park KK, Chun KS, Lee JM, Lee SS, Surh YJ (1998.) Inhibitory effects of [6]-gingerol, a major pungent principle of ginger, on phorbol ester-induced inflammation, epidermal ornithine decarboxylase activity and skin tumor promotion in ICR mice. *Cancer Lett*, **129**, 139-44.
- Pereira MM, Haniadka R, Chacko PP, Palatty PL, Baliga MS (2011). *Zingiber officinale* Roscoe (ginger) as an adjuvant in cancer treatment: a review. *J BUON*, **16**(3), 414-424.
- Phuphathanaphong, L (1979). Thai medicinal plants. 2nd ed., Department of Forestry, Bangkok.
- Rost D, Konig J, Weiss G, et al (2001). Expression and localization of the multidrug resistance proteins MRP2 and MRP3 in human gallbladder epithelia. *Gastroenterol*, **121**, 1203-8.
- Sausville EA, Burger AM (2006). Contributions of human tumor xenografts to anticancer drug development. *Cancer Res*, **66**, 3351-4.
- Schoonen WG, de Roos JA, Westerink WM, Débiton E (2005). Cytotoxic effects of 110 reference compounds on HepG2 cells and for 60 compounds on HeLa, ECC-1 and CHO cells. II Mechanistic assays on NAD(P)H, ATP and DNA contents. *Toxicol In Vitro*, **19**, 491-503.
- Shukla Y, Singh M (2007). Cancer preventive properties of ginger: a brief review. *Food Chem Toxicol*, **45**, 683-90.
- Sirica AE, Zhang Z, Lai GH, et al (2008). A novel "patient-like" model of cholangiocarcinoma progression based on bile duct inoculation of tumorigenic rat cholangiocyte cell lines. *Hepatol*, **47**, 1178-90.
- Somchai P, Yusuke H, Ning M, Puangrat Y, Reiji S (2004). Mechanism of NO-mediated oxidative and nitrative DNA damage in hamsters infected with *Opisthorchis viverrini*: a model of inflammation-mediated carcinogenesis. *Nitric Oxide*, **11**, 175-83.
- Sriamporn S, Pisani P, Pipitgool V, Suwanrungruang K (2004). Prevalence of *Opisthorchis viverrini* infection and incidence of cholangiocarcinoma in Khon Kaen, Northeast Thailand. *Trop Med Int Hlth*, **9**, 588-94.

- Stavrovskaya AA (2000). Cellular mechanisms of multidrug resistance of tumor cells. *Biochem (Mosc)*, **65**, 95-106.
- Surh TJ (2003). Cancer chemoprevention with dietary phytochemicals. *Nat Rev Cancer*, **3**, 768-80.
- Surh YJ, Park KK, Chun KS, et al (1999). Anti-tumor-promoting activities of selected pungent phenolic substances present in ginger. *J Environ Pathol Toxicol Oncol*, **18**, 131-9.
- Surh YJ, Lee JY, Choi KJ, Ko SR (2002). Effects of selected ginsenosides on phorbol ester-induced expression of cyclooxygenase-2 and activation of NF-kappaB and ERK1/2 in mouse skin. *Ann New York Acad Sci*, **973**, 396-401.
- Suzuki F, Kobayashi M, Komatsu Y, Kato A, Pollard RB (1997). Keishi-ka-kei-to, a traditional Chinese herbal medicine, inhibits pulmonary metastasis of B16 melanoma. *Anticancer Res*, **17**, 873-8.
- Szabo MR, Iditoiu C, Chambre D, Lupea AX (2007). Improved DPPH determination for antioxidant activity spectrophotometric assay. *Chem*, **61**, 214-6.
- Tengchaisri T, Chawengkirtikul R, Rachaphaew N, et al (1998). Antitumor activity of TPL against cholangiocarcinoma growth *in vitro* and in hamsters. *Cancer Lett*, **133**, 169-75.
- Tepsiri N, Chaturat L, Sripa B, et al (2005). Drug sensitivity and drug resistance profiles of human intrahepatic cholangiocarcinoma cell lines. *World J Gastroenterol*, **11**, 2748-53.
- Tesana S, Takahashi Y, Sithithaworn P, et al (2007). Ultrastructural and immunohistochemical analysis of cholangiocarcinoma in immunized Syrian golden hamsters infected with *Opisthorchis viverrini* and administered with dimethylnitrosamine. *Parasitol Int*, **49**, 239-51.
- Tew KD (1994). Glutathione-associated enzymes in anticancer drug resistance. *Cancer Res*, **54**, 4313-20.
- Thamavit N, Bhamarapavati S, Sahaphong S, Vajrasthira T, Angsubhakorn S (1978). Effects of dimethylnitrosamine on induction of cholangiocarcinoma in *Opisthorchis viverrini* infected Syrian golden hamsters. *Cancer Res*, **38**, 4634-9.
- Thamavit W, Pairojkul C, Tiwawech D, et al (1993). Promotion of cholangiocarcinogenesis in the hamster liver by bile duct ligation after dimethylnitrosamine initiation. *Carcinogen*, **14**, 2415-7.
- Vapiwala N, Mick R, Hampshire MK, Metz JM, DeNittis AS (2006). Patient initiation of complementary and alternative medical therapies (CAM) following cancer diagnosis. *Cancer J*, **12**, 467-74.
- Voskoglou-Nomikos T, Pater JL, Seymour L (2003). Clinical predictive value of the *in vitro* cell line, human xenograft, and mouse allograft preclinical cancer models. *Clinical Cancer Res*, **9**, 4227-39.
- Wang H, Nair MG, Strasburg GM, et al (1999). Antioxidant and antiinflammatory activities of anthocyanins and their aglycon, cyanidin, from tart cherries. *J Nat Prod*, **62**, 294-6.
- Weng CH, Wu CF, Huang HW, Ho CT (2010). Anti-invasion effects of 6-shogaol and 6-gingerol, two active components in ginger, on human hepatocarcinoma cells. *Molecular Nutr Food Res*, **54**, 1618-27.
- Xu X, Kobayashi S, Qiao W, et al (2006). Induction of intrahepatic cholangiocellular carcinoma by liver-specific disruption of Smad4 and Pten in mice. *J Clin Invest*, **116**, 1843-52.
- Young LC, Campling BG, Voskoglou-Nomikos T, et al (1999). Expression of multidrug resistance protein-related genes in lung cancer: correlation with drug response. *Clin Cancer Res*, **5**, 673-80.