

RESEARCH ARTICLE

Is Sunlight a Predisposing Factor for Triple Negative Breast Cancer in Turkey?

Hasan Mutlu^{1*}, Abdullah Büyükçelik², Taner Çolak³, Mustafa Özdoğan⁴, Abdülşamet Erden⁵, Tuncay Aslan⁵, Zeki Akca⁶

Abstract

Intraduction: There is known to be a relationship between vitamin D level and more aggressive breast cancer subtypes, especially triple-negative breast cancer (TNBC). It was reported that sunlight exposure has an effect on the prognosis of patients with cancer, possibly related to the conversion of vitamin D to its active form with sunlight. We aimed to evaluate the effect of sunlight exposure on patients with TNBC. **Materials-Methods:** A total of 1,167 patients with breast cancer from two different regions of Turkey (Antalya and Kayseri, regions having different climate and sunlight exposure intensity over the year) were analysed retrospectively. The ratio of patients with TNBC was identified in those two regions. **Results:** The ratio of patients with TNBC was 8% and 12% for Kayseri and Antalya regions, respectively, and this difference between the two groups was statistically significant ($p=0.021$). **Discussion:** Sunlight exposure may be associated with more prevalent TNBC. This finding should be investigated with a prospective study.

Keywords: Breast cancer - sunlight exposure - season - triple negative

Asian Pacific J Cancer Prev, 14 (2), 801-803

Introduction

Breast cancer is the most common cancer type in women (Parkin et al., 1999). Triple-negative breast cancer (TNBC) is defined by absent expression of the estrogen receptor (ER), progesterone receptor (PR), and human epidermal growth factor receptor 2 (HER2). Approximately, TNBC accounts for 20% of all breast cancers (Swain, 2008) and the prognosis of patients with TNBC is usually worse than other types of breast cancer. Some factors such as parity, obesity, age at first pregnancy, BRCA mutations were reported as risk factors for TNBC (Trivers et al., 2009; Gonzalez-Angulo et al., 2011).

It was previously reported that sufficient sunlight exposure and vitamin D levels may decrease the morbidity and mortality due to breast cancer (Freedman et al., 2002; Abbas et al., 2008). The incidence and mortality of breast cancer were found to be lower in the regions with higher sunlight exposure (Mohr et al. 2008; Rhee et al., 2009).

The conversion of vitamin D to its active form was mediated via sunlight exposure in the skin. Vitamin D binds to and activates the vitamin D receptor (VDR). VDR acts as a transcription factor to modulate gene expression affecting cell cycle proliferation, cell cycle arrest, induction of differentiation, and activation of apoptosis (Peppone et al., 2012).

In this study, we aimed to evaluate whether more sunlight exposure has an effect on the ratio of patients with TNBC.

Materials and Methods

A total of 1167 patients with breast cancer from two different regions of Turkey (Antalya and Kayseri, those regions have different climate and sunlight exposure intensity for a year) were included in this study. The exposure hours to sunshine per day in those regions was given in Table 1. The patients were divided into two groups: Kayseri region which has a colder climate, and Antalya region with a warmer climate and more sunlight exposure than Kayseri region. Of patients 743 were from the Akdeniz University located in Antalya region and 424 patients were from the Kayseri Training and Research Hospital located in Kayseri were analysed retrospectively, using hospital records. The ratio of patients with TNBC was calculated. The age, stage, menopausal status, histological and nuclear-grade were evaluated in patients with TNBC. Statistical analysis was done using SPSS version 16.0 (Statistical Product and Service Solutions, SPSS Inc, Chicago, IL, USA). Frequencies analysis was carried out and independent two samples t-test and chi square test were used.

¹Department of Medical Oncology, Acibadem Kayseri Hospital, ²Department Internal Medicine, Kayseri Training And Research Hospital, Kayseri, ³Department of Internal Medicine, Acibadem University School of Medicine, Istanbul, ⁴Department of General Surgery, ⁵Department of Medical Oncology, Akdeniz University School of Medicine, Antalya, ⁶Department of Radiation Oncology, Zeki Akca, Mersin Government Hospital, Mersin, Turkey *For correspondence: doktorhasanmutlu@gmail.com

Table 1. Hours of Exposure to Sunshine Per Day in Kayseri and Antalya

Months	Hours of exposure to sunshine per day		P Value
	Kayseri	Antalya	
January	3	5	
February	4	7	
March	5	7	
April	6	9	
May	8	11	
June	10	12	
July	12	13	
August	11	12	
September	9	11	
October	7	9	
November	5	7	
December	3	5	
Day/Hours (mean)	6.9/207	9/270	0.097

Table 2. Characteristics of Patients with TNBC in Two Region

Variables		Kayseri (n:33)	Antalya (n:90)	P value
Age (mean)		55.7±9.8	49.0±10.1	0.001
Menopausal Status	Premenopausal	8 (24%)	46 (51%)	0.028
	Perimenopausal	0 (0%)	1 (1%)	
	Postmenopausal	25 (76%)	43 (48%)	
Stage	1	4 (12%)	23 (26%)	0.054
	2	17 (52%)	48 (53%)	
	3	10 (30%)	12 (13%)	
	4	0 (0%)	0 (0%)	
	Unkonwn	2 (6%)	7 (8%)	
Histological Grade	1	1 (3%)	0 (0%)	0.155
	2	8 (24%)	33 (36%)	
	3	9 (27%)	32 (36%)	
	Unknown	15 (46%)	25 (28%)	
Nuclear Grade	1	1 (3%)	10 (11%)	0.569
	2	11 (33%)	38 (42%)	
	3	6 (18%)	19 (21%)	
	Unknown	15 (46%)	23 (26%)	

Results

The patients characteristics are given in Table 2. The mean age of patients with TNBC in Kayseri and Antalya regions were 55,7±9,8 and 49,0±10,1 year, respectively (p=0.001). There was a significant difference in menopausal status between two regions (p=0.028). While the majority of patients were postmenopausal in Kayseri region, the premenopausal patients were more common in Antalya region. We did not find a difference in stage, histologic and nuclear grades (p=0.054, p=0.155, p=0.569, respectively) between two groups. The ratios of patients with TNBC in Kayseri and Antalya regions were 8% and 12% for, respectively and there was a statistically significant difference between two groups (p=0.021).

Discussion

In present study, we evaluated whether sunlight exposure has an impact on the ratio of TNBC. We find that the ratio of patients with TNBC was higher in the region with a more sunlight exposure. The effect of sunlight

exposure on the prognosis of patients with cancer is related to the conversion of vitamin D. Many studies reported that the adequate vitamin D level and more sunlight exposure decreased the incidence and mortality of breast cancer. In addition, it was reported that there was a relationship between low level of vitamin D and more aggressive breast cancer subtypes, especially TNBC (Rainville et al., 2009; Peppone et al., 2012; Yao et al., 2012). It was also reported that patients with TNBC had lower vitamin D level, and probably low level of vitamin D may be a characteristic of TNBC. The sunlight exposure is an important factor for the conversion of vitamin D to its active form, and in turn, the sufficient level of vitamin D have a positive effect on the prognosis of patients with cancer. Our results showed that the ratio with patients with TNBC were significantly higher in Antalya region with more sunlight exposure. However, in vitro studies showed that vitamin D decreases the estrogen and progesteron receptor expression on the breast cancer cells (Swami et al., 2000; 2003; Peng et al., 2007). Also, it was thought that the anti-proliferative effects of the vitamin D analogue, EB1089 on breast cancer cell line was through the down regulation of estrogen receptor (James et al., 1994; Davoodi et al., 1995). The potential activities of vitamin D on ErbB2 overexpressing have not been detailed in literature but it was previously reported that vitamin D analogue, BXL0124 inhibited the growth of ErbB2 overexpressing mammary tumors through regulating the ErbB2/AKT/ERK signaling pathways (Lee et al., 2010). We previously reported that the positivity of progesteron receptor was significantly lower among patients with breast cancer in Antalya region with more sunlight exposure than Kayseri region and there was a trend for the higher positivity of estrogen receptor (Mutlu et al., 2011). Due to the effect of vitamin D on estrogen and progesteron receptor during cancer development, the ratio of patients with TNBC might be higher in the region with more sunlight exposure despite the decrease of incidence and mortality of all patients with breast cancer. According to our results, while the patients with TNBC were mostly premenopausal status in Antalya with more sunlight exposure, the majority of patients with TNBC were postmenopausal in Kayseri. In previous studies, it was reported that the TNBC subtype was more prevalent among premenopausal (Carey et al., 2006; Lin et al., 2012). The more sunlight exposure via vitamin d decreased ER and PR expression may be contribute that TNBC is more commonly at young age.

It is not fully understood that the function of sunlight and vitamin D on the development of cancer. The further studies are warranted to evaluate the effect of sunlight and vitamin D on the development of cancer.

References

Abbas S, Linseisen J, Slanger T, et al (2008). Serum 25-hydroxy vitamin D and risk of postmenopausal breast cancer-results of a large case control study. *Carcinogenesis*, **29**, 93-9.
 Carey LA, Perou CM, Livasy CA, et al (2006). Race, breast cancer subtypes, and survival in the carolina breast cancer study. *JAMA*. **7**, 2492-502.
 Davoodi F, Brenner RV, Evans SRT, et al (1995). Modulation

- of vitamin D receptor and estrogen receptor by 1,25(OH)₂-vitamin D₃ in T47D human breast cancer cells. *J Steroid Biochem Mol Biol*, **54**, 147-53.
- Freedman D, Dosemeci M, McGlynn K (2002). Sunlight and mortality from breast, ovarian, colon, prostate, and non-melanoma skin cancer: a composite death certificate based case-control study. *Occup Environ Med*, **59**, 2567-2.
- Gonzalez-Angulo AM, Timms KM, Liu S, et al (2011). Incidence and outcome of BRCA mutations in unselected patients with triple receptor-negative breast cancer. *Clin Cancer Res*, **17**, 1082.
- James SY, Mackey AG, Binderup L, Colston KW (1994). Effects of a new synthetic vitamin D analogue, EB1089, on the oestrogen-responsive growth of human breast cancer cells. *J Endocrinol*, **141**, 555-63.
- Lee HJ, So JY, DeCastro A, et al (2010). Gemini vitamin D analog suppresses ErbB2-positive mammary tumor growth via inhibition of ErbB2/AKT/ERK signaling. *J Steroid Biochem Mol Biol*. **121**, 408-12.
- Lin NU, Vanderplas A, Hughes ME, et al (2012). Clinicopathologic features, patterns of recurrence, and survival among women with triple-negative breast cancer in the national comprehensive cancer network. *Cancer*, **15**, 5463-72
- Mohr SB, Garland CF, Gorham ED, et al (2008). Relationship between low ultraviolet B irradiance and higher breast cancer risk in 107 countries. *Breast J*, **14**, 255-60.
- Mutlu H, Colak T, Ozdoğan M, Altuner TY, Akça Z (2011). The effect of seasonal differences on prognostic factors in Turkish patients with breast cancer. *Eur J Cancer Prev*. **20**, 475-7.
- Parkin DM, Pisani P, Ferlay J (1999). Global cancer statistics. *CA Cancer J Clin*. **49**, 39.
- Peng X, Jhaveri P, Hussain-Hakimjee EA, Mehta RG (2007) Overexpression of ER and VDR is not sufficient to make ER-negative MDA-MB231 breastcancer cells responsive to 1alpha-hydroxyvitamin D₅. *Carcinogenesis*, **28**, 1000-7.
- Peppone LJ, Rickles AS, Janelins MC, Insalaco MR, Skinner KA (2012). The association between breast cancer prognostic indicators and serum 25-OH vitamin D levels. *Ann Surg Oncol*. **19**, 2590-9.
- Rainville C, Khan Y, Tisman G (2009). Triple negative breast cancer patients presenting with low serum vitamin D levels: a case series. *Cases J*, **21**, 8390.
- Rhee HV, Coebergh JW, Vries ED (2009). Sunlight, vitamin D and the prevention of cancer: a systematic review of epidemiological studies. *Eur J Cancer Prev*, **18**, 458-75.
- Swami S, Krishnan AV, Feldman D (2000). 1alpha,25-Dihydroxyvitamin D₃ downregulates estrogen receptor abundance and suppresses estrogen actions in MCF-7 human breast cancer cells. *Clin Cancer Res*, **6**, 3371-9.
- Swami S, Raghavachari N, Muller UR, Bao YP, Feldman D (2003). Vitamin D growth inhibition of breast cancer cells: gene expression patterns assessed by cDNA microarray. *Breast Cancer Res Treat*, **80**, 49-62.
- Swain S (2008). Triple-Negative Breast Cancer: Metastatic Risk and Role of Platinum Agents 2008 ASCO Clinical Science Symposium, 2008.
- Trivers KF, Lund MJ, Porter PL, et al (2009). The epidemiology of triple-negative breast cancer, including race. *Cancer Causes Control*. **20**, 1071-82.
- Yao S, Ambrosone CB (2012). Associations between vitamin D deficiency and risk of aggressive breast cancer in African-American women. *J Steroid Biochem Mol Biol*, **12**, 166-5.