

RESEARCH ARTICLE

Experiences of Turkish Women with Breast Cancer During the Treatment Process and Facilitating Coping Factors

Neslihan Partlak Günüşen*, Figen Şengün İnan, Besti Üstün

Abstract

Background: Breast cancer is the most common type of cancer among women in Turkey and around the world. Treatment adversely affects women's physical, psychological, and social conditions. The purpose of this study was to identify the experiences of Turkish women with breast cancer and the facilitating coping factors when they receive chemotherapy. **Methods:** A phenomenological approach was used to explain the experiences and facilitating factors of breast cancer patients during the treatment period. Data were collected through individual semi-structured interviews. The sample comprised 11 women with breast cancer receiving treatment. **Results:** At the end of the interviews conducted with women with breast cancer, two main themes were identified: adjustment and facilitating coping factors. The adjustment main theme had two sub-themes: strains and coping. Women with breast cancer suffer physical and psychological strains as well as stress related to social and health systems. While coping with these situations, they receive social support, turn to spirituality and make new senses of their lives. The facilitating coping factors main theme had four sub-themes: social support, disease-related factors, treatment-related factors and relationships with nurses. It has been determined that women receiving good social support, having undergone preventive breast surgery and/or getting attention and affection from nurses can cope with breast cancer more easily. **Conclusions:** Women with breast cancer have difficulty in all areas of their lives in the course of the disease and during the treatment process. Therefore, nurses should provide holistic care, teaching patients how to cope with the new situation and supporting them spiritually. Since family support is very important in Turkish culture, patients' relatives should be informed and supported at every stage of the treatment.

Keywords: Breast cancer - adjustment - coping - social support - Turkey

Asian Pacific J Cancer Prev, 14 (5), 3143-3149

Introduction

Breast cancer, the most common type of cancer among women all over the world (28%), is the second leading cause of cancer-related deaths (15%) (American Cancer Society: ACS, 2010). The most frequent type of cancer among women in Turkey is breast cancer (37.6 per hundred thousand) (TC. Ministry of Health, 2006). Cancer diagnosis and treatment have a profound impact on an individual's physical, psychological, and social life.

During the process of diagnosis and treatment, women with breast cancer face many physical and psychological problems in the family, at work and in social life, and their quality of life is adversely affected (Liu et al., 2006). Breast cancer is a complex disease affecting millions of people around the world. In recent years, there have been important advances in prevention, diagnosis and treatment methods of breast cancer. Survival rates in women with breast cancer are increasing and developments in breast

surgery affect women positively (Sertöz-Önen et al., 2004). With these developments, women experience changes in their perceptions of the disease and requirements as well.

Treatment methods used in the treatment of breast cancer extend life of patients; on the other hand, they can cause problems such as infection, anemia, loss of appetite, nausea, vomiting, pain and fatigue. Women diagnosed with breast cancer need help to adapt themselves to this situation (Liu et al., 2006; Remmers et al., 2010). Fear, anxiety, and uncertainty of future life experienced due to the known and unknown characteristics of the disease may lead to psychological distress in women with breast cancer (Schmid-Buchi et al., 2008; Taleghani et al., 2008). Women with breast cancer strive to cope with the adverse situations suffered during the diagnosis and treatment processes of the disease. It has been reported that women with breast cancer involve in religion more during the course of the disease and that they try to cope with the disease by turning to spiritual life (Taleghani et al., 2008).

Psychiatric and Mental Health Nursing Department, Faculty of Nursing, Dokuz Eylül University, İzmir, Turkey *For correspondence: n.gunusen@yahoo.com

Another factor stated to help women with breast cancer cope with the disease is the social support. Women with breast cancer consider the positive support from the family and spouses important (Doumit et al., 2010). If the social support is enough, it makes it easy to cope with the disease, but if not, it has a negative effect on coping (Henderson et al., 2003).

The support women with breast cancer receive from nurses during the diagnosis and treatment within the adjustment process is important. Women with breast cancer stated that their relationships with nurses greatly contributed to meet their requirements in the diagnosis and treatment processes of the disease. Women with breast cancer expect to receive support from nurses. They want to feel valued and to receive emotional support (Remmers et al., 2010). In a study conducted to determine the support needs of women with breast cancer, it was identified that what they needed most was to get enough information, which is followed by requirements necessary in the management of daily life and emotions (Griesser et al., 2011). In a study carried out by Lei et al. (2011), it was determined that there were differences between nurses and women with breast cancer regarding their perception of information needs. While women with breast cancer care for the recurrence of the disease, metastasis, the treatment plan and the side effects of the treatment, nurses give priority to the side effects of the treatment, the duration of the disease and the impact of the disease on the patient's life. During the first interview with women with breast cancer, the areas they considered difficult should be carefully determined. However, it has been reported that the needs of women with breast cancer are often overlooked (Pigot et al., 2009). It has been determined that 53% of the women with breast cancer reported that their requirements related to patient education in the process of diagnosis and treatment are not met (von Heymann-Homan et al., 2013). In another study, women with breast cancer suffered problems in areas such as physical functioning, psychological functioning, self and body image, sexuality, relationship with the partner, relationship with others, work, returning to work and social security (Pauwels et al., 2013).

Nurses play key roles in the treatment of women with breast cancer and in their adaptation to the treatment process. Nurses provide care for patients from many different cultures in the globalizing world. To know the factors that help women with breast cancer from different cultures to cope with the disease gains importance. In parallel with the changes and developments in the diagnosis and treatment processes; experiences of women with breast cancer, their coping methods and determining whether the sources of support have changed will significantly contribute to the planning of services. In this sense, it is worth understanding the experiences of Turkish Muslim women with breast cancer have during the course of the disease and the treatment process. It is thought that the results of this study can promote the ways through which nurses provide nursing care in parallel with the needs of women with breast cancer.

The purpose of this study is to identify the experiences

of women with breast cancer and the facilitating coping factors while they receive chemotherapy.

Materials and Methods

A phenomenological approach was used to explain the experiences and facilitating factors of Turkish women with breast cancer in the treatment period. The data were collected through semi-structured individual interviews.

Participants

Women over the age of 18 who were treated for breast cancer at a university hospital and were able to provide in-depth information on the research questions comprised the sample of this study. Of the breast cancer patients, those who were known to have recurrent or metastatic cancer were excluded from the study. The data collection was based on the principles of saturation (Yıldırım ve Şimşek 2006). The final sample comprised 11 breast cancer patients in the treatment period. Mean age of the patients was 48.82.

Data collection

The data were collected through semi-structured in-depth interviews. Interviews were conducted by two trained researchers and each interview was audiotape-recorded. The interviews were conducted in the chemotherapy unit. An appropriate physical environment (quiet, safe) was created for the interviews. Women with breast cancer were asked questions about experiences related to the breast cancer treatment. The research questions were: What are the changes in your life in the treatment period? How did you cope with these situations? What facilitated your coping?

Data analysis

The researchers transcribed the audiotape-recorded interviews and combined them with notes taken during the interviews. The transcripts were read through several times to get a general sense of the information and to ensure the accuracy of the data analysis. During the coding phase, two researchers independently read all transcripts and determined codes. The codes were categorized and then themes and sub-themes. After coding was completed, the researchers discussed the codes and agreed on the thematic statements (Creswell, 2003).

Trustworthiness

To achieve trustworthiness of the data were based the strategies (credibility, transferability, dependability, confirmability) established by Lincoln and Guba (1985). For the semi-structured interview questions, two experts' opinions were received. In order to access the richer variation of phenomena, variations in sampling were provided on qualifications such as age, marital status and employment status. The research method (characteristics of the participants, data collection and analysis) were clearly defined. The interviews and data analysis were performed by two researchers. The differences between the two researchers were discussed, and then the data were

organized and documented by the researchers.

Ethical considerations

Ethical approval to conduct the study was obtained from the Ethical Committee of the Dokuz Eylul University Institute of Health Sciences (decision numbered 154 and dated 03.03.2011). The patients were informed about the purpose of the study verbally and in written, and assured that they had the right to refuse to participate in the study before interview.

Results

The results about the experiences of women with breast cancer in the treatment period were classified in two main themes and six sub-themes. These two themes are adjustment and facilitating coping factors (Table 1).

Theme: adjustment

Two sub-themes (strains, and coping) were identified under the adjustment main theme.

Strains

The interviews revealed that women with breast cancer suffer physical, psychological strains and strains related to social and health systems. Physical strains experienced in the treatment process are as follows: fatigue, pain, oral and dental problems, skin / nail problems, nausea / vomiting, weight loss, loss of appetite, hair loss, and sleep problems.

"Chemotherapy is not an easy thing. I cannot eat. I have fatigue. I suffer nausea, burning sensation under my feet, weight loss, hair loss, and mouth and teeth problems. For example, my heart beats very fast. By experiencing all these, I understood what chemotherapy is." (MÇ)

"My hair shed very much. I thought I wouldn't live like this, I tried hard and went to the hair dresser and asked her to cut my hair shortest. She did." (MÇ)

"Drugs caused acnes (pimples) all over my face. Drugs caused nail problems too." (ÜS)

"You find something to keep you busy during daytime. But nights, you're alone, the most pessimistic, negative thoughts come into your mind, you cannot go to sleep." (ÇD)

Psychological strains specified by the women were uncertainty, changes in body image, loneliness, and changes in the emotional state.

One patient defined uncertainty as *"I experience emotional tides; I question myself; what am I doing; what's gonna happen to me."* (MÇ)

One of the areas uncertainty is experienced is about the effects of chemotherapy. A patient defined the chemotherapy-related uncertainty she experienced as

"now I am having treatment, I wonder if my hair will shed, if I will suffer nausea or if I won't be able to eat? I am thinking of things like these." (SY)

Women with breast cancer suffer fear of pain, metastasis, and death.

"I have fears about the next phases of the process. I do not want to be in pain. I do not want to suffer from aches." (ÇD)

"I have fears. For example, when somewhere else in my body hurts, I wonder if it has metastasized." (ÜS)

"Death is cold, and cancer reminds you death. Cancer leads to fear of death." (SK)

During the interviews, women said that they experienced body image problems due to removal of breasts and hair loss.

"One of my breasts is different. This affects me badly. For example, when I take a shower I feel bad." (SY)

"What I felt sorry about most was loss of my hair. My hair was long. I had my hair cut; my head looked different; I cried a lot." (FK)

"At the moment, I have problems with my hair, my hair sheds. This affects me negatively. I say to my friend "I want you to see me when I am well, to remember me when I look healthy." (ÇD)

Another psychological strain is loneliness. One patient defined the loneliness she experienced as *"For example, I tell someone something. Then, everyone turns back his/her own life, and you find yourself alone."* (ÜS)

Interviews conducted with women with breast cancer revealed that they experienced such changes as anger, sadness, incompetence and regret in their emotional lives.

"I wasn't like this before. When there was a problem I used to talk about it calmly, in a friendly way, but now I get furious." (SK)

"Dying is not a problem for me but I have regrets up behind me and hopes for the future. You say I wish I had done this or that. You cannot give up your hopes. That's my sorrow." (HGS)

"You want your kids not to be told, you want others not to worry, but you feel that you are shattered." (ÜS)

Women with breast cancer regarded difficulties arising in the family or behaviors like staying away from other people as social strains.

"My skin looked terrible, some people did not come close to me, did not want to kiss me. I felt this and I did not kiss anyone. But I felt bad, of course." (ÜS)

"It is just natural that everyone looks at the operated site. Has your hair or eyebrow hair shed? Are you gonna have chemotherapy? So many questions, I don't want to hear such questions so I don't talk to people." (SY)

"I had my surgery and now it's over for me. But hearing people around you saying "don't eat it this or that. You're sick" annoys me." (SY)

"My older son was negatively affected. For example, he doesn't study his lessons, he doesn't go to school. He cut classes for 5 or 6 days. I'm here, he's there." (MÇ)

Women with breast cancer regarded topics like having difficulty reaching physicians, lack of support provided for patients and their relatives, and transportation problems as strains related to the health care system.

"Here, we are looking for the doctor, with the files in

Table 1. Themes and Sub-themes

Themes	Sub-Themes
Adjustment	Strains
	Coping
Facilitating coping factors	Social Support
	Disease-Related Factors
	Treatment-Related Factors

our hands. Not being able to reach the doctor is the biggest problem. And sometimes doctors become extremely angry for no reason.” (SK)

“There is no place where women can easily reach nor do they receive psychological consultancy. For example, if there were a place the patient’s family could consult, they could get counseling while waiting for the women. It seems to me that patients’ relatives are worn out more than the patients are.” (NÇ)

“For example, tomorrow, I’m gonna take medicine, after taking medicine, getting on a vehicle, or moving around makes me very sad. It is too hard, because I cannot go home right away.” (MT)

Coping

It was determined that women with breast cancer employed the following methods to cope with the difficulties they encounter during the treatment: denial, suppression, crying, fatalistic approach, positive thinking, getting social support, turning to religion, taking up a hobby, setting priorities in life, acquiring information and rearranging their lifestyles.

“I sense the disease is taking things away from me. But I still haven’t accepted the disease.” (MÇ)

“I cried a lot, but I managed to continue my life too. I was sad, but I continued.” (ÜS)

“Disease is something that comes from God. There exists death. We’re all gonna die someday.” (ŞB)

“I’m reasoning the event out on my own. I say it is nothing. I’ll overcome this too. I’m telling these to myself. I mean, the key to achievement, of course, is in your own hands.” (MÇ)

“I share the difficulties with my husband. I can easily tell him, he supports me.” (ÜS)

“When the test results appeared to be good, I started to go to the hospital willingly. I give thanks to my God.” (HGS)

“I love crocheting, I spend all my time crocheting, when I am bored, I go out and get around.” (SY)

“I’ve always tried hard, worked hard, I burned myself out, I’ve always thought of other people; I’ve always placed myself in the background, but now I say it is I who is in the first place.” (ÇD)

“Because I don’t know how much time I have left, I cling to life more. I try to share more things with the kids. I am living life to the fullest, I wasn’t like this before.” (MÇ)

“What is chemotherapy, how it affects the human body, what I will experience before and after chemotherapy, what psychological problems I will have ... I research all these on the internet. In addition, I exercise; I’m trying to do what the doctor has told to me to do.” (NÇ)

Theme: facilitating coping factors

Four sub-themes were determined under the “Facilitating Coping Factors” main theme: social support, disease-related factors, treatment-related factors and relationship with nurses.

Social support

The participants in this study consider social support

as one of the most important factors that facilitate their coping with difficulties.

“My family is very supportive, I continue my everyday life, I have never been introverted because I am sick, and they didn’t allow me to become introverted. That’s why I’m alright.” (SY)

“My neighbors are very supportive. When I come here for the therapy, my husband looks after my elder son and my sister in law looks after my younger son.” (MÇ)

Disease related factors

The participants regarded high breast cancer survival rates as a protective factor in the disease-fighting process.

“As long as you take care of yourself, no deaths from breast cancer will occur. That’s why I’m telling myself I’m fine and I’ll be fine.” (SY)

Treatment related factors

The participants regarded breast-conserving surgery, the positive response to the treatment and the positive approach displayed by the health care team as protective factors.

“I just heard my test results are good. After getting these results, having nice talks with the doctors makes me feel better.” (MC)

“My breast was not completely removed. That’s why I’m psychologically more comfortable.” (SY)

Relationships with nurses

During the interviews, the women indicated the relationship with the nurses during the treatment period as a facilitating factor in coping with the disease. The participants expect nurses to be helpful and friendly.

“The nurses are performing our treatments; they show us where to go” (E.Ö)

“Nurses are very supportive, they say you’re gonna be well again, they give morale. They give us our medicines; they say “good morning; they talk to us.” (FK)

I expect attentiveness and a smiling face from them. I don’t expect anything else.” (ÇD)

Discussion

In this study, two main themes were determined: adjustment and facilitating coping factors. The adjustment theme has two sub-themes: strains and coping. The women participating in the study suffered physical, psychological, social and system-related strains. These findings account for the women’s needs related to these four areas.

The participants stated that they were adversely affected due to the side effects of chemotherapy. Women with breast cancer suffer physical symptoms such as changes in taste, alopecia, nausea, vomiting, mucositis and fatigue due to chemotherapy (Stein et al., 2003). These side effects can lead to psychological problems which affect the patient’s activities of daily living (Erci and Karabulut, 2007). In addition to the psychological distress caused by being diagnosed cancer, chemotherapy-related side effects cause psychological distress too. Psychological strains suffered by women with breast cancer most are uncertainty, fear of cancer recurrence, negatively affected

body image, loneliness, anger and sadness. These findings are similar to the findings the literature. It has been indicated that cancer diagnosis and treatment cause the following psychological problems in women with breast cancer: anxiety, depression, anger, uncertainty about the future, fear of cancer recurrence, body image distortion, fear of losing female identity (Doumit et al., 2010; Denieffe and Gooney, 2011; Cebeci et al., 2012). In the literature, uncertainty and fear of recurrence are stated to be the most challenging experiences that women with breast cancer have in the process of diagnosis and treatment of the disease (Landmark et al., 2001; Doumit et al., 2010; Elmir et al., 2010). In our study too, women with breast cancer experienced problems such as uncertainty about the future and fear of cancer recurrence intensely. In their qualitative study of women with breast cancer, Doumit et al. (2010) determined that fear of cancer recurrence and changing body image had a negative impact on patient's coping with the disease.

This study also determined that women with breast cancer suffered social strain as well. Women with breast cancer have trouble in the family and move away from social environments. It is reported women with breast cancer are not sure how people around them would react to their illness; therefore, they do not share the disease and stay away from social environments (Brede, 1999; Louma and Hakamies-Blomqvist, 2004). In our study too, patients stay away from social environments not to hear questions of curious people or due to changes in their appearances. In Turkish society, familial and neighborly relationships are very strong. These relationships might be supporting, but on the other hand, they might lead to such problems as curiosity about and/or interfering in someone else's life. This leads to difficulties arising from the environment.

In a study of women with breast cancer conducted by Remmers et al. (2010), it was reported that especially those with small children or those who have family members who need care worried about their family members rather than about themselves. In our study too, it was determined that patients had difficulties in their family lives. In a Turkish family, women play an important role in ensuring and maintaining family functioning. Turkish women are responsible for the care of the children, all of the housework and the maintenance of intra-family relationships. Therefore, changes in the woman's health result in changes in the life of the whole family. It has been determined that women with breast cancer experience physical problems due to the disease and the treatment of the disease and that these physical problems affect them both psychologically and socially. Therefore, it is very important for health professionals to holistically evaluate women with breast cancer and to be aware of the fact that changes occurring in one area of their lives can affect the other areas of their lives as well.

Difficulties experienced by women with breast cancer are not limited to disease- and treatment-related difficulties. Patients have problems such as obtaining treatment, reaching physicians and struggling with bureaucratic barriers. According to the results of a review regarding women living in rural regions by Bettencourt et al. (2007), women with breast cancer experience problems

of transportation and accommodation. In addition, patients are negatively affected because they cannot adequately communicate with physicians and have to wait for a long time to receive the test results. In a study of women with breast cancer by Remmers et al. (2010), patients stated that they needed a friendly environment all over the hospital so that they could feel comfortable. System-related problems increase stressors in women with breast cancer who have already been suffering from depressive experiences, which affects them physiologically and psychologically too.

Another sub-theme of the adjustment main theme of the study is coping. Women with breast cancer struggle to cope with the strains they suffer. It was determined that patients resort to positive thinking, social support, turning to religion, taking up a hobby, setting priorities in life, acquiring information and rearranging their lifestyles for coping. It has been determined that Iranian Muslim women tend to involve in religious activities such as reading the Qur'an more after being diagnosed breast cancer. In addition, women with breast cancer reported that believing in that "diseases come from God" relieved them (Taleghani et al., 2008). In a study conducted by Cebeci et al. (2012), it was found that after having cancer, women experienced positive changes in their lives and their perspectives of the world changed. Women with breast cancer reported that they clung to life more than they did before the diagnosis and treatment process and that life became more valuable for them. The process of diagnosis and treatment of breast cancer causes a traumatic effect on individuals. After this experience, many changes occur in individuals. The meaning they attribute to life and priorities in their lives may change (Landmark et al., 2001). Tedeschi and Calhoun (2004) define this as post-traumatic growth. Post-traumatic growth can be defined as "positive psychological changes experienced as a result of struggle with highly challenging life circumstances" (Tedeschi and Calhoun, 2004). Major domains of posttraumatic growth are: greater appreciation of life and changed sense of priorities, warmer, more intimate relationship with others, a greater sense of personal strength, recognition of new possibilities or paths for one's life and spiritual development (Tedeschi and Calhoun, 2004). Women with breast cancer tend to consider materialistic factors less important after experiencing the disease. While the importance attributed to the demands of everyday life diminishes, developing better relationships with their relatives and friends becomes more and more important.

Another main theme of the study is the factors affecting coping. This main theme has four sub-themes: social support, disease-related factors, treatment-related factors and relationships with nurses. The majority of women with breast cancer participating in the study indicated that social support was the most important factor for their coping with the disease. While those who had adequate social support stated that they were able to cope with the disease better, those who did not have adequate social support stated that this lack of social support made their coping harder. The participants of the study stated that they wanted to survive for the sake of their children and that having children motivated them to fight the disease. These findings support the findings of other studies indicating

that the positive family support plays an important role in coping with breast cancer. In the literature too, it is indicated that social support received from the family and friends during the healing process is of great importance for women with breast cancer. It is stated that social support is an important factor that facilitates coping (Connell et al., 2006). In a study evaluating hopelessness levels of women with breast cancer, it was determined that women receiving social support had a lower level of hopelessness (Öztunç et al., 2013). In a study carried out by Doumit et al. (2010), positive support from family and colleagues was indicated to be one of the factors that help women with breast cancer to cope with the disease better. In their study of women with breast cancer, Henderson et al. (2003) pointed out that support from the family and/or spouse was a facilitating coping factor, but not in all cases. In their study investigating the experiences of women with breast cancer, Cebeci et al. (2012) stated that all the participants of their study received physical, psychological and social support from their families and emphasized the importance of the family in Turkish society. In Turkish family structure, social support is of great importance. A patient's needs are met by the other family members. On the other hand, some participants, though not many, stated that they did not receive adequate social support. However, most of the women stated that the other members of the family fulfilled the responsibilities they were not able to do due to the disease. Families support the women with breast cancer not only by performing their roles but also by motivating them psychologically, which makes it easier for the women with breast cancer to cope with the disease better.

Disease- and treatment-related factors make it easier for women with breast cancer to cope with the disease too. In this study, it was determined that women with breast cancer perceived breast cancer more positively than they did other types of cancer. Although the incidence of breast cancer is high, women's survival time has prolonged in recent years thanks to effective treatment methods. A review of survival rates of different cancer types unique to women reveals that the highest survival rate (40%) is among women with breast cancer (Knobf, 2007). This situation has widespread media coverage and the survival rate is known to be high, which reduce women's negative perceptions of breast cancer diagnosis. The participants considered undergoing breast-conserving surgery or removal of their breasts as a facilitating coping factor. In their study conducted to evaluate the impact of the type of the breast surgery on women with breast cancer, Önen-Sertöz et al. (2004) determined that total mastectomy affected the women's perception of breast and self-esteem negatively. Furthermore, half of the women who underwent total mastectomy said that they would have preferred breast-conserving surgery if they had had another opportunity. Changes in the body image occurring after mastectomy may be an additional stressor for the patient in the treatment of the disease, which itself is already difficult. Therefore, it is thought that surgeries like breast-conserving surgery do not create stressors for women and facilitate their coping with the disease. These results suggest that recent developments in breast cancer

treatment have changed breast cancer women's perception of the disease.

The relationship between nurses and women with breast cancer was determined to be one of the factors that facilitate these women's coping with cancer. Women with breast cancer expect to get sympathy and a smiling face from nurses. These results are similar to those of the studies carried out in Turkey. However, studies conducted especially in Western countries indicate that breast cancer women's expectations of nurses have increased. According to women in Turkey, if nurses are smiling this would be enough, but in western countries, nurses are expected to be professionally competent too. In a study of women with breast cancer by Remmers et al. (2010), patients expected nurses to stand by them in their hard times, to give them emotional support and to have professional competence. Gilbert et al. (2010) evaluated positive and negative interactions between health professionals and relatives of cancer patients. As a result, of the health personnel, those who are warm, courteous, accessible and approachable and make patients feel relaxed are considered to interact with patients positively. On the other hand, patients do not have good perception of health personnel who do not empathize with them well enough or do not give enough information on the management of cancer. In our study too, patients indicated that nurses' friendly approaches relieved them. In order for women with breast cancer to talk about their needs freely, nurses are required to establish a friendly and reliable environment. Unless this environment is established, patients will not share their fears and sex-related problems with nurses.

Özsoy and Özgür (2007) reviewed the literature in order to investigate the studies carried out in Turkey on women's expectations of nurses. As a result, they determined that women expect nurses to be smiling, understanding, affectionate, courteous, and helpful. On the other hand, it was determined that nurses' professional competence was not among the women's expectations of nurses. This suggests that if nurses establish a positive relationship with women this would content women. However, this fact, in other words, Turkish people's only expectation of nurses is their establishing a positive relationship with patients but no professional effort, is worrisome. For women, nurses' displaying a humanistic approach is good enough. This may be related to the society's perception of the nursing profession. Since women do not know much about roles and functions of nurses, they do not have any expectations regarding care.

The results of this study are expected to shed light both on giving effective culture-specific nursing care and on cross-cultural comparison studies. Although there are changes in the treatment of breast cancer, women with breast cancer still have difficulty in every area of their lives over the course of the disease and treatment. Therefore, nurses should provide holistic care and be aware of patients' strains. Besides being supportive, nurses should have enough knowledge about how to teach ways of coping with the disease and how to spiritually (psychologically) strengthen women with breast cancer. Nurses should be aware of the importance of information for women with breast cancer, should give them the

opportunity to ask questions freely and should plan the training programs in line with the information needs of women with breast cancer. Nurses should also provide information, skills, and support that the other family members need.

Since family support is very important in Turkish culture, patients' relatives should be informed and supported at every stage of the treatment. The results of this study demonstrated once again the importance of dealing with women and their family together. It is considered that conducting studies on the experiences of relatives of women with breast cancer in the future would be beneficial.

References

- American Cancer Society (2010). Cancer facts and figures 2010. Atlanta, American Cancer Society.
- Bettencourt BA, Schlegel RJ, Talley AE, Molix, L (2007). The breast cancer experience of rural women: a literature review. *Psychooncology*, **16**, 875-87.
- Bredin M (1999). Mastectomy, body image and therapeutic massage: a qualitative study of women experience. *J Adv Nurs*, **29**, 1113-20.
- Cebeci F, Balci YH, Tekeli, A (2012). Life experiences of women with breast cancer in South Western Turkey: a qualitative study. *Eur J Oncol Nurs*, **16**, 406-12.
- Connell S, Patterson C, Newman B (2006). Issues and concerns of young Australian women with breast cancer. *Supportive Care Cancer*, **14**, 419-26.
- Creswell JW (1998). Qualitative inquiry and research design: choosing among five traditions. 2nd Edition. Sage Publications, 190-6.
- Denieffe S, Gooney M (2011). A meta-synthesis of women's symptoms experience and breast cancer. *Eur J Cancer Care*, **20**, 424-35.
- Doumit MAA, Huijjer HA, Kelley JH, El Saghir N, Nassar A (2010). Coping with breast cancer a phenomenological study. *Cancer Nursing*, **33**, 33-9.
- Elmir R, Jackson D, Beale B, Schmied V (2010). Against all odds: Australian women's experiences of recovery from breast cancer. *J Clin Nur*, **19**, 2531-8.
- Erci N, Karabulut N (2007). Appraising the self- assessed support needs of Turkish women with breast cancer. *Eur J Oncol Nursing*, **14**, 42-8.
- Gilbert E, Ussher JM, Perz J, Hobbs K, Kirsten L (2010). Positive and negative interactions with health professionals. *Cancer Nursing*, **33**, 1-9.
- Griesser AC, Vlastos G, Morel L, et al (2011). Socio-demographic predictors of high support needs in newly diagnosed breast cancer patients. *Eur J Cancer Care*, **20**, 466-74.
- Henderson PD, Gore SV, Davis BL, Condon EH (2003). African American women coping with breast cancer: a qualitative analysis. *Oncol Nurs Forum*, **30**, 641-7.
- Knobf MT (2007). Psychosocial responses in breast cancer survivors. *Semin Oncol Nurs*, **23**, 71-83.
- Landmark BT, Strandmark M, Wahl AK (2001). Living with diagnosed breast cancer - the meaning of existential issues. *Cancer Nurs*, **24**, 220-6.
- Lei CP, Har YC, Abdullah KL (2011). Informational needs of breast cancer patients on chemotherapy: differences between patients and nurses perceptions. *Asian Pac J Cancer Prev*, **12**, 797-802.
- Lincoln SY, Guba GE (1985). Naturalistic Inquiry. Sage Publications Beverly Hills, 289-300.
- Liu L, Li C, Tang Huang CS, Chiou AF (2006). Role of continuing supportive cares in increasing social support and reducing perceived uncertainty among women with Newly Diagnosed Breast Cancer in Taiwan. *Cancer Nurs*, **29**, 273-82.
- Luoma M, Hakamies-Blomqvist L (2004). The meaning of quality of life in patients being treated for advanced breast cancer: a qualitative study. *Psychooncology*, **13**, 729-39.
- Önen-Sertöz Ö, Elbi-Mete H, Noyan A, Alper M, Kapkaç M (2004). Meme kanserinde ameliyat tipinin beden algısı, cinsel işlevler, benlik saygısı ve eş uyumuna etkileri: kontrollü bir çalışma. (Effects of Surgery Type on Body Image, Sexuality, Self-Esteem, and Marital Adjustment in Breast Cancer: A Controlled Study) *Türk Psikiyatri Derg*, **15**, 264-75.
- Özsoy SA, Özgür G, Durmaz Akyol A (2007). Patient expectation and satisfaction with nursing care in Turkey: a literature review. *Int Nurs Rev*, **54**, 249-55.
- Öztunç G, Yeşil P, Paydaş S, Erdoğan S (2013). Social support and hopelessness in patients with breast cancer. *Asian Pac J Cancer Prev*, **14**, 571-8.
- Pauwels EEJ, Charlier C, De Bourdeaudhuij I, Lechner L, Van Hoof E (2013). Care needs after primary breast cancer treatment. Survivors' associated sociodemographic and medical characteristics. *Psychooncology*, **22**, 125-32.
- Pigot C, Pollard A, Thomson K, Aranda S (2009). Unmet needs in cancer patients: development of supportive needs screening tool (SNST). *Support Care Cancer*, **17**, 33-45.
- Remmers H, Holtgrawe M, Pinkert C (2010). Stress and nursing care needs of women with breast cancer during primary treatment: A qualitative study. *Eur J Oncol Nurs*, **14**, 11-6.
- Scmid-Buchi S, Halfens RJG, Dassen T, Van Den Borne B (2008). A review of psychosocial needs of breast-cancer patients and their relatives. *J Clin Nurs*, **17**, 2895- 909.
- Stein KD, Denniston M, Baker F, et al (2003). Validation of modified rotterdam symptom checklist for use with cancer patients in the United States. *J Pain Symptom Manage*, **26**, 975-89.
- Taleghani F, Yekta ZP, Nasrabadi NA, Kappeli S (2008). Adjustment process in Iranian women with breast cancer. *Cancer Nurs*, **31**, 32-41.
- Tedeschi RG, Calhoun LG (2004). Posttraumatic growth: conceptual foundations and empirical evidence. *Psychol Inq*, **15**, 1-18.
- Turkey Ministry of Health (2008). Cancer statistics of Turkey in 2008, available at <http://kanser.thsk.gov.tr/index.php/daire-faaliyetleri/kanser-istatistikleri.html> (Accessed:01 April 2013).
- Von Heymann-Horan AB, Dalton OS, Dziekanska A, et al (2013). Unmet needs of women with breast cancer during and after primary treatment: a prospective study in Denmark. *Acta Oncol*, **52**, 382-90.
- Yıldırım A, Şimşek H (2006). Sosyal Bilimlerde Nitel Araştırma Yöntemleri. (Qualitative research methods in social sciences) 5th ed, Seçkin press, Ankara, Turkey, 114-115.