

## RESEARCH ARTICLE

# Lack of Association Between CYP1A1 Polymorphisms and Risk of Bladder Cancer: a Meta-analysis

Yu Lu, Xiao-Lian Zhang, Li Xie, Tai-Jie Li, Yu He, Qi-Liu Peng, Yan Deng, Jian Wang, Xue Qin\*, Shan Li

## Abstract

**Background:** The effects of CYP1A1 gene polymorphisms on the risk of bladder cancer (BC) remain controversial. We carried out a meta-analysis to clarify the role of CYP1A1 gene polymorphisms in BC. **Material and Methods:** A comprehensive literature search was conducted up to November 20, 2013. Odds ratios (ORs) with 95% confidence intervals (CIs) were used to estimate the strength of the association. Meta-regression, subgroup analysis, sensitivity analysis and publication bias were also performed. **Results:** Eight studies involving 1,059 BC cases and 1,061 controls were included. The meta-analysis showed that there was no significant association between the two common mutations of CYP1A1 and BC risk. For the Ile462Val A/G polymorphism with GG vs. AA the OR was 1.47 (95% CI= 0.70-3.07,  $P=0.308$ ). For the MspI T/C polymorphism, though a slight trend was found this was not statistically nonsignificant (CC vs. TT, OR = 1.24, 95% CI= 0.98-1.58,  $P=0.078$ ). Subgroup analyses by ethnicity also found no obvious association between CYP1A1 and BC risk. **Conclusion:** The present meta-analysis suggests that CYP1A1 polymorphism is not associated with bladder cancer risk.

**Keywords:** Bladder cancer - CYP1A1 - Ile462Val - mspI – polymorphisms - meta-analysis

*Asian Pac J Cancer Prev*, 15 (9), 4071-4077

## Introduction

Bladder cancer (BC) remains the second leading malignancy of the genitourinary tract and the seventh most frequently diagnosed cancer in males, accounting for approximately 5.9% (386,300) of the new cancer cases and 3.4% (150,200) of the estimated cancer deaths annually worldwide. Bladder is the major cancer sites for males in Jordan (Ismail et al., 2013); and with an age-specific incidence rate of about 11.2/100,000, BC is also the fifth most common cancer among men in Iran (Karbakhsh et al., 2013). The etiology and carcinogenesis of BC is unclear, but smoking, especially opium abusing simultaneously (Shakhssalim et al., 2010), is thought to be a major risk factor followed by occupational exposure (Zeegers et al., 2000). Over 60 carcinogens, including polycyclic aromatic hydrocarbons (PAHs) such as benzo[a]pyrene and aromatic amines, have been identified in cigarette smoke (Luch et al., 2005). These carcinogens, which have been proven to be associated with BC in aluminum workers exposure to PAHs and coal gasification workers exposure to aromatic amines (Boffetta et al., 1997), can be converted, in vivo, into more hydrophilic and chemically active derivatives by the CYP450 enzyme superfamily (Grando et al., 2009).

CYP1A1, an important phase I xenobiotic metabolizing

enzyme of the cytochrome P450 superfamily, is well known for its involvement in the metabolic activation of tobacco procarcinogens such as PAHs and aromatic amines (Vineis et al., 2007), and has been found in many epithelial tissues (Romkes et al., 1996). It is a highly polymorphic gene with more than 11 alleles thought to lead to amino acid changes (Shimada et al., 2006). One of the most common mutations described is a single-base exchange from A to G at position 2455 of exon 7, also called CYP1A1\*2B or m2 (Hayashi et al., 1991). Polymorphism of this heme-binding region causes a substitution of isoleucine with valine in codon 462 (Ile462Val), resulting in an alteration of the functional protein and thus increasing enzyme activity (Kawajiri et al., 1996). Furthermore, mutation of this gene exhibits strong potential linkage disequilibrium with another mutation of CYP1A1-MspI (m1) polymorphism (Bartsch et al., 2000), a point mutation (T to C) at the MspI site in the 3'-untranslated region, and has also been proven to be associated with increased catalytic activity.

Such genetic differences in enzymes involved in the biotransformation of environmental risk factors are believed to play a pivotal role in an individual's susceptibility to environmentally induced cancer and their relationship with various malignancies has been extensively studied (Bartsch et al., 2000). An initial study

regarding the association between CYP1A1 variations and cancer risk was conducted by Kellerman et al. in 1973 (Kellermann et al., 1973), investigating the correlation between benzo[a]pyrene hydroxylase inducibility and bronchogenic carcinoma. Since then, numerous studies on CYP1A1 polymorphisms and various cancers have been conducted, including lung cancer (Ji et al., 2012), head and neck cancer (Hiyama et al., 2008), brain cancer (Wahid et al., 2013), and breast cancer (Sergentanis et al., 2010), among others (Dai et al., 2009). However, less attention has been paid on the relationship between CYP1A1 variations and BC, with only a few studies having been conducted and their conclusions remaining controversial (Bartsch et al., 2000). As the statistical power of an individual study may be too weak to identify associations between CYP1A1 polymorphisms and BC risk, a meta-analysis combining data from all published studies may be more convincing. We thus carried out a meta-analysis to evaluate the association between CYP1A1 polymorphisms and BC risk including all eligible publications to date.

## Materials and Methods

### Literature Search

A comprehensive literature search was carried out by two independent investigators. The PubMed, EMBASE, China National Knowledge Infrastructure (CNKI), and WanFang databases were searched using the following keywords: “CYP1A1”, “bladder cancer”, and “polymorphism”, in both English and Chinese. Synonyms and different search term styles were also used to obtain every relevant paper. Further, the bibliographies of all retrieved articles were manually checked for other relevant publications to find additional eligible studies. Searches were not restricted in language or publication date, although no efforts were made to obtain unpublished studies. The final search was updated on November 20, 2013.

### Selection Criteria

All identified articles were reviewed according to the inclusion criteria before further analysis, namely any type of comparative study that i) assessed the association between CYP1A1 gene polymorphisms and BC, and ii) provided the frequencies of the CYP1A1 variants in both cases and controls, or provided sufficient data to estimate the odds ratio (OR) with their 95% confidence intervals (95% CIs). Studies were excluded from the meta-analysis if they met any of the following: i) without sufficient data; ii) a lack of control population; and/or iii) assessed the association between CYP1A1 and BC in rare alleles. In the event of overlapping data, either the study with higher quality or the most recent one was included in the analysis.

### Data Extraction

For each eligible study, the following information was extracted: first author, year of publication, study region, ethnicity, BC confirmation, sample size (including number of cases and controls), source of control (together with matching criteria), polymorphisms of CYP1A1, methods used for genotyping, genotype distribution in

cases and controls, and whether *P* value for the control population deviated from the Hardy-Weinberg equilibrium (HWE). Such extraction was performed by two authors independently. In the cases of conflict evaluation, a third author was consulted.

### Quality Assessment

We evaluated the quality of included studies using the set of predetermined criteria derived by Thakkinstian et al. (Thakkinstian et al., 2005). This set of predetermined criteria was structured as a 20-item list with scores ranging from 0 to 12 by Peng et al. (Li et al., 2013) (Supplement Table S2), and has been quoted by several meta-analyses (Lu et al., 2013, Peng et al., 2013). As previously performed in other meta-analyses, we also considered articles with scores <8 as low quality studies, while the rest were high quality.

### Statistical Analysis

The strength of association between CYP1A1 gene polymorphisms and BC risk was assessed by calculating crude ORs with the corresponding 95% CIs under the additive, dominant, and recessive genetic models. The stratification analysis was also conducted by ethnicity to evaluate the effect of CYP1A1 gene polymorphisms on the susceptibility to BC in different populations (categorized as Turks, Caucasians, and other, based on the main racial group of the included studies). To assess heterogeneity in each combined analysis, Cochran's *Q* test and *I*<sup>2</sup> statistics were carried out, where  $P_Q < 0.10$  or  $I^2 > 50\%$  indicated significant heterogeneity. In the situation of high heterogeneity, the random-effects model was used to pool the data; otherwise, the fixed-effects model was used. Additionally, if significant heterogeneity was detected, logistic meta-regression was performed to identify four possible sources of heterogeneity including ethnicity, BC confirmation method, genotyping method, and quality score. Moreover, Galbraith plots were also used to further explore the sources of heterogeneity among studies.

A sensitivity analysis was conducted to assess the influence of a single study on the overall result of the meta-analysis, especially for studies whose genotype frequencies in the control populations were inconsistent with the HWE, given that they may generate possible bias. Potential publication bias was evaluated using funnel plots and Egger's regression tests to provide both graphical and statistical evidence. All data analysis was completed by using STATA software version 11.0 (Stata Corp, College Station, TX, USA). A value of  $P < 0.05$  was considered as statistically significant, and all *P* values were two-sided.

## Results


### Study Characteristics

The systematic literature search generated a total of 39 citations based on the search strategy mentioned above (31 in PubMed and EMBASE, and 8 in the CNKI and WanFang databases), 16 of which were excluded after screening of the titles and abstracts, for they were not relevant to our study. Therefore, 23 articles were considered of potential value and the full text was

**Table 1. Detail Characteristics of Studies Included in This Meta-analysis**


Author, Year	Region	Ethnicity	case/control	Genotyping methods	BC confirmation	Source of control	PI	HWE (Yes/No)	QS
Brockmüller J, 1996	Germany	Caucasian	368/359	AS-PCR	HD	HB (matched for age, gender and race)	CYP1A1*2B A/G (I1e462Val)	Yes	6
Grando JP, 2009	Brazil	Mix	100/100	AS-PCR	HD	HB (matched for age, gender, race, and smoking status)	CYP1A1*2B A/G (I1e462Val)	Yes	9
Fontana L, 2009	French	Caucasian	51/45	PCR	HD	HB (age and sex-matched cancer-free subjects)	CYP1A1*2B A/G (I1e462Val)	No	7
Öztürk T, 2011	Turkish	Non-Caucasian	176/97	PCR-RFLP	PD	HB (matched for age)	CYP1A1*2B A/G (I1e462Val)	No	6
Berber U, 2013	Turkish	Non-Caucasian	114/114	AS-PCR	HD	HB (age and sex-matched cancer-free subjects)	CYP1A1*2B A/G (I1e462Val)	Yes	8
Fu J, 2013	China	Asian	99/100	PCR-RFLP	PD	HB (matched for age, gender, race, and smoking status)	CYP1A1*2B A/G (I1e462Val)	Yes	8
Brockmüller J, 1996	Germany	Caucasian	369/360	PCR-RFLP	HD	HB (matched for age, gender and race)	CYP1A1*2A T/C (MspI)	Yes	6
Yang LX, 2007	China	Asian	44/85	PCR-RFLP	PD	HB (matched for age and race)	CYP1A1*2A T/C (MspI)	Yes	5
Srivastava DS, 2008	North India	Non-Caucasian	106/160	PCR-RFLP	HD	HB (matched for age and race)	CYP1A1*2A T/C (MspI)	Yes	8

AS-PCR, Allele specific polymerase chain reaction; PCR-RFLP, Polymerase chain reaction-restriction fragment length polymorphism; HD, Histologically diagnosed; PD, Pathologically diagnosed; HB, Hospital-based; PB, Population-based; PI, Polymorphism (s) investigated; HWE, Hardy-Weinberg equilibrium; QS, quality score

**Figure 1. Process of Data Extraction**

retrieved for a detailed evaluation. An additional 15 articles were excluded following full text review; nine were review articles or conference abstracts, three were without sufficient information to access ORs and 95% CIs, two assessed the association between CYP1A1 polymorphisms and urothelial cancer, and one assessed the association between CYP1A1 polymorphisms and BC in three rarely studied alleles. Manual search of the reference lists did not reveal any additional eligible studies. Thus, eight relevant studies focused on two of the most studied polymorphisms in CYP1A1 (CYP1A1\*2B A/G (I1e462Val) and CYP1A1\*2A T/C (MspI) and risk of BC were included in our meta-analysis (Brockmoller et al., 1996; Li et al., 2007; Srivastava et al., 2008; Grando et al., 2009; Fontana et al., 2009; Ozturk et al., 2011; Fu et al., 2013; Berber et al., 2013). Of these, the study by Brockmüller et al. (Brockmoller et al., 1996) identified the relationship between CYP1A1 polymorphisms and BC risk in both alleles, and were therefore treated independently. Consequently, nine separate comparisons involving 1,059 BC cases and 1,061 controls were included. The process of data extraction is illustrated in Figure 1.

From the nine studies reporting an association between CYP1A1 polymorphisms and BC risk, six studies, with a total of 908 cases and 815 controls, evaluated the

**Figure 2. Forest Plots of CYP1A1 Polymorphisms and Risk of Bladder Cancer.** a Forest plots of CYP1A1\*2B A/G (I1e462Val) polymorphism and bladder cancer risk (contrast GG vs. AA); b Forest plots of CYP1A1\*2A T/C (MspI) polymorphism and bladder cancer risk (contrast CC vs. TT).

association of BC risk in I1e462Val A/G mutation (two in Caucasians, two in Turks, one in Asian, and one was a mixed population); three studies, with a total of 519 cases and 605 controls, evaluated MspI T/C mutation (one in Caucasians, one in Turks, and one in Asian). Of all the eligible studies, the majority of BC patients were histologically diagnosed (six studies) for their genotype using PCR-RFLP assays (five studies), the rest were pathologically diagnosed and genotyped with allele-specific PCR or standard PCR methods. After testing for concordance with the HWE principle, two studies were found to deviate from HWE. The quality score of all eligible studies ranged from 5 to 9, with four studies being graded as high quality and five as low quality; the detailed characteristics are shown in Table 1.

### Meta-Analysis Results

**CYP1A1\*2B A/G (I1e462Val) and Bladder Cancer Risk:** As established above, six studies, with a total of 908 cases and 815 controls, evaluated the association of BC risk in I1e462Val A/G mutation (Brockmoller et al., 1996; Fontana et al., 2009; Grando et al., 2009; Ozturk et al., 2011; Berber et al., 2013; Fu et al., 2013). According to our results, no significant association between this polymorphism and BC risk was found in all genetic models in the overall populations: i) G vs. A (OR = 1.04, 95% CI = 0.74–1.47,  $P = 0.810$ ); ii) GG vs. AA (OR = 1.47, 95% CI = 0.70–3.07,  $P = 0.308$ ); iii) AG vs. AA (OR =

**Table 2. Meta-analysis of the CYP1A1 Gene Polymorphisms on Bladder Cancer Risk**

Genetic model	Population	No. of studies	Test of association		Mode	Heterogeneity		Publication bias
			OR(95%CI)	<i>p</i> value		I <sup>2</sup> (%)	<i>p</i> value	
CYP1A1*2B A/G(I1e462Val)								
G vs. A	Overall	6	1.04(0.74-1.47)	0.810	R	47.2	0.092	0.451
	Turkish	2	0.89(0.63-1.28)	0.539	F	31.1	0.228	—
	Caucasian	2	0.69(0.37-1.27)	0.230	F	47.9	0.166	—
	Other	2	1.39(0.98-1.98)	0.068	F	37.4	0.206	—
GG vs. AA	Overall	6	1.47(0.70-3.07)	0.308	F	0.0	0.789	0.416
	Turkish	2	1.60(0.41-6.20)	0.498	F	0.0	0.803	—
	Caucasian	2	0.61(0.10-3.63)	0.586	F	36.9	0.208	—
	Other	2	1.89(0.67-5.35)	0.232	F	0.0	0.890	—
AG vs. AA	Overall	6	0.97(0.74-1.27)	0.819	F	40.9	0.132	0.792
	Turkish	2	0.77(0.51-1.17)	0.224	F	48.7	0.163	—
	Caucasian	2	0.73(0.37-1.42)	0.351	F	0.0	0.918	—
	Other	2	1.42(0.76-2.64)	0.271	R	50.2	0.157	—
GG+AG vs. AA	Overall	6	1.01(0.68-1.48)	0.970	R	47.2	0.092	0.709
	Turkish	2	0.82(0.54-1.22)	0.326	F	48.3	0.164	—
	Caucasian	2	0.70(0.37-1.33)	0.280	F	0.0	0.407	—
	Other	2	1.44(0.95-2.19)	0.084	F	47.0	0.170	—
GG vs. AG+AA	Overall	6	1.45(0.70-3.02)	0.317	F	0.0	0.809	0.445
	Turkish	2	1.79(0.46-6.87)	0.399	F	0.0	0.906	—
	Caucasian	2	0.61(0.10-3.65)	0.592	F	37.4	0.206	—
	Other	2	1.71(0.61-4.80)	0.308	F	0.0	0.992	—
CYP1A1*2A T/C(MspI)								
C vs. T	Overall	3	1.24(0.98-1.58)	0.078	F	1.6	0.362	0.953
CC vs. TT	Overall	3	1.93(0.95-3.93)	0.071	F	0.0	0.787	0.310
TC vs. TT	Overall	3	1.19(0.89-1.60)	0.24	F	0.0	0.398	0.644
CC+TC vs. TT	Overall	3	1.24(0.93-1.64)	0.14	F	16.0	0.304	0.673
CC vs. TC+TT	Overall	3	1.67(0.84-3.34)	0.145	F	0.0	0.883	0.430

R, random effects model; F, fixed effects model

0.97, 95% CI = 0.74–1.27,  $P = 0.819$ ); iv) GG+AG vs. AA (OR = 1.01, 95% CI = 0.68–1.48,  $P = 0.970$ ); v) GG vs. AG+AA (OR = 1.45, 95% CI = 0.70–3.02,  $P = 0.317$ ). In the stratification analysis based on ethnicity, we also failed to find any significant association between the I1e462Val A/G polymorphism and BC risk in all comparison models (Table 2). A similar result was also found when limiting data to studies whose genotype in the control population was consistent with HWE (data not show).

As significant heterogeneity was found in the G vs. A model, the GG+AG vs. AA model in the overall population, and the AG vs. AA model in the “other” population subgroup, the random-effects model was used to pool these data. Meta-regression analyses and Galbraith plot analysis were also performed to explore the source of heterogeneity between the three comparison models. The results of meta-regression analyses revealed that sources of heterogeneity were not from any of the factors mentioned above (ethnicity, BC confirmation method, genotyping method, and quality score) in all three genetic models (data not shown). Galbraith plots identified that the study by Grando et al. (Grando et al., 2009) was the outlier and main contributor to heterogeneity in the three comparison models (Figure 2). Separate forest plots omitting the outlier study were conducted in each genetic model, the degree of heterogeneity decreased significantly (G vs. A:  $I^2 = 15.4%$ ,  $P_Q = 0.317$ ; GG+AG vs. AA:  $I^2 = 0.0%$ ,  $P_Q = 0.437$ ) with the insignificance result remaining the same (G vs. A: OR = 0.92, 95% CI = 0.71–1.18,  $P = 0.507$ ; GG+AG vs. AA: OR = 0.86,


95% CI = 0.64–1.15,  $P = 0.305$ ). However, we could not calculate the heterogeneity for the AG vs. AA model in the “other” population subgroup because only one study remained after the outlier was excluded.

#### CYP1A1\*2A T/C (MspI) and Bladder Cancer Risk:


Three studies, including a total of 519 cases and 605 controls, evaluated the association between the MspI T/C variant and BC risk (Brockmoller et al., 1996; Li et al., 2007; Srivastava et al., 2008). However, though a slight trend was observed, we still failed to identify any significant association between the MspI T/C polymorphism and BC risk in all comparison models in the overall populations: i) C vs. T (OR = 1.24, 95% CI = 0.98–1.58,  $P = 0.078$ ); ii) CC vs. TT (OR = 1.24, 95% CI = 0.98–1.58,  $P = 0.078$ ); iii) TC vs. TT (OR = 1.19, 95% CI = 0.89–1.60,  $P = 0.240$ ); iv) CC+TC vs. TT (OR = 1.24, 95% CI = 0.93–1.64,  $P = 0.140$ ); v) CC vs. TC+TT (OR = 1.67, 95% CI = 0.84–3.34,  $P = 0.145$ ). No subgroup analysis was conducted in this gene due to the limited number of studies. Because no significant heterogeneity was found among the studies, the fixed-effects model was used to pool the data, and meta-regression and Galbraith plot analyses were not carried out.

#### *Sensitivity Analysis*

Sensitivity tests for both polymorphisms showed that no single study greatly influenced the estimates of overall risk by using the leave-one-out analysis and recalculating the ORs and the 95% CIs. Although genotype frequencies



**Figure 3. Galbraith Plots of CYP1A1\*2B A/G (Ile462Val) Polymorphisms and BC Risk in Different Genetic Models.** The study of Grando et al was outlier among the contrast: a. G vs. A; b. AG vs. AA; c. GG+AG vs. AA



**Figure 4. Funnel Plot Analysis to Detect Publication Bias.** Each point represents a separate study for the indicated association. a Funnel plot for contrast GG vs. AA; b Funnel plot for contrast CC vs. TT

in the two studies of Ile462Val A/G polymorphism deviated significantly from HWE (Fontana et al., 2009; Ozturk et al., 2011), the pooled ORs did not significantly change with or without the inclusion of these studies.

#### Publication Bias

To evaluate the possible publication bias, the Begg's funnel plot and Egger's test were performed. The funnel plots were symmetrical, suggesting that there was no significant publication bias among all the compared models (Figures 3 and 4). Egger's test, which statistically assesses the funnel plot symmetry, still did not reveal any potential publication bias, indicating that our results were relatively stable.

## Discussion

Environmental carcinogens, such as air pollution and tobacco smoke, are well-established sources of DNA damage, including oxidative base damage and abasic sites as well as single- and double-strand breaks (BosettiPiraLa Vecchia 2005). Through their metabolism in the organism, these risk factors can either become more carcinogenic or be detoxified. CYP1A1 is an important phase 1 enzyme involved in the biotransformation of tobacco procarcinogens (Guengerich et al., 1998), and according to Giri et al., CYP1A1 polymorphisms are modulator of genetic damage in occupational settings (Giri et al., 2012). Consequently, it might affect the detoxification process

of environmental carcinogens and lead to the alteration of an individual's BC susceptibility.

In this study, we analyzed the data from eight case-control studies whose results are conflicting regarding the association between CYP1A1 polymorphisms and BC susceptibility. The majority of the eligible studies included in our meta-analysis observed no significant association between CYP1A1 mutations and BC risk. On the other hand, Srivastava et al. (Srivastava et al., 2008) found a slight association with CYP1A1 (2A\* T/C) for risk of BC (OR = 1.56), although it was not statistically significant ( $P = 0.093$ ). In fact, there is evidence that CYP1A1 is involved in the carcinogenic progress of environmentally induced BC. A study conducted in human BC by Murai et al. (Murray et al., 1995) demonstrated an increased expression of CYP1A1 using immunohistochemistry. An in vitro investigation carried out by Wolf et al. (Wolf et al., 2005) found that cells treated with benzo[a]pyrene showed a dramatic increase in the expression of CYP1A1. Similar conclusions were also found in other experimental animals whose bladder cells were exposed to tobacco containing compounds such as benzo[a]pyrene and arylamines (Kogevinas et al., 2003; Dorrenhaus et al., 2007). However, we failed to reveal any evidence of an association between the two CYP1A1 polymorphisms and BC susceptibility overall, regardless of the additive, dominant, or recessive genetic models.

It has been reported that the distribution of the Val allele of Ile462Val varies extensively between different races, with the highest rate among East Asians (25%) and rarest among Caucasians (5%); the frequency of homozygous CYP1A1 MspI has also been found to be significantly different between various ethnicities (Cascorbi et al., 1996; Ji et al., 2012). Therefore, a further meta-analysis was performed among subgroups stratified by ethnicity. However, our result still failed to identify any significant association in this subgroup analysis.

Taken together, it may be concluded that CYP1A1 polymorphisms are not associated with BC risk in the overall population. Nevertheless, these results should be taken with caution as there is increasing evidence that metabolizing enzymes do not act alone and that single polymorphisms cannot explain complex multifactorial malignancies (Manuguerra et al., 2007). BC is a multifactorial disease that develops gradually following complex interactions among many genetic, environmental factors and other disease such as Diabetes Mellitus (Pollard et al., 2010; Botelho et al., 2010; Yang et al., 2013). The analysis of various combinations of genotypes to predict the risk of BC is now urgently required, since few studies have been performed studying double or triple polymorphisms of CYP1A1 combined with other genes to explore their association with various malignancies.

Data provided by Grando et al. (Grando et al., 2009) suggest that the GSTT1 null genotype and the A/G and G/G variants of the CYP1A1 genes may increase the risk for BC. A study by Öztürk et al. (Ozturk et al., 2011) also found that the CYP1A1 Ile/Ile (A/A) genotype combined with GSTM1 null genotype may contribute to the development of BC in the Turkish population, and that the CYP1A1Val genotype with GSTM1 null genotype

combinations, though not directly associated with risk of BC, is associated with higher grade tumors. However, due to the limited number of studies, meta-analyses on gene-gene combinations and BC risk were not conducted.

To summarize, this is the first meta-analysis assessing whether polymorphisms in CYP1A1 are associated with BC risk. A total of nine separate comparisons consisting of 1,059 BC cases and 1,061 controls were included, leading to a greater statistical power than for a single study. Our results were consistent with the majority of the eligible studies that indicated a non-significant association between CYP1A1 polymorphisms and BC risk. Nevertheless, some limitations to this meta-analysis should be considered. Firstly, there were only eight studies and therefore the limited sample size did not represent the global population. Thus, the null association between CYP1A1 mutations and BC risk may be due to the small sample size as it provides low statistical power. As is universally known, smoking is an important risk factor for BC, smokers with the CYP1A1 A/G polymorphism have been reported to have more PAH-DNA adducts than smokers without the polymorphism (Mooney et al., 1997), and some studies have found cigarette smoking to be strongly associated with tumor grade of BC (Jiang et al., 2012). However, only two studies included in our meta-analysis presented detailed frequencies of CYP1A1 polymorphisms with smoking habit. Therefore, a further analysis to clarify the effect of smoking behavior was not conducted. BC is a multifactorial disease, other risk factors such as occupational exposure and chronic irritation may modify the relationship between CYP1A1 polymorphisms and BC risk; however, these factors were not taken into consideration. Finally, a potential publication bias may exist because no effort was made to obtain unpublished studies, although the results of the Begg's funnel plot and Egger's test showed no evidence of publication bias among all comparison models.

In conclusion, our meta-analysis indicates that there is no relationship between CYP1A1\*2B A/G, 2A\* T/C polymorphisms and BC risk. Nevertheless, we have to emphasize the need for larger studies with a more rigorous design, especially studies taking the gene-gene and gene-environment interactions into consideration. In this way, the limitations mentioned above could be overcome and a more precise conclusion on the association between CYP1A1 polymorphisms and BC risk could thus be drawn.

## Acknowledgements

No funding was provided for the analysis.

## References

- Bartsch H, Nair U, Risch A, et al (2000). Genetic polymorphism of CYP genes, alone or in combination, as a risk modifier of tobacco-related cancers. *Cancer Epidemiol Biomarkers Prev*, **9**, 3-28.
- Berber U, Yilmaz I, Yilmaz O, et al (2013). CYP1A1 (Ile462Val), CYP1B1 (Ala119Ser and Val432Leu), GSTM1 (null), and GSTT1 (null) polymorphisms and bladder cancer risk in a Turkish population. *Asian Pac J Cancer Prev*, **14**, 3925-9.
- Boffetta P, Jourenkova N, Gustavsson P (1997). Cancer risk from occupational and environmental exposure to polycyclic aromatic hydrocarbons. *Cancer Causes Control*, **8**, 444-72.
- Bosetti C, Pira EL, Vecchia C (2005). Bladder cancer risk in painters: a review of the epidemiological evidence, 1989-2004. *Cancer Causes Control*, **16**, 997-1008.
- Botelho MC, Machado J, Cda Costa JM (2010). Schistosoma haematobium and bladder cancer: what lies beneath? *Virulence*, **1**, 84-7.
- Brockmoller J, Cascorbi I, Kerb R, et al (1996). Combined analysis of inherited polymorphisms in arylamine N-acetyltransferase 2, glutathione S-transferases M1 and T1, microsomal epoxide hydrolase, and cytochrome P450 enzymes as modulators of bladder cancer risk. *Cancer Res*, **56**, 3915-25.
- Cascorbi I, Brockmoller J, Roots I (1996). A C4887A polymorphism in exon 7 of human CYP1A1: population frequency, mutation linkages, and impact on lung cancer susceptibility. *Cancer Res*, **56**, 4965-9.
- Dai LP, Wang YP, Wu XB, et al (2009). Study on the association of cytochrome P450 polymorphisms and the risk of esophageal cancer: a meta-analysis. *Zhonghua Liu Xing Bing Xue Za Zhi*, **30**, 1198-202 (in Chinese).
- Dorrenhaus A, Muller TR, Roos PH (2007). Increased CYP1A1 expression in human exfoliated urothelial cells of cigarette smokers compared to non-smokers. *Arch Toxicol*, **81**, 19-25.
- Fontana L, Delort L, Joumard L, et al (2009). Genetic polymorphisms in CYP1A1, CYP1B1, COMT, GSTP1 and NAT2 genes and association with bladder cancer risk in a French cohort. *Anticancer Res*, **29**, 1631-5.
- Fu Jing CB-c (2013). Relationship between genetic polymorphisms of CYP1A1, NQO1, and EPHX1 and susceptibility to bladder cancer. *Chin J Cancer Prev Treat*, **20**, 10-4.
- Giri SK, Yadav A, Kumar A, et al (2012). CYP1A1 gene polymorphisms: modulator of genetic damage in coal-tar workers. *Asian Pac J Cancer Prev*, **13**, 3409-16.
- Grando JP, Kuasne H, Losi-Guembarovski R, et al (2009). Association between polymorphisms in the biometabolism genes CYP1A1, GSTM1, GSTT1 and GSTP1 in bladder cancer. *Clin Exp Med*, **9**, 21-8.
- Guengerich FP, Shimada T (1998). Activation of procarcinogens by human cytochrome P450 enzymes. *Mutat Res*, **400**, 201-13.
- Hayashi S, Watanabe J, Nakachi K, et al (1991). Genetic linkage of lung cancer-associated MspI polymorphisms with amino acid replacement in the heme binding region of the human cytochrome P450IA1 gene. *J Biochem*, **110**, 407-11.
- Hiyama T, Yoshihara M, Tanaka S, et al (2008). Genetic polymorphisms and head and neck cancer risk (Review). *Int J Oncol*, **32**, 945-73.
- Ismail SI, Soubani M, Nimri JM, et al (2013). Cancer incidence in Jordan from 1996 to 2009--a comprehensive study. *Asian Pac J Cancer Prev*, **14**, 3527-34.
- Ji YN, Wang Q, Suo LJ (2012). CYP1A1 Ile462Val polymorphism contributes to lung cancer susceptibility among lung squamous carcinoma and smokers: a meta-analysis. *PLoS One*, **7**, e43397.
- Jiang X, Castela JE, Yuan JM, et al (2012). Cigarette smoking and subtypes of bladder cancer. *Int J Cancer*, **130**, 896-901.
- Karbakhsh M, Dabbagh N, Shabani A, et al (2013). Age at diagnosis in bladder cancer: does opium addiction play a role? *Asian Pac J Cancer Prev*, **14**, 4723-5.
- Kawajiri K, Eguchi H, Nakachi K, et al (1996). Association of CYP1A1 germ line polymorphisms with mutations of the p53 gene in lung cancer. *Cancer Res*, **56**, 72-6.

- Kellermann G, Shaw CRL, Luyten-Kellerman M (1973). Aryl hydrocarbon hydroxylase inducibility and bronchogenic carcinoma. *N Engl J Med*, **289**, 934-7.
- Kogevinas M, Mannetje A, Cordier S, et al (2003). Occupation and bladder cancer among men in Western Europe. *Cancer Causes Control*, **14**, 907-14.
- Li S, Peng Q, Chen Y, et al (2013). DNA repair gene XRCC1 polymorphisms, smoking, and bladder cancer risk: a meta-analysis. *PLoS One*, **8**, e73448.
- Li X, Cui WP, Bai SF (2007). Analysis of the polymorphism of CYP1A1-MSPI gene in patients with bladder cancer. *China Prac Med*, **2**, 3-5.
- Lu Y, Peng Q, Zeng Z, et al (2013). CYP2D6 phenotypes and Parkinson's disease risk: A meta-analysis. *J Neurol Sci*, **336**, 161-8.
- Luch A (2005). Nature and nurture - lessons from chemical carcinogenesis. *Nat Rev Cancer*, **5**, 113-25.
- Manuguerra M, Matullo G, Veglia F, et al (2007). Multi-factor dimensionality reduction applied to a large prospective investigation on gene-gene and gene-environment interactions. *Carcinogenesis*, **28**, 414-22.
- Mooney LA, Bell DA, Santella RM, et al (1997). Contribution of genetic and nutritional factors to DNA damage in heavy smokers. *Carcinogenesis*, **18**, 503-9.
- Murray GI, Taylor VE, McKay JA, et al (1995). Expression of xenobiotic metabolizing enzymes in tumours of the urinary bladder. *Int J Exp Pathol*, **76**, 271-6.
- Ozturk T, Kahraman OT, Toptas B, et al (2011). The effect of CYP1A1 and GSTM1 gene polymorphisms in bladder cancer development in a Turkish population. *In Vivo*, **25**, 663-8.
- Peng Q, Mo C, Tang W, et al (2013). DNA repair gene XRCC3 polymorphisms and bladder cancer risk: a meta-analysis. *Tumour Biol*, **35**, 1933-44.
- Pollard C, Smith SC, Theodorescu D (2010). Molecular genesis of non-muscle-invasive urothelial carcinoma (NMIUC). *Expert Rev Mol Med*, **12**, e10.
- Romkes M, White C, Johnson J, et al (1996). Expression of cytochrome P450 mRNA in human lung, head and neck tumours, and normal adjacent tissues. *Proc Am Assoc Cancer Res*, **37**, 105.
- Sergentanis TN, Economopoulos KP (2010). Four polymorphisms in cytochrome P450 1A1 (CYP1A1) gene and breast cancer risk: a meta-analysis. *Breast Cancer Res Treat*, **122**, 459-69.
- Shakhssalim N, Hosseini SY, Basiri A, et al (2010). Prominent bladder cancer risk factors in Iran. *Asian Pac J Cancer Prev*, **11**, 601-6.
- Shimada T (2006). Xenobiotic-metabolizing enzymes involved in activation and detoxification of carcinogenic polycyclic aromatic hydrocarbons. *Drug Metab Pharmacokinet*, **21**, 257-76.
- Srivastava DS, Mandhani A, Mittal RD (2008). Genetic polymorphisms of cytochrome P450 CYP1A1 (\*2A) and microsomal epoxide hydrolase gene, interactions with tobacco-users, and susceptibility to bladder cancer: a study from North India. *Arch Toxicol*, **82**, 633-9.
- Thakkinstian A, McElduff P, D'Este C, et al (2005). A method for meta-analysis of molecular association studies. *Stat Med*, **24**, 1291-306.
- Vineis P, Veglia F, Garte S, et al (2007). Genetic susceptibility according to three metabolic pathways in cancers of the lung and bladder and in myeloid leukemias in nonsmokers. *Ann Oncol*, **18**, 1230-42.
- Wahid M, Mahjabeen I, Baig RM, et al (2013). Expression of CYP1A1 and GSTP1 in human brain tumor tissues in Pakistan. *Asian Pac J Cancer Prev*, **14**, 7187-91.
- Wolf A, Kutz A, Plottner S, et al (2005). The effect of benzo (a) pyrene on porcine urinary bladder epithelial cells analyzed for the expression of selected genes and cellular toxicological endpoints. *Toxicology*, **207**, 255-69.
- Yang XQ, Xu C, Sun Y, et al (2013). Diabetes mellitus increases the risk of bladder cancer: an updated meta-analysis. *Asian Pac J Cancer Prev*, **14**, 2583-9.
- Zeegers MP, Tan FE, Dorant E, et al (2000). The impact of characteristics of cigarette smoking on urinary tract cancer risk: a meta-analysis of epidemiologic studies. *Cancer*, **89**, 630-9.