

RESEARCH ARTICLE

Construction of a Protein-Protein Interaction Network for Chronic Myelocytic Leukemia and Pathway Prediction of Molecular Complexes

Chao Zhou¹, Wen-Jing Teng², Jing Yang¹, Zhen-Bo Hu³, Cong-Cong Wang³, Bao-Ning Qin², Qing-Liang Lv³, Ze-Wang Liu², Chang-Gang Sun^{1*}

Abstract

Background: Chronic myelocytic leukemia is a disease that threatens both adults and children. Great progress has been achieved in treatment but protein-protein interaction networks underlining chronic myelocytic leukemia are less known. **Objective:** To develop a protein-protein interaction network for chronic myelocytic leukemia based on gene expression and to predict biological pathways underlying molecular complexes in the network. **Materials and Methods:** Genes involved in chronic myelocytic leukemia were selected from OMIM database. Literature mining was performed by Agilent Literature Search plugin and a protein-protein interaction network of chronic myelocytic leukemia was established by Cytoscape. The molecular complexes in the network were detected by Clusterviz plugin and pathway enrichment of molecular complexes were performed by DAVID online. **Results and Discussion:** There are seventy-nine chronic myelocytic leukemia genes in the Mendelian Inheritance In Man Database. The protein-protein interaction network of chronic myelocytic leukemia contained 638 nodes, 1830 edges and perhaps 5 molecular complexes. Among them, complex 1 is involved in pathways that are related to cytokine secretion, cytokine-receptor binding, cytokine receptor signaling, while complex 3 is related to biological behavior of tumors which can provide the bioinformatic foundation for further understanding the mechanisms of chronic myelocytic leukemia.

Keywords: Protein-protein interaction networks - chronic myelocytic leukemia - molecular complexes - pathways

Asian Pac J Cancer Prev, 15 (13), 5325-5330

Introduction

Chronic myelogenous leukemia (CML) is a kind of chronic myeloproliferative disorders of primitive pluripotent hematopoietic stem cells. The special cytogenetic features of Chronic myelogenous leukemia cells is Philadelphia chromosome t(9; 22)(q34; q11). The proto-oncogene c-ABL in the long arm of Chromosome 9 translocates to the breakpoint cluster pool (BCR) of the long arm of chromosome 22, forming fusion gene BCR-ABL. Which encodes a protein having a strong activity of tyrosinase. The tyrosine kinase pathway can be phosphorylated by a variety of genes such as RAS, myc, c-CBL, phthalocyanine inositol phospholipid 3-kinase, and NF- κ B. Downstream of tyrosinase stimulation is the expression of oncogenes including c-fos, c-jun, etc. which stimulate cell non-growth factor-dependent proliferation and block apoptosis eventually leading to uncontrolled cell proliferation. BCR-ABL gene is considered as the molecular basis of the pathogenesis of CML and as an effective indicator diagnosis, efficacy, prognosis. The gene targeted drugs for the treatment of leukemia has

played a good effect but can not be cured, and gradually develop resistance. It requires our understanding of CML systematically to find more key genes and signaling pathways, and research on multi-target therapy or multi-drug combination therapy can bring a hope for CML cure..

Recently, systems biology approaches such as network-based methods have been successfully applied to elucidate the mechanism of diseases (Ideker et al, 2008; Zhao et al., 2008). From the systematic perspective, analysis of CML related bio-molecular interaction networks will improve the understanding of the complexity of molecular pathways underlying CML and will help to uncover the dynamic processes of disease progression.

In this study, we screened the confirmed disease-related genes using online Mendelian Inheritance in Man database (OMIM), created a protein-protein interaction networks based on biological function using Cytoscape software, detected molecule complex which may be included in the network, and conducted prediction of the relevant pathways that may be involved. We provided a basis for further explaining the mechanism of chronic myeloid leukemia development.

¹WeiFang Traditional Chinese Hospital, ²Shandong University of Chinese Traditional Medicine, ³Wei Fang Medical University *For correspondence: zhongliuyike@163.com

Materials and Methods

Design: Study on enrichment of genomic biological pathways.

Data acquisition

OMIM is a comprehensive, authoritative, daily updated human phenotype database, containing more than 12 000 genes of all human genetic diseases, and mainly focusing on hereditary diseases. Text messages, related reference information, sequence records, maps, and related databases are available for each gene. (Hamosh et al., 2005; Amberger et al., 2009).

The construction of gene/protein interaction networks

Chronic myelogenous leukemia associated genes were submitted to Cytoscape 2.8.2 plug-in Agilent Literature Search 2.7.7 (USA Agilent Technologies company) and Pubmed (Vailaya et al., 2005). False positive interaction information was removed from retrieval results. Then, gene/protein interaction relations were read in Cytoscape 2.8.2 and visualized (Shannon et al., 2003).

Network analysis

MCOMD algorithm in Cytoscape 2.8.2 web analytics plug-in Clusterviz of 1.2 was administrated to make the correlation analysis for the area of the construction of biological networks (Rintaro et al., 2012). By analyzing the network structure, proteins were grouped to form molecular compounds in the entire network and were shown in Cytoscape according to the correlation integral value. The areas with integral value higher than 3 were regarded as molecular compounds. The gene/protein names contained in the molecular compounds were submitted to The Database for Annotation, Visualization and Integrated Discovery (Huang et al., 2009). By retrieving Kyoto Encyclopedia of Genes and Genomes (KEGG) Database, biological pathways involved in chronic myelogenous leukaemia heredity were identified.

Main outcome measures

Protein networks were constructed based on the chronic myelogenous leukemia -related genes, nodes (proteins) and edges (interaction between), molecular complexes in the network and its associated interaction points and nodes (protein) and the edge (interaction between), analyze the biological pathways involved in the molecular complexes.

Results

Chronic myeloid leukemia related genes in OMIM Through the OMIM database retrieval, we found that 79 genes that were reported to be genes were identified: ABL1, ABL2, AKT1, ANPEP, APAF1, AQP5, ASXL1, AURKA, BAX, BCL11b, Bcl2, BCL2L1, BCR, CBL, CBLB, CD27, CDH13, CDKN1A, DOK2, EBAG9, EXOSC5, Fbxw7, FLT3, FN1, FUT4, FYN, H2afx, HMGA2, HOXA10, HOXA11, HRAS, IDH1, IDH2, IFNA1, Il6, Irf4, Irf8, JAK2, JUNB, KBTBD11, KIT, Lcn2, LOC425684, LOC514358, LOC556890, LOC560982, LYN, MAP2K7,

Map3k1, MCL1, MIR21, MIR328, MME, MPL, MTHFR, MUC1, Myc, NEWENTRY, NFKB1, NTRK1, NUDCD1, PCBP2, PDCD5, PTPN6, RALA, RBM15, RPS6KB1, SIGLEC8, SIRT1, SOCS3, SOCS6, SPHK1, STAT1, STAT3, STAT5A, TET2, TP53, USP9X, Zfp423.

Protein interaction networks

Through text mining 79 genetic-related genes showed that there was a network diagram with 638 nodes (proteins) and 1830 edges. As shown in Figure 1, the triangles represented OMIM genetic disease related proteins, while the diamonds represented the proteins obtained from text mining.

Network topology attribute analysis

Network topology attribute analysis shows that the connectivity of nodes in the network (the number of nodes in the network) obeys descending distribution, i.e. with the increase of edges connected to the node, correspondingly the number of nodes decrease, so it can be seen that the gene - protein interaction networks are scale-free networks (Burkard et al., 2010). We found that the connectivity of nodes in the network greater than or equal to 20 corresponds to a sharp reduction in the number of nodes. Therefore, we regarded the nodes which the connectivity is greater than / equal to 20 as the key nodes (hub). Key nodes included: TP53, AKT1, MAPK8, STAT3, EPHB2, MYC, TNFSF13, PIK3CA, JUN, JAK2, MAPK3, SRC, ABL1, ERVK2, IL6, MAPK14, FRAP1, BCL2L1, HCC, PCBP2, TNFSF10, BCR, CD4, CDKN1A, PARP1 etc.

Comparison (The horizontal axis represents betweenness, and the ordinate represents the connectivity degree. And the graphic in the table represents each node in the network). As can be seen that the connectivity (the number of nodes in the network) of nodes in the network obeys descending distribution, while the connectivity is greater than /equal to 20, the number of nodes corresponding to a sharp decrease.




Figure 1. Network Map of Chronic Myeloid Leukemia Protein Interaction (Overall + partial)




Figure 2. Connectivity Degree of Each node Degree and betweenness Comparison.

Table 1. Enrichment of Pathways Related to Molecular Complexes

Complex 1 Category	Term	P Value	Genes	FDR
BIOCARTA	h_inflamPathway:Cytokines and Inflammatory Response	3.44E-06	CSF3, IFNA1, IL6, IFNB1, IFNG, IL1A	0.003662
KEGG_PATHWAY	hsa04060:Cytokine-cytokine receptor interaction	1.74E-05	CSF3, IFNA1, IL6, IFNB1, IFNG, TGFB2, IL1A, CXCL10	0.017515
KEGG_PATHWAY	hsa04620:Toll-like receptor signaling pathway	1.95E-05	IFNA1, IL6, IFNB1, NFKBIA, TLR3, CXCL10	0.019171
KEGG_PATHWAY	hsa04623:Cytosolic DNA-sensing pathway	3.35E-05	IFNA1, IL6, IFNB1, NFKBIA, CXCL10	0.033809
KEGG_PATHWAY	hsa04623:Cytosolic DNA-sensing pathway	3.35E-05	IFNA1, IL6, IFNB1, NFKBIA, CXCL10	0.033809
BIOCARTA	h_il1rPathway:Signal transduction through IL1R	1.57E-04	IFNA1, IL6, IFNB1, NFKBIA, IL1A	0.166774
Complex3				
Category	Term	P Value	Genes	FDR
KEGG_PATHWAY	hsa04660:T cell receptor signaling pathway	2.12E-08	AKT1, NRAS, HRAS, KRAS, MAPK14, JUN, CBL, CD4, MAP2K7, IL10	2.22E-05
KEGG_PATHWAY	hsa04012:ErbB signaling pathway	6.70E-08	AKT1, NRAS, HRAS, PTK2, KRAS, JUN, CBL, MAP2K7, SRC	7.01E-05
KEGG_PATHWAY	hsa05215:Prostate cancer	8.02E-08	E2F1, AKT1, NRAS, HRAS, CCND1, HSP90AA1, KRAS, CREB1, PTEN	8.40E-05
KEGG_PATHWAY	hsa05220:Chronic myeloid leukemia	4.47E-07	E2F1, AKT1, NRAS, HRAS, CCND1, KRAS, CBL, BCL2L1	4.68E-04
KEGG_PATHWAY	hsa05214:Glioma	2.91E-06	E2F1, AKT1, NRAS, HRAS, CCND1, KRAS, PTEN	0.003048
PANTHER_PATHWAY	P04393:Ras Pathway	5.45E-06	AKT1, NRAS, HRAS, KRAS, MAPK14, JUN, STAT1, MAP2K7	0.004987
KEGG_PATHWAY	hsa04630:Jak-STAT signaling pathway	5.76E-06	STAT6, AKT1, CCND1, IL2RA, CBL, BCL2L1, STAT1, IL10, EPO	0.006026
KEGG_PATHWAY	hsa05218:Melanoma	5.91E-06	E2F1, AKT1, NRAS, HRAS, CCND1, KRAS, PTEN	0.006185
KEGG_PATHWAY	hsa05219:Bladder cancer	7.02E-06	E2F1, NRAS, HRAS, CCND1, FGFR3, KRAS	0.007344
PANTHER_PATHWAY	P04398:p53 pathway feedback loops 2	8.11E-06	AKT1, NRAS, HRAS, KRAS, MAPK14, STAT1, PTEN	0.007423
KEGG_PATHWAY	hsa04370:VEGF signaling pathway	8.15E-06	AKT1, NRAS, HRAS, PTK2, KRAS, MAPK14, SRC	0.008532
PANTHER_PATHWAY	P00018:EGF receptor signaling pathway	1.09E-05	STAT6, AKT1, NRAS, HRAS, KRAS, MAPK14, CBL, STAT1, MAP2K7	0.009929
REACTOME_PATHWAY	REACT_18266:Axon guidance	1.40E-05	NRAS, HRAS, PTK2, KRAS, CREB1, SRC	0.011615
KEGG_PATHWAY	hsa05222:Small cell lung cancer	1.58E-05	E2F1, AKT1, CCND1, PTK2, BCL2L1, PTEN, ITGA2B	0.0165
KEGG_PATHWAY	hsa05213:Endometrial cancer	2.04E-05	AKT1, NRAS, HRAS, CCND1, KRAS, PTEN	0.021389
KEGG_PATHWAY	hsa05223:Non-small cell lung cancer	2.46E-05	E2F1, AKT1, NRAS, HRAS, CCND1, KRAS	0.025779
PANTHER_PATHWAY	P00036:Interleukin signaling pathway	2.82E-05	STAT6, AKT1, NRAS, HRAS, IL2RA, KRAS, FOXP3, STAT1, SRC, IL10	0.025805
KEGG_PATHWAY	hsa04912:GnRH signaling pathway	3.82E-05	NRAS, HRAS, KRAS, MAPK14, JUN, MAP2K7, SRC	0.039968
PANTHER_PATHWAY	P00005:Angiogenesis	4.09E-05	AKT1, NRAS, HRAS, PTK2, FGFR3, KRAS, MAPK14, JUN, STAT1, SRC	0.037446
REACTOME_PATHWAY	REACT_16888:Signaling by PDGF	5.24E-05	STAT6, NRAS, HRAS, KRAS, STAT1, SRC	0.04343
PANTHER_PATHWAY	P00031:Inflammation mediated by chemokine and cytokine signaling pathway	8.07E-05	STAT6, AKT1, NRAS, HRAS, PTK2, KRAS, JUN, JUN, MYH14, STAT1, PTEN	0.073851
KEGG_PATHWAY	hsa05212:Pancreatic cancer	1.00E-04	E2F1, AKT1, CCND1, KRAS, BCL2L1, STAT1	0.10462
REACTOME_PATHWAY	REACT_11061:Signaling by NGF	1.07E-04	AKT1, NRAS, HRAS, KRAS, MAPK14, CREB1, PTEN, SRC	0.088951
KEGG_PATHWAY	hsa04722:Neurotrophin signaling pathway	1.43E-04	AKT1, NRAS, HRAS, KRAS, MAPK14, JUN, MAP2K7	0.149705
BIOCARTA	h_il2rPathway:IL-2 Receptor Beta Chain in T cell Activation	1.46E-04	E2F1, AKT1, HRAS, IL2RA, CBL, BCL2L1	0.173676
KEGG_PATHWAY	hsa04664:Fc epsilon RI signaling pathway	1.47E-04	AKT1, NRAS, HRAS, KRAS, MAPK14, MAP2K7	0.153361
KEGG_PATHWAY	hsa04530:Tight junction	2.19E-04	AKT1, NRAS, HRAS, KRAS, MYH14, PTEN, SRC	0.229339
REACTOME_PATHWAY	REACT_9417:Signaling by EGFR	2.35E-04	NRAS, HRAS, KRAS, CBL, SRC	0.194347
PANTHER_PATHWAY	P00048:PI3 kinase pathway	2.77E-04	AKT1, NRAS, HRAS, CCND1, KRAS, FOXP3, PTEN	0.253426
KEGG_PATHWAY	hsa04010:MAPK signaling pathway	2.86E-04	AKT1, NRAS, HRAS, FGFR3, KRAS, MAPK14, JUN, JUN, MAP2K7	0.298882
KEGG_PATHWAY	hsa04510:Focal adhesion	2.94E-04	AKT1, HRAS, CCND1, PTK2, JUN, PTEN, SRC, ITGA2B	0.307381
PANTHER_PATHWAY	P00056:VEGF signaling pathway	3.77E-04	AKT1, NRAS, HRAS, PTK2, KRAS, MAPK14	0.344702
KEGG_PATHWAY	hsa05221:Acute myeloid leukemia	5.48E-04	AKT1, NRAS, HRAS, CCND1, KRAS	0.571819
PANTHER_PATHWAY	P00021:FGF signaling pathway	6.37E-04	AKT1, NRAS, HRAS, FGFR3, KRAS, MAPK14, MAP2K7	0.581001
KEGG_PATHWAY	hsa05216:Thyroid cancer	8.86E-04	NRAS, HRAS, CCND1, KRAS	0.923857




Figure 3. Molecular Complexes Obtained by MCOMD Algorithm Analysis

The detection of molecular complexes

MCOMD algorithm analysis, there is a total of five molecular complexes whose correlation integral values are higher than 3.

Molecular complex pathway enrichment

Submit the 5 names of protein molecule complexes online to obtain the relevant passage in Table 1.

Discussion

In recent years, scientists have done a lot of studies of molecular mechanisms to CML: K Sailaja etc. suggested that the CYP3A5*3 gene polymorphism is significantly associated with the risk of CML development and disease progression (K Sailaja et al., 2010), Gulzar etc. proved that the presence of GSTT1 genotype may have a protective role against the CML (Gulzar et al., 2012), Khaldon Bodoor etc. highlighted an essential role of DAPK methylation in chronic leukemia in contrast to p15 methylation in the acute cases (Khaldon et al., 2014). Previous studies mainly used regression/variance analysis without considering the entire biological network analysis and targeted therapy prediction. Our results provide new understanding of CML pathogenesis and prognosis with the hope of offering theoretical support for future therapeutic studies.

Biomolecular network analysis is an important direction of systematic biology research. Understanding the complexity of molecular networks is an important challenge to the post-genomic era. The function of biomolecular function is often dependent on modularization, the network module which is made by a number of nodes in conjunction with each other and has a stable structure which can often reflect a similar nature between nodes (Chen et al., 2010; Kovács et al., 2010).

Analyzing the function module is the most common method to analyze biological molecular network. For example, Jin Gu etc. used gene network modules response to study inflammation and angiogenesis, and had predicted that ADAM17, CD40, ETS1, FOXO1, SMAD3 and TLR4 are the target genes of mir-145 (Jin et al., 2010), Francisco Azuaje and others have found that ITGB1 and FN1 are potential biomarkers associated with heart failure through analyzing related protein network module (Francisco et al., 2010), Zhu XL etc. have used the Weighted Gene Co-expression Network Analysis method to identify prognostic markers for EC and characterized underlying molecular mechanisms by KEGG pathway enrichment and transcriptional regulation analyses (Zhu et al., 2012). Feng DQ etc. have characterized different KEGG pathways and GO terms by altered gene expression with complex pathway analysis, and identified potentially subjected to miRNA regulation (Feng et al., 2012). It can be seen, that the analysis of functional module can reflect the overall web properties of biological molecules.

There are three main categories to construct Biomolecular regulatory network: constructing gene regulatory networks through a mathematical model; building a network through literature mining and integrating multiple data (Friedman et al., 2000; Hwang et al., 2005; Mohamed et al., 2009). Liu Y etc. built a PPI network using the nearest neighbor expansion method from existing database, and analyzed the key protein nodes (Liu et al., 2013). But in our experiment we built network with the method of literature mining, then analyzed modules to draw relevant signaling pathways and hub proteins, the new method and multi-angles deepened disease research. Building a network through literature mining means using bioinformatics, computational biology and other fields of computer science to analyze the knowledge of the literatures, and build biomolecules regulatory networks by using the relationship between gene/protein interaction of the existing literatures. The advantage of this method is that establishing the relationship between the regulation accurately and the network stably.

Therefore, in this paper we use the OMIM genes information and literature mining constructs to construct the chronic myeloid leukemia protein interaction networks. Although, the data of this experiment is genetic genes, but to elaborate and prove its functionality as well as the relationship with other molecules is mostly at the protein level in the literatures. For this reason, this network constructed combining with literature data mining should be a protein interaction network. In Cytoscape, each protein is represented as a node, and each interaction between two nodes is represented as an edge. To increase the liabilities and define the protein perturbation to a certain level, the network reconstruction was limited to the first protein neighbors (Wu et al., 2014).

According to 79 genes provided by OMIM, this study has built chronic myelogenous leukemia protein interaction network, containing 638 nodes (proteins), 1830 edge (interaction). Can the network describe molecular regulation of chronic myelogenous leukemia in the development? According to the existing literature (Biethahn et al., 1999; Reddy et al., 2000; Reilly et al.,

2002;Margaret, 2002; Bader et al., 2003;Donald et al., 2005; Jayaraman et al., 2012) BCR, jun is the main reason of chronic myelogenous leukemia genes; as a cancer gene, myc is downstream signaling molecules of BCR ABL fusion protein, involved in the pathogenesis of chronic myelogenous leukemia; Number-growing evidence suggests the role of Stat protein family in the initiation and development of leukemia, and the activation of the Stat proteins are associated with poor prognosis and shortened survival in patients; CD4's immune inhibitory effect make cancer gene to escape immune surveillance; IL - 6 is a pleiotropic cytokine that can regulate a variety of cellular functions: including cell proliferation, cell differentiation, immune defense mechanisms and blood cell production, etc.; the abnormalities of tumor suppressor gene p53 and oncogene myc is associated with rapid change of chronic myelogenous leukemia ;the mutation of JAK2 V617F in chronic myeloid leukemia sex is rare, and CML patients with this mutation have a longer survival, JAK2 V617F mutation may indicate a good prognosis; Experiments which use the radiolabeled molecules LymphoRad131 TNFSF13B to assist cancer chemotherapy has entered clinical trials; the inordinate expression of apoptosis-related gene bcl-x, anti-apoptotic gene akt1, proto-oncogene scr in time and space make their protein product change in the quality and quantity , and this is an important cause of tumorigenesis. These are relational pathogenesis of chronic myelogenous leukemia, We constructed a network which comprise these genes or proteins , this proves that genes - protein interaction network we constructed has a certain authenticity that can be used to describe the interaction of chronic myeloid leukemia in molecules.

As the network is very large, the experimental introduced MCOMD algorithm to evaluate the network's regional integration through the correlation integral. Correlation integral is described of proteins associated with the degree within the region. Proteins of the same molecular complex generally have the same biological function, so we can discover unknown gene functions or new molecular functional groups. Research shows 5 molecular complexes that is more than 3 in correlation integral. This knowledge DAVID is not only extensive genes annotation in different species, but also enriched to biological information of a single gene. The Protein molecule biological pathways of complexes 2, 4, 5 are not exist, There are two reasons speculated: Although the relevance of these molecular complexes correlation integral is higher, but it can not prove that it contains a protein with similar biological functions; Existing studies have not revealed their biological pathways involved. Molecular complex 1, 3 involved many biological pathways, Figure 3 shows its complexity, Where there are 6 biological pathways whose FDR < 1 in molecular complex 1, and 36 in molecular complex 3.

Molecular complex 1 can be predicted to be relative to complex regulation of cytokines, specifically cytokines and inflammation, cytokine receptor interaction and biological receptor signal transduction pathways. Existing literature indicates myeloproliferative diseases are highly sensitive to cytokine, these genes of signaling pathway

may provide research direction for molecular therapy of CML. Molecular complex 3 Shows the complex biological behavior of chronic myeloid leukemia and extensive correlation with other cancers. CML is demonstrated not simply controlled by a particular gene or signaling pathways, but by the complex process of network system coordinately regulated which consisted of a variety of signaling pathways and multiple genes. In the signaling network , it is likely there is some “ key regulatory point”.

The experiment dig out a variety of signaling pathways and genes, and with the application of protein network greatly expanded, such as transcriptome network analysis and pathway crosstalk analysis (Pan, 2012) , it can provide reliable directions for molecular mechanism research of CML treatment.

References

- Amberger J, Bocchini CA, Scott AF, et al (2009). McKusick's Online Mendelian Inheritance in Man (OMIM). *Nucleic Acids Res*, **37**, 793-6.
- Burkard TR, Rix U, Breitwieser FP, Superti-Furga G, Colinge J (2010). A computational approach to analyze the mechanism of action of the kinase inhibitor bafetinib. *PLoS Comput Biol*, **18**, 1001001.
- Bader GD, Hogue CW(2003). An automated method for finding molecular complexes in large protein interaction networks. *BMC Bioinformatics*, **4**, 2.
- Biethahn S, Alves F, Wilde S, et al (1999). Expression of granulocyte colony_stimulated factor and granulocyte macrophage colony stimulating factor associated signal transduction proteins of the JAK/ Stat passway in normal granulopoiesis and in blast cells of acute myelogenous leukemia. *Exp Hematol*, **27**, 885-94.
- Chen L, Wang H, Zhang L, et al (2010). Uncovering packaging features of co-regulated modules based on human protein interaction and transcriptional regulatory networks. *BMC Bioinformatics*, **7**, 392.
- Dennis G Jr, Sherman BT, Hosack DA, et al (2003). DAVID: database for annotation, visualization, and integrated discovery. *Genome Biol*, **4**, 3.
- Donald P, Mc Lornan, Melanie J (2005). Chronic neutrophilic leukemia with an associated V617F JAK2 tyrosine kinase mutation. *Haematologica*, **90**, 1696-7.
- Francisco Azuaje, Yvan Devaux, Daniel R Wagner (2010). Coordinated modular functionality and prognostic potential of a heart failure biomarker- driven interaction network. *BMC Systems Biology*, **4**, 60
- Friedman N, Linial M, Nachman I, et al (2000). Using Bayesian networks to analyze expression data. *J Comput Biol*, **7**, 601-20.
- Feng DQ, Huang B, Li J, et al (2013). Selective miRNA expression profile in chronic myeloid leukemia K562 cell-derived exosomes. *Asian Pac J Cancer Prev*, **14**, 7501-7.
- Gulzar Bhat, Ashaquallah Bhat, Aadil Wani, et al (2012). Polymorphic variation in GST genes and risk of CML in the kashmiri population. *Asian Pac J Cancer Prev*, **15**, 69-73.
- Hamosh A, Scott AF, Amberger JS, et al (2005). Online Mendelian Inheritance in Man (OMIM), a knowledgebase of human genes and genetic disorders. *Nucleic Acids Res*, **33**, 514-7.
- Huang da W, Sherman BT, Lempicki RA (2009). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc*, **4**, 44-57.
- Hwang D, Smith JJ, Leslie DM, et al (2005). A data integration

- methodology for systems biology: experimental verification. *Proc Natl Acad Sci USA*, **102**, 17302-7.
- Ideker T, Sharan R(2008). Protein networks in disease. *Genome Res*, **18**, 644-52.
- Kovács IA, Palotai R, Szalay MS, et al (2010). Community landscapes:an integrative approach to determine overlapping network module hierarchy, identify key nodes and predict network dynamics. *PLoS One*, **5**, 12528.
- Sailaja K, Surekha D, Nageswara Rao D, et al (2010). Polymorphisms in indian chronic myeloid leukemia patients. *Asian Pac J Cancer Prev*, **11**, 781-4.
- Khalid Bodoor, Yazan Haddad, Asem Alkhateeb, et al (2014). DNA hypermethylation of cell cycle and apoptotic genes in peripheral blood of leukemia patients. *Asian Pac J Cancer Prev*, **15**, 75-84.
- Liu Y, Liu CX, Wu ZT, et al (2013). Mining proteins associated with oral squamous cell carcinoma in complex networks. *Asian Pac J Cancer Prev*, **14**, 4621-5.
- Mohamed-Hussein ZA, Harun S (2009). Construction of a polycystic ovarian syndrome (PCOS) pathway based on the interactions of PCOS-related proteins retrieved from bibliomic data. *Theor Biol Med Model*, **6**, 18.
- Margaret CF (2002). Src in cancer:deregulation and consequences for cell behaviour. *Biochim Biophys Acta*, **1602**, 114-30.
- Pan XH (2012). Pathway crosstalk analysis based on protein-protein network analysis in ovarian cancer. *Asian Pac J Cancer Prev*, **13**, 3905-9.
- Reddy EP, Korapati A, Chaturvedi P, et al (2000). IL-3 signaling and the role of Src kinases, JAKs and STATs:a covert liaison unveiled. *Oncogene*, **19**, 2532-47.
- Reilly JT (2002). Chronic neutrophilic leukaemia: a distinct clinical Entity. *Br J Haematol*, **11**, 10-8.
- Rintaro Saito, Michael E Smoot, Keiichiro, et al (2012). A travel guide to Cytoscape plugins, *Nat Methods*, **9**, 1069-76.
- Shannon P, Markiel A, Ozier O, et al (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res*, **13**, 2498-504.
- Vailaya A, Bluvias P, Kincaid R, et al(2005). An architecture for biological information extraction and representation. *Bioinformatics*, **21**, 430-8.
- Wu BL (2014). Protein-protein Interaction network analyses for elucidating the roles of LOXL2-delta72 in esophageal squamous cell carcinoma. *Asian Pac J Cancer Prev*, **15**, 2345.
- Zhao XM, Wang RS, Chen L, Aihara K(2008). Uncovering signal transduction networks from high-throughput data by integer linear programming. *Nucleic Acids Res*, **36**, 48.
- Zhu XL, Ai ZH, Wang J, et al (2012). Weighted gene co-expression network analysis in identification of endometrial cancer prognosis markers. *Asian Pac J Cancer Prev*, **13**, 4607-11.