

RESEARCH ARTICLE

Flavonoids from *Orostachys Japonicus A. Berger* Induces Caspase-dependent Apoptosis at Least Partly through Activation of p38 MAPK Pathway in U937 Human Leukemic Cells

Won Sup Lee^{1&*}, Jeong Won Yun^{1&}, Arulkumar Nagappan¹, Ji Hyun Jung¹, Sang Mi Yi¹, Dong Hoon Kim², Hye Jung Kim³, GonSup Kim⁴, Chung Ho Ryu^{5&}, Sung Chul Shin⁶, Soon Chan Hong⁷, Yung Hyun Choi⁸, Jin-Myung Jung⁹

Abstract

Background: *Orostachys japonicus A. Berger (A. Berger)* is commonly used as a folk remedy for cancer therapy. However, the mechanisms of its anti-cancer activity are poorly investigated in human cancer cells. In this study, we investigated whether flavonoids extracted from *Orostachys japonicus A. Berger* (FEOJ) might have anticancer effects in human leukemia cells, focusing on cell death mechanisms. **Materials and Methods:** U937 human leukemic cancer cells were used. **Results:** FEOJ induced apoptosis in a dose-dependent manner in human U937 cancer cells. Flow cytometry revealed significant accumulation of cells with sub-G1 DNA content at the concentrations of 200 $\mu\text{g/mL}$ and 400 $\mu\text{g/mL}$. FEOJ-induced apoptosis was caspase-dependent through loss of mitochondrial membrane potential (MMP, $\Delta\Psi\text{m}$) in human U937 cancer cells, which might be associated with suppression of Bcl-2 and XIAP proteins. FEOJ induced the p38 MAPK signaling pathway, playing at least in part an important role in FEOJ-induced apoptosis. **Conclusions:** This study suggested that FEOJ may induce caspase-dependent apoptosis in human leukemic cells by regulating MMP ($\Delta\Psi\text{m}$) through suppressing Bcl-2 and X-IAP. In addition, the results indicated that upstream p38 MAPK signaling regulates the apoptotic effect of FEOJ. This study provides evidence that FEOJ might have anti-cancer potential for human leukemic cells.

Keywords: *Orostachys japonicus A. Berger* - flavonoids - U937 - apoptosis - p38 MAPK - leukemia cells

Asian Pac J Cancer Prev, 16 (2), 465-469

Introduction

Apoptosis is an active-energy requiring process (a type I programmed cell death) with a distinctive phenotype characterized by cytoplasmic shrinkage, blebbing of the plasma membrane, chromatin condensation, DNA degradation, and nuclear fragmentation (Kerr et al., 1972; Walker and Sikorska, 1997). Most of apoptosis occurs through two major pathways (intrinsic pathway - mitochondria-mediated apoptosis and extrinsic pathway - death receptor-mediated apoptosis) (Kumi-Diaka et al., 2004).

Despite enormous efforts to control cancer development, the cancer incidence is still increasing (Jung et al., 2014). Given the results, we obviously need a paradigm shift to reduce the cancer incidence. One of the pragmatic ways is chemoprevention because the

dietary agents and some phytochemicals are reported to induce apoptosis or enhance anti-cancer effects without significant toxicities in the normal cells (Palasap et al., 2014). This safely showing or enhancing anti-cancer activity derived from the dietary agents or phytochemicals has drawn our interests to seek the possibility of controlling cancer with minimal toxicity. The extracts of *Orostachys japonicus A. Berger (A. Berger)* has been used as a folk remedy for cancer treatment. *Orostachys japonicus A. Berger* belongs to a member of the family Liliaceae, and has plentiful flavonoids, which may exhibit anti-cancer properties. However, few studies have been conducted regarding the anti-cancer effects of flavonoids extracted from lyophilized *A. Berger* (FEOJ) and the molecular mechanisms of the effects are poorly understood in human cancer cells. In addition, with the growth of ecological movements, natural products have become more popular

Departments of ¹Internal Medicine, ²Emergency Medicine, ³Pharmacology, ⁷Surgery, ⁹Neurosurgery, Institute of Health Sciences, ⁵Division of Applied Life Science (BK 21 Program), Institute of Agriculture and Life Science, ⁶Department of Chemistry, Research Institute of Life Science, ⁴School of Veterinary Medicine, Gyeongsang National University School of Medicine, Jinju, ⁸Department of Biochemistry, Dongeui University College of Oriental Medicine, Busan, Korea *Equal contributors *For correspondence: lwshmo@hanmail.net, lwshmo@gsnu.ac.kr

for the prevention or treatment of cancer.

With this background information, we assumed that FEOJ are major ingredients have the anti-cancer effects of the extracts of *Orostachys japonicus* A. Berger (A. Berger) that has been used as a folk remedy for cancer treatment. Therefore, we isolated and fully characterized FEOJ, and investigated the mechanisms of anticancer effects of FEOJ on human leukemic cells.

Materials and Methods

Preparation of FEOJ

FEOJ isolated from the plants of *Orostachys japonicus* A. Berger was a generous gift from Dr. S.C. Shin (Department of Chemistry, Gyeongsang National University, Korea). FEOJ were extracted with polar organic solvents followed by RP-chromatography. FEOJ contains 16 flavonoids; (1): Procyanidin dimer gallate isomer; (2): Procyanidin dimer gallate isomer (3): Epigallocatechin-3-gallate (4): Procyanidin dimer digallate isomer (5): Epicatechin (6): Quercetin 3-O-rhamnosyl-7-O-glucoside (7): Epicatechin gallate (8): Kaempferol 3-O-rhamnosyl-7-O-glucoside (9): Procyanidin dimer digallate isomer (10): Myricetin 3-O-glucoside (11): Quercetin 3-O-glucoside (12): Kaempferol-3-O-rutinoside (13): Quercetin 7-O-rhamnoside (14): Kaempferol dideoxyhexosylhexoside (15): N1,N5,N10-tri-p-(E,E,E)-coumaroyl spermidine (16): Kaempferol.

Cells and reagents.

U937 human leukemic cells from the American type culture collection (Rockville, MD, USA) were cultured in RPMI 1640 medium (Invitrogen Corp, Carlsbad, CA, USA) supplemented with 10% (v/v) fetal bovine serum (FBS) (GIBCO BRL, Grand Island, NY, USA), 1 mM L-glutamine, 100 U/mL penicillin, and 100 µg/mL streptomycin at 37°C in a humidified atmosphere of 95% air and 5% CO₂. Antibodies against Bcl-2, Bid, inhibitors of apoptosis protein-1 (IAP-1), IAP-2, X-linked IAP (XIAP), pro-caspase 3, pro-caspase 8, and pro-caspase 9 were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Antibody against poly (ADP-ribose) polymerase (PARP) was purchased from PharMingen (San Diego, CA, USA). Antibody against β-actin was from Sigma (Beverly, MA). Peroxidase-labeled donkey anti-rabbit and sheep anti-mouse immunoglobulin, and an enhanced chemiluminescence (ECL) kit were purchased from Amersham (Arlington Heights, IL). Caspase activity assay kits were purchased from R&D systems (Minneapolis, MN, USA). All other chemicals not specifically cited here were purchased from Sigma Chemical Co. (St. Louis, MO, USA).

Cell viability

The cells were seeded onto 24-well plates at a concentration of 1×10⁵ cells/ml, and then treated with the indicated concentration of FEOJ for 48 h. 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT, 0.5 mg/ml) was subsequently added to each well. After additional 2 hr-incubation, 100 µL of a solution containing 10% SDS (pH 4.8) plus 0.01 N HCl was added

to dissolve the crystals. The absorption values at 570 nm were determined with an enzyme-linked immunosorbent assay (ELISA) plate reader.

Flow cytometry analysis for measurement of sub-G1 phase

The cells were plated at a concentration of 2×10⁵ cells/well in six-well plates. After treatment of FEOJ, the cells were collected, washed with cold PBS, and then centrifuged. The pellet was fixed in 75% (v/v) ethanol at 4°C for 1 h. The cells were washed once with PBS, and re-suspended in cold PI solution (50 µg/ml) containing RNase A (0.1 mg/ml) in PBS (pH 7.4) for 30 min in the dark. The cellular DNA was then stained by applying 250 µl of propidium iodide (50 mg/ml) for 30 min at room temperature. The stained cells were analyzed by fluorescent activated cell sorting (FACS) on a FACScan flow cytometer (Becton Dickinson, San Jose, CA) for relative DNA content based on red fluorescence.

Western blot analysis

The suspending cells were harvested and lysed in the lysis buffer. The cells were disrupted by sonication and extracted at 4°C for 30 min. The protein concentrations were quantified using the BioRad protein assay (BioRad Lab., Hercules, CA, U.S.A.). The proteins of the extracts were resolved by electrophoresis, electrotransferred to a polyvinylidene difluoride membrane (Millipore, Bedford, MA), and then the membrane was incubated with the primary antibodies followed by a conjugated secondary antibody to peroxidase. After washing, the membranes were developed by enhanced chemiluminescence.

Caspase activity assay.

Caspase activity was determined by a colorimetric assay according to the manufacturer's protocol in a kit for caspase activity (R&D Systems, Minneapolis, Minnesota, USA). In brief, the cells were lysed in the supplied lysis buffer. The supernatants were collected and incubated with the supplied reaction buffer containing dithiothreitol and substrates at 37°C. The reaction was measured by determining the change in absorbance at 405 nm using the microplate reader.

Mitochondrial membrane potential (MMP, ΔΨ_m) assay

The MMP (ΔΨ_m) in living cells was measured by flow cytometry with the lipophilic cationic probe JC-1, which is a ratiometric, dual-emission fluorescent dye. There are two excitation wavelengths, 527 nm for the monomer form and 590 nm for the J-aggregate form. The magnitude of green fluorescent signals reflects the amount of damaged mitochondria. The cells were trypsinized and the cell pellets were re-suspended in 500 µl of PBS, incubated with 10 µM JC-1 for 20 min at 37°C. The cells were subsequently washed once with cold PBS, suspended in a total volume of 500 µl, and analyzed by flow cytometry.

Statistical analyses

Data represent means ± standard deviations. Statistical significance was determined using the one-way analysis of variance (ANOVA) with post-test Neuman-Keuls for more than two groups and Student's t test for two groups.

Results

FEOJ induces apoptosis in U937 human leukemic cells

To investigate the anti-cancer activity of FEOJ, U937 cells were treated with various concentrations of FEOJ for 48 h. The cell growth was assessed by MTT assay. The MTT assay revealed that the growth of U937 cells were inhibited by FEOJ treatment in a dose-dependent manner, and the 50% inhibition of cell growth (IC₅₀) was approximately 180 µg/mL (Figure 1A). To determine whether the decrease in cell viability was derived from FEOJ-induced apoptosis, we measured cells with sub-G1 DNA content by flow cytometry. The cells with sub-G1 DNA content was increased in a dose-dependent (Figure 1B and 1C).




Figure 1. Effects of FEOJ on the Cell Viability of U937 Cells.

The cells were seeded at the density of 1×10^5 cells per ml and incubated with the indicated concentrations of FEOJ for 48h at the indicated concentrations of FEOJ. **A)** Cell viability was assessed by the MTT assay. Data are expressed as mean±SD of three independent experiments (*p<0.05 versus control). **B and C)** By flow cytometry, the cells with sub-G1 DNA content representing the cells undergoing cell death were also analyzed.




Figure 2. Activation of Caspases During the FEOJ-Induced Apoptosis in U937 Cells.


U937 cells were incubated at indicated concentrations of FEOJ for 48 h. **A)** Equal amounts of cell lysate (30µg) were resolved by SDS-polyacrylamide gels and transferred onto nitrocellulose membranes. The membranes were probed with the anti-caspase-3, -8, -9 and anti-PARP antibodies. The proteins were visualized using an ECL detection system. β-actin was used as an internal control. **B and C)** The cell lysates from the cells treated with FEOJ for 48 h were assayed for in vitro caspase-3, -8 and -9 activity using DEVD-pNA, IETD-pNA and LEHD-pNA respectively, as substrates. The released fluorescent products were measured. Each bar graph represents mean±SD of three independent experiments. *p<0.05 between the treated with FEOJ and the untreated group

Next, we investigated whether FEOJ-induced apoptosis is caspase dependent. Then we assessed the effects of FEOJ on caspases and their substrates (PARP). FEOJ decreased the expression levels of pro-caspase-3, pro-caspase-8, and pro-caspase-9 in a dose-dependent manner. With the decrease of pro-caspases, the cleavages of PARP were prominent at the concentration of 400 µg/mL of FEOJ (Figure 2A). These findings suggest that FEOJ induce apoptosis through caspase activation. To confirm this finding, we assessed proteolytic activities of caspases using colorimetric assay kits. Unfortunately, the caspase activity assay only showed that FEOJ increased proteolytic activities of caspases at the concentration of 400 µg/mL (Figure 2B). The caspase activity assay revealed that FEOJ increased proteolytic activities of caspases and induce caspase-dependent apoptosis.


FEOJ induces apoptosis at least partly through mitochondrial pathway and modulates Bcl-2 and IAP family members.

Mitochondria play a central role in apoptosis. As the early event of apoptosis, mitochondrial depolarization occurs. Therefore, we investigated the role of mitochondria in FEOJ-induced apoptosis by measuring MMP ($\Delta\Psi_m$). As shown in Figure 3A, 48 h-treatment of FEOJ significantly induced the loss of MMP ($\Delta\Psi_m$) in a dose-dependent manner (78.5% at 400µg/ml vs 37% at the

A)



B)



C)




Figure 3. Effects of FEOJ on Mitochondrial Depolarization, and Bcl-2, and IAP Family Members in U937 Cells.

The cells were treated with FEOJ for 48 h at the indicated concentrations. **(A)** The cells were stained with JC-1 and incubated at 37°C for 20 min. The mean JC-1 fluorescence intensity was detected using a flow cytometer. **(B and C)** Equal amounts of cell lysate (30µg) were resolved by SDS-polyacrylamide gels and transferred onto nitrocellulose membranes. The membranes were probed with the indicated antibodies and detected by the ECL system. The results are from a representative experiment of at least two independent experiments that showed similar patterns

control), indicating that FEOJ increases depolarization of the MMP ($\Delta\Psi_m$). The loss of MMP ($\Delta\Psi_m$) elicited by FEOJ lead to activation of caspase-3, which has been shown to play a pivotal role in the terminal execution phase of apoptosis, and also activate caspase 8 through the activation of Bid protein (Ashkenazi, 2002). Therefore, we checked Bid cleavages to verify the feedback pathway to the activation of caspase 8. Western blot data showed that FEOJ induced cleavage of Bid (activation of Bid protein) in a dose-dependent manner (Figure 3B). Next, we assessed the levels of Bcl-2 and IAP family proteins which play a crucial role in apoptosis to elucidate further underlying mechanisms of FEOJ-induced apoptosis. Western blotting revealed that FEOJ suppressed Bcl-2 and XIAP (anti-apoptotic proteins) in a dose-dependent manner (Figure 3B) whereas the suppression of CIAP-1 and CIAP-2 was not observed (Figure 3C). These findings suggest that the loss of MMP ($\Delta\Psi_m$) and activation of Bid protein are associated with down-regulation of Bcl-2 and X-IAP protein.

FEOJ induced apoptosis at least in part by activating p38 MAPK activity.

Mitogen-activated protein kinase (MAPK) (Robinson and Cobb, 1997) and PI3K/Akt pathway (Luo et al., 2003) play important roles in regulating apoptosis, differentiation and cell death. To determine whether Mitogen-activated protein kinase (MAPK) and PI3K/Akt pathway is associated with FEOJ-induced apoptosis, we investigated the effects of FEOJ on MAPK and Akt in U937 cells. Since the activity of MAPK and Akt is regulated by phosphorylation, we assessed the levels of phosphorylated MAPK and Akt in FEOJ-treated U937 cells. The Western blot analysis revealed that FEOJ significantly activated phosphorylation of p38 MAPK and suppressed the phosphorylation of Akt (Figure




Figure 4. The Role of p38 MAPK Signaling in FEOJ-Induced Apoptosis. The cells were seeded at the density of 1×10^5 cells per ml and incubated at the indicated concentrations of FEOJ for the indicated time duration. (A) Western blot analysis for Akt and p38 MAPK. Equal amounts of cell lysate (30 µg) were resolved by SDS-polyacrylamide gels and transferred onto nitrocellulose membranes. The results are from at least two independent experiments that showed similar patterns. (B) Cell viability was assessed by the MTT assay. Data are expressed as mean \pm SD of three independent experiments (* $p < 0.05$ versus control, ** $p < 0.05$ versus FEOJ treatment)




Figure 5. Schematic Representation on Apoptotic Effects of FEOJ on U937 Human Leukemic Cells.

FEOJ induced apoptosis through mitochondrial pathway by inducing loss of MMP ($\Delta\Psi_m$), and then activates caspases. During this process, the suppression of Bcl-2 and X-IAP contributes the augmentation of the apoptosis. In addition, p38 MAPK activation may serve as one of the upstream signaling involved in FEOJ-induced apoptosis. Taken together, this study suggested that FEOJ induced caspase-dependent apoptosis through mitochondrial pathway by modulating Bcl-2 and X-IAP proteins, and that p38 MAPK is at least one of the important upstream signalings that regulate the apoptotic effect of FEOJ in human leukemic U937 cells

4A). To confirm this finding, we evaluated the effects of FEOJ with and without treatment of Akt inhibitor (LY294002) and p38 MAPK inhibitor (SB203580). MTT assay revealed that LY294002 and p38 MAPK inhibitor (SB203580) reduced the effect of FEOJ on cell viability (Figure 4B). Taken together, these findings suggest that FEOJ induce apoptosis at least in part by activating p38 MAPK signaling pathway.

Discussion

This study was designed to determine whether FEOJ has anti-cancer properties in human leukemia cells, and further investigate the underlying mechanisms of the apoptotic effect of FEOJ. Indeed, we demonstrated that FEOJ induced apoptosis in a dose-dependent manner in U937 human leukemic cells. The FEOJ-induced apoptosis was caspase dependent. This finding was confirmed by caspase activity assay. Many well-known phytochemicals that can safely modulate physiological function and enhance anti-cancer activity induce apoptosis through mitochondrial pathway (Chun et al., 2003; Gupta et al., 2013). Similarly, FEOJ induced apoptosis through induction of loss of MMP ($\Delta\Psi_m$). The loss of MMP ($\Delta\Psi_m$) may be related with suppression of BCL-2. Next, we assessed IAP family members which are important anti-apoptotic proteins that regulate caspases. We demonstrated that FEOJ suppressed X-IAP in a dose-dependent manner, which can be associated with induction of apoptosis and activation of Bid protein. All these finding suggested that FEOJ induces caspase-dependent apoptosis at least in part through inhibiting anti-apoptotic protein expression.

When we look into the components of FEOJ, quercetin was one of the major components (Wiczowski et al., 2008; Lee and Mitchell, 2011). Previous reports demonstrated

Flavonoids from Orostachys Japonicus A. Berger Induce Caspase-dependent Apoptosis in U937 Human Leukemic Cells **2011**, 591356.

that quercetin induced apoptosis in several types of cancer cells (Psahoulia et al., 2007), and it augmented apoptosis by up-regulating DR5 expression (Jang et al., 2003). These findings are consistent with our results even though we did not demonstrate the up-regulation of DR5 by FEOJ in that FEOJ also induced caspase 8 activation. In addition, quercetin increases apoptosis by altering anti-apoptotic proteins, which supports our finding that FEOJ suppressed an anti-apoptotic factor Bcl-2, and X-IAP in U937 cells. Regarding cIAP-1, our data showed that FEOJ did not influence the expression of cIAP-1 which is consistent with previous reports showing that quercetin treatment did not affect the levels of cIAP-1 (Psahoulia et al., 2007; Kim et al., 2008). In previous studies, quercetin also leads to reduction in MMP ($\Delta\Psi_m$), and subsequent activation of caspases, thus ultimately inducing apoptosis via the intrinsic pathway in several types of cancer cells (Mouria et al., 2002).

Here, we also demonstrated that FEOJ-induced apoptosis is closely related to activation of p38 MAPK pathway. MAPK plays important roles in regulating apoptosis, and cell death (Robinson and Cobb, 1997), and quercetin plays a critical role in modulating cancer cell survival through p38 MAPK pathway (Chen et al., 2012). These findings support that p38 MAPK plays a critical role in FEOJ-induced apoptosis. In addition, the merit of FEOJ as compared to the quercetin is that the bioavailability of quercetin in FEOJ is much better than that of quercetin aglycone (Manach et al., 2005; Gibellini et al., 2011), because quercetin in FEOJ is in the form of quercetin glucosides, which are easily absorbed in the apical membrane of enterocyte (O'Leary et al., 2003).

In summary, this study demonstrated that FEOJ induced caspase-dependent apoptosis in U937 cells. The apoptosis was triggered through mitochondrial pathway by inducing loss of MMP ($\Delta\Psi_m$), and through modulating Bcl-2 and X-IAP proteins. In addition, FEOJ induces caspase-dependent apoptosis at least in part through the activation of p38 MAPK pathway (Figure 5). This study provides evidence that FEOJ might have anticancer property on human leukemic cells.

Acknowledgements

This work was supported by the grants from the National R & D Program for Cancer Control, Ministry of Health & Welfare, Republic of Korea (0820050).

References

Ashkenazi A (2002). Targeting death and decoy receptors of the tumour-necrosis factor superfamily. *Nat Rev Cancer*, **2**, 420-30.

Chen SF, Nieh S, Jao SW, et al (2012). Quercetin suppresses drug-resistant spheres via the p38 MAPK-Hsp27 apoptotic pathway in oral cancer cells. *PLoS One*, **7**, 49275.

Chun KH, Kosmeder JW, Sun S, et al (2003). Effects of deguelin on the phosphatidylinositol 3-kinase/Akt pathway and apoptosis in premalignant human bronchial epithelial cells. *J Natl Cancer Inst*, **95**, 291-302.

Gibellini L, Pinti M, Nasi M, et al (2011). Quercetin and cancer chemoprevention. *Evid Based Complement Alternat Med*,

Gupta RK, Banerjee A, Pathak S, et al (2013). Induction of mitochondrial-mediated apoptosis by *Morinda citrifolia* (Noni) in human cervical cancer cells. *Asian Pac J Cancer Prev*, **14**, 237-42.

Jang YJ, Park KS, Chung HY, et al (2003). Analysis of the phenotypes of Jurkat clones with different TRAIL-sensitivities. *Cancer Lett*, **194**, 107-17.

Jung KW, Won YJ, Kong HJ, et al (2014). Cancer statistics in Korea: incidence, mortality, survival, and prevalence in 2011. *Cancer Res Treat*, **46**, 109-23.

Kerr JF, Wyllie AH, Currie AR (1972). Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br J Cancer*, **26**, 239-57.

Kim YH, Lee DH, Jeong JH, et al (2008). Quercetin augments TRAIL-induced apoptotic death: involvement of the ERK signal transduction pathway. *Biochem Pharmacol*, **75**, 1946-58.

Kumi-Diaka J, Saddler-Shawnette S, Aller A, et al (2004). Potential mechanism of phytochemical-induced apoptosis in human prostate adenocarcinoma cells: Therapeutic synergy in genistein and beta-lapachone combination treatment. *Cancer Cell Int*, **4**, 5.

Lee J, Mitchell AE (2011). Quercetin and isorhamnetin glycosides in onion (*Allium cepa* L.): varietal comparison, physical distribution, coproduct evaluation, and long-term storage stability. *J Agric Food Chem*, **59**, 857-63.

Luo HR, Hattori H, Hossain MA, et al (2003). Akt as a mediator of cell death. *Proc Natl Acad Sci U S A*, **100**, 11712-7.

Manach C, Williamson G, Morand C, et al (2005). Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *Am J Clin Nutr*, **81**, 230-42.

Mouria M, Gukovskaya AS, Jung Y, et al (2002). Food-derived polyphenols inhibit pancreatic cancer growth through mitochondrial cytochrome C release and apoptosis. *Int J Cancer*, **98**, 761-9.

O'Leary KA, Day AJ, Needs PW, et al (2003). Metabolism of quercetin-7- and quercetin-3-glucuronides by an in vitro hepatic model: the role of human beta-glucuronidase, sulfotransferase, catechol-O-methyltransferase and multi-resistant protein 2 (MRP2) in flavonoid metabolism. *Biochem Pharmacol*, **65**, 479-91.

Palasap A, Limpaboon T, Boonsiri P, et al (2014). Cytotoxic effects of phytochemicals from *Caesalpinia mimosoides* Lamk on cervical carcinoma cell lines through an apoptotic pathway. *Asian Pac J Cancer Prev*, **15**, 449-54.

Psahoulia FH, Drosopoulos KG, Doubravska L, et al (2007). Quercetin enhances TRAIL-mediated apoptosis in colon cancer cells by inducing the accumulation of death receptors in lipid rafts. *Mol Cancer Ther*, **6**, 2591-9.

Robinson MJ, Cobb MH (1997). Mitogen-activated protein kinase pathways. *Curr Opin Cell Biol*, **9**, 180-6.

Walker PR, Sikorska M (1997). New aspects of the mechanism of DNA fragmentation in apoptosis. *Biochem Cell Biol*, **75**, 287-99.

Wiczowski W, Romaszko J, Bucinski A, et al (2008). Quercetin from shallots (*Allium cepa* L. var. *aggregatum*) is more bioavailable than its glucosides. *J Nutr*, **138**, 885-8.