

## RESEARCH ARTICLE

# Effects of Analgesine on Oxaliplatin-Induced Neurotoxicity in Patients with Gastrointestinal Cancer

Meng-Yan Liu, Xin-En Huang\*

### Abstract

**Background:** As the third generation of platinum-based antineoplastic agent against gastrointestinal cancer, oxaliplatin is considered to be associated with severe sensory neurotoxicity. According to previous studies, vitamin E, intravenous Ca/Mg and glutamine may partly reduce the incidence and severity of oxaliplatin-induced neurotoxicity. The aim of this study was to investigate the safety and efficacy of analgesine for preventing oxaliplatin-induced neurotoxicity in the patients with gastrointestinal tumors. **Method:** In this study, patients undergoing oxaliplatin-based chemotherapy were assigned to analgesine (experimental) group or control group. Analgesine 6ml was administered once a day for seven days from the day of oxaliplatin treatment. The National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE; version 3) was used to evaluate oxaliplatin-induced neurotoxicity. The incidence rates and grade of neurotoxicity of patients were assessed before and during (after four and eight cycles) treatment. **Results:** Totally, 82 patients were enrolled in this study, 42 in experimental group and 40 in control group. The occurrence of each grade neurotoxicity in the experimental group was significantly lower than that in control group. The overall occurrence rate was 31% vs 55% ( $P=0.043$ ) after 4 cycles and 52% vs 75% ( $P=0.050$ ) after 8 cycles. **Conclusion:** Analgesine appears could be effective in reducing oxaliplatin-induced neurotoxicity and be applied for patients with gastrointestinal tumors who would be treated with oxaliplatin-based chemotherapy.

**Keywords:** Analgesine - oxaliplatin-induced neurotoxicity

*Asian Pac J Cancer Prev*, 16 (10), 4465-4468

### Introduction

The incidence and mortality rates of gastrointestinal carcinoma, including gastric and colorectal cancers, are still high in China. Among them, colorectal cancer is the third most commonly cancer in males and the second in females, with over 1.2 million new cancer cases and 608,700 deaths estimated to occur in 2008 (Wu et al., 2014). Gastric cancer is also a leading cause of cancer-related deaths, with the highest incidence in Korea, Japan and China (Liu et al., 2014). Regarding that, oxaliplatin-based chemotherapy is a crucial regimen in both adjuvant and palliative settings for patients with gastrointestinal cancer, e.g., FOLFOX (Wang et al., 2007; Axel et al., 2010). Neurotoxicity is the main dose-limiting side effect of oxaliplatin-based chemotherapy, in case of grade III-IV neurotoxicity, dose reductions and interruption of treatment should be considered. It was reported that oxaliplatin-induced peripheral neurotoxicity is varying from 82% to 98%, and severe events occur in 10% to 20% of patients (Toru et al., 2013). The neurotoxicity of oxaliplatin is demonstrated in two distinct forms: acute and chronic. The former includes transient symptoms, e.g., cold related paresthesia, dysesthesia or jaw stiffness, and muscle cramps. The latter causes numbness and

tingling, affecting hands and feet, and vibration sensation (Axel et al., 2010; Cath et al., 2013). Various kinds of regimens have been tested to control the neurotoxicity of oxaliplatin, e.g., vitamin E, intravenous Ca/Mg and GM1 (Ganglioside-monosialic acid) (Lisa et al., 2011; Axel et al., 2010; Zhu et al., 2013). However, the treatment effect of these regimens were dissatisfied. Therefore, there is an urgent need to develop new medications to relieve oxaliplatin-induced neurotoxicity.

Analgesine (Extracts from Rabbit Skin Inflamed by Vaccinia Virus for Injection) is a biological agent with neurotropism. It has shown therapeutic effects for the control of neuropathic pain, especially in patients with diabetes mellitus, by playing an important role in improving peripheral circulation, anti-allergic reaction, and repairing cellular damage (Li et al., 2009). The possible mechanism of analgesine is considered as follows: first, directly affect hypothalamus to regulate the activities of neuroendocrine and cellular automaticity, consequently relieve the symptoms of numbness; second, improving peripheral circulation by regulating autonomic nerves, thus recovering peripheral circulation disorders (Li et al., 2009; Fu et al., 2010). The aim of this study was to assess the safety and efficacy of analgesine in controlling the oxaliplatin-induced neurotoxicity.

Department of Chemotherapy, the Affiliated Jiangsu Cancer Hospital of Nanjing Medical University & Jiangsu Institute of Cancer Research, Nanjing, China \*For correspondence: [huangxinen06@163.com](mailto:huangxinen06@163.com)

## Materials and Methods

### Eligibility criteria

Patients were eligible in this clinical trial if they had histologically or cytologically confirmed gastrointestinal tumors and were scheduled to receive oxaliplatin-based adjuvant or palliative chemotherapy in Jiangsu Cancer Hospital and Research Institute. All eligible patients were required to have a good performance status (ECOG 0-2), life expectancy 6months, routine blood test performed 0 to 3 days before chemotherapy and normal hematopoietic function as evidenced by white blood cell count 3000/ul and platelet count 100000/ul, normal hepatic function test (aspartate aminotransaminase and alanine aminotransferase less than 1.5 times of the upper limit of normal values), renal function test (serum total bilirubin<1.5mg/dl and creatinine<1.5mg/dl). Exclusion criteria included history of alcoholic intoxication, diabetes, central nervous system metastasis, preexisting neuropathy from any case, had received prior treatment with neurotoxic chemotherapy or radiation therapy, and patients who were pregnant or nursing.

### Neurotoxicity Evaluation

The incidence rates of neurotoxicity in this study were assessed at baseline and respectively after four and eight cycles of treatment, the grade of neurotoxicity was determined according to The National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE; version 3), which describe the four grades as follows: grade 1, loss of deep tendon reflexes or paresthesia, including tingling, but not interfering with function; grade

2, objective sensory alteration or paresthesia, including tingling, interfering with function, but not interfering with activities of daily living (ADL); grade 3, sensory alteration or paresthesia interfering with ADL; and grade4, permanent sensory losses that are disabling (Toru et al., 2013).

### Statistical analysis

All dates in this study were processed using the Stata.11 software. The rates of neurotoxicity were calculated for each grade and compared between experimental and control group. The difference in the grades of neurotoxicity between two groups was compared using Chi-square test and  $P<0.05$  defined as statistically significant.

## Results

In this study, 82 patients with gastrointestinal cancer enrolled from Jiangsu Cancer Hospital &Research Institute and divided into two groups, the experimental group (N=42), and the control group group (N=40).

As showed in Table 1, there were no significant difference between two groups in age, gender, performance status, primary tumor and tumor stage (Chi-square test). In our study, the incidence rates and grading scales of neurotoxicity in two groups were compared by Chi-square test. As showed in Table 2, after four cycles of treatment, the occurrence rates of grade 1, 2, 3, 4 neurotoxicity were 16.7%, 9.5% , 4.8% , 0, in experimental group respectively, while those in control group were 25%, 15%, 12.5%, 0.5%, respectively. The overall incidence of occurrence of neurotoxicity in experimental group (31%) was significantly lower than that in control group (55%), ( $p=0.043$ ). Moreover, after eight cycles of treatment, the incidence rates of grade1 neurotoxicity in experimental and control group was 28.6% versus 25%, that of grade 2 was 16.7% versus 20%, which of grade 3 was 9.5% versus 17.5%, and grade 4 was 2.3% versus 12.5%, respectively. Totally, 52% patients in experimental group experienced neurotoxicity while that in control group was 75%, (52%versus75%,  $P=0.050$ ). The proportion of patients who received analgesic experienced significant lower neurotoxicity than those who did not.

**Table 1. Patients Characteristics**

Characteristics	Experimental group N=42 (%)	Control group N=40 (%)	P Value
Age			
50	29 (72%)	28 (77%)	0.817
50	13 (28%)	12 (23%)	
Gender			
Male	14 (38%)	13 (27%)	1.021
Female	28 (62%)	27 (73%)	
Performance status			
0	23 (56%)	24 (67%)	0.727
1,2	19 (44%)	16 (33%)	
Primary tumor			
Gastric tumor	15 (31%)	12 (23%)	0.378
Colorectal tumor	27 (69%)	28 (77%)	
Tumor stage			
Locally advanced	14 (28%)	15 (40%)	0.550
Metastatic	28 (72%)	25 (60%)	

## Discussion

Oxaliplatin is one of the crucial components of chemotherapeutic regimens in the treatment of patients with gastrointestinal cancer. However, the majority

**Table 2. The Occurrence and Grade of Neurotoxicity Between Experimental Group and Control Group after Four, Eight Cycles**

Group	N	0 (%)	I (%)	II (%)	III (%)	IV (%)	occurrence rate	P value
After four weeks								0.043
Control group	40	18 (45)	10 (25)	5 (15)	5 (12.5)	2 (0.5)	55%	
Experimental group	42	29 (69)	7 (16.7)	4 (9.5)	2 (4.8)	0 (0)	31%	
After eight weeks								0.050
Control group	40	10 (25)	10 (25)	8 (20)	7 (17.5)	5 (12.5)	75%	
Experimental group	42	18 (42.9)	12 (28.6)	7 (16.7)	4 (9.5)	1 (2.3)	52%	

of patients receiving oxaliplatin-based chemotherapy developing different grades of neurotoxicity, which can seriously affect the quality of life, even leads to dose-reduction or interruption of chemotherapy regimens, consequently limiting treatment effect.

Analgesine is a kind of purified non-protein physiologically substances obtained by inflammation and immune reaction, which has a wide spectrum of pharmacologic actions (Liu et al., 2005). Analgesine exerts its neuroprotective function by affecting hypothalamus to regulate the activities of neuroendocrine as well as cellular automaticity, and through regulating autonomic nervous, consequently improving peripheral circulation. Clinical trials and animal studies have indicated that analgesine supplementation has remarkable efficacy in controlling chronic pain, nerve damage, numbness and other symptoms of nervous system (Fu et al., 2010; Li et al., 2009). Analgesine supplementation may relieve neurologic symptoms of diabetic peripheral neuropathy, which may also exert its curative effect in ameliorating the postherpetic neuralgia and relieving facial neuritis in combination with glucocorticoid (Liu et al., 2009; Li et al., 2011). Many other neuroprotective agents have been proved to be effective in reducing the neurotoxicity induced by oxaliplatin-based chemotherapy. Ca/Mg infusion has been supposed to have remit oxaliplatin associated neurotoxicity (Axel et al., 2010). However, there are still controversies in this field among oncologists. The CONCEPT trial, which assigned patients to receive Ca/Mg infusion prior or after oxaliplatin, seemed active against acute neurotoxicity at first. Unfortunately, this trial was prematurely aborted because of a low tumor response rate in the Ca/Mg arm. However, another French NEUROXA study demonstrated equivalent antitumor activity in patients receiving Ca/Mg and those receiving placebo (Game et al., 2005). In another prospective randomized, double-blind, placebo-controlled study, the antidepressant drug venlafaxine, has been reported to remarkably reduce the symptoms of oxaliplatin-induced neuropathy. Unfortunately, the occurrence of grade 1-2 vomiting was observed more frequently in patients who received venlafaxine (Toru et al., 2013; Dura et al., 2012). Compared with venlafaxine, analgesine has reassured security. Because analgesine is a kind of non-protein physiologically substances exerted from rabbit skin inflamed by vaccinia virus for injection, only working for the injured body and having no adverse effects on normal physiological function. According to the testified studies, the security of analgesine has no between-group differences compared with the placebos (Liu et al., 2009). What's more, analgesine is a kind of easily administered agents which is more convenient for the patients, with another additional advantage of the cost of analgesine is affordable for the majority of patients.

In our study, the overall incidence rate of neurotoxicity in experimental group was 31% after 4 cycles and 52% after 8 cycles, while that in control group was 55% and 75% respectively. The grade of neurotoxicity in experimental group was notably lower than that in control group after 4 cycles. There was significant difference in the occurrence rate of oxaliplatin-induced neurotoxicity

between two groups after 4 cycles ( $P=0.043$ ). However, there was no adequate statistical power detected after 8 cycles ( $P=0.050$ ). We speculated that this phenomenon was due to the modest sample size, single centre design and the chronic, cumulative oxaliplatin-induced neurotoxicity.

In summary, our data confirmed that analgesine could be effective in delaying the occurrence of oxaliplatin-induced neurotoxicity and reducing the grade of neurotoxicity. Larger prospective studies are needed to verify the clinical application of analgesine in treatment of oxaliplatin associated neurotoxicity.

## Acknowledgements

Dr. Xin-En Huang is supported by Traditional Chinese Medicine Scientific Research Project (LZ11091) and Jiangsu Province fourth stage "333 high-level Personnel Training Project" third levels of talent cultivating object.

## References

- Durand JP, Deplanque G, Montheil V, et al (2012). Efficacy of venlafaxine for the prevention and relief of oxaliplatin-induced acute neurotoxicity: results of eFFOX, a randomized, double-blind, placebo-controlled phase III trial. *Ann Oncol*, **23**, 200-5.
- Fu XR, Dong XH, Sun ZC, et al (2010). Clinical Observation to Combination Treatment of Lipoic Acid and Analgesine for Chemotherapy-induced Peripheral Neuropathy. *Chin General Pract*, 1007-9572.
- Gamelin L, Boisdrion-Celle M, Morel A, et al (2005). Oxaliplatin-related neurotoxicity: Interest of calcium/magnesium infusion and no impact on its efficacy. *J Clin Oncol*, **26**, 1188-9.
- Grothey A, Nikcevich DA, Sloan JA, et al (2010). Intravenous calcium and magnesium for oxaliplatin-induced sensory neurotoxicity in adjuvant colon cancer: NCCTG N04C7. *J Clin Oncol*, **29**, 421-7.
- Han CH, Khwaounjoo P, Kilfoyle DH, Hill A, McKeage MJ (2013). Phase I drug-interaction study of effects of calcium and magnesium infusions on oxaliplatin pharmacokinetics and acute neurotoxicity in colorectal cancer patients. *BMC Cancer*, **13**, 495
- Kono T, Hata T, Morita S, et al (2013). Goshajinkigan oxaliplatin neurotoxicity evaluation (GONE): a phase 2, multicenter, randomized, double blind, placebo controlled trial of goshajinkigan to prevent oxaliplatin induced neuropathy. *Cancer Chemo Pharmacol*, **72**, 1283-90.
- Li YH, Chen J, Xu X, et al (2009). Clinical observation of treatment diabetic peripheral neuropathy with analgesine. *Guide China Med*, 1671-8194, 2009, 13-0035-03.
- Lisa A. Kottschade, Jeff A. Sloan, Miroslaw A. Mazurczak, et al (2011). The use of vitamin E for the prevention of chemotherapy-induced peripheral neuropathy: results of a randomized phase III clinical trial. *Support Care Cancer*. 2011 November, **19**, 1769-77.
- Liu J, Huang XE, Feng JF, et al (2014). Further Study on Pemetrexed based chemotherapy in Treating Patients with Advanced Gastric Cancer (AGC). *Asian Pac J Cancer Prev*, **15**, 6587-90.
- Liu ZB, et al (2005). Extract from Rabbit Skin Inflamed by Vaccinia Virus for Injection- Analgesine. *Qilu Pharmacol Affairs*, 1672-7738 (2005) 08- 0509- 02.
- Wang WS, Lin JK, Chen PM, et al (2007). Oral Glutamine Is Effective for Preventing Oxaliplatin-Induced Neuropathy

- in Colorectal Cancer Patients. *The Oncologist*, **12**, 312-9.
- Wu XY, Huang XE, et al (2014). A Predictive Model for Evaluating Responsiveness to Pemetrexed Treatment in Patients with Advanced Colorectal Cancer. *Asian Pac J Cancer Prev*, **15**, 5941-4.
- Wu XY, Huang XE, You SX, et al (2013). Phase II study of pemetrexed as second or third line combined chemotherapy in patients with colorectal cancer. *Asian Pac J Cancer Prev*, **14**, 2019-22.
- Xiao Y, Liu J, Liu YC, Huang XE, et al (2014). Phase II Study on EANI Combined with Hydrochloride Palonosetron for Prevention of Chemotherapy-induced Nausea and Vomiting Following Highly Emetogenic Chemotherapy. *Asian Pac J Cancer Prev*, **15**, 3951-4.
- Yu H, Ji J, Ma CH, et al (2013). Efficacy of glutathione in prevention of neurotoxicity induced by oxaliplatin. *Modern Oncology*, **21**, 1342-3.
- Zhu YY, Yang JL, Jiao SC, et al (2007). Ganglioside-monosialic acid (GM1) prevents oxaliplatin-induced peripheral neurotoxicity in patients with gastrointestinal tumors. *World J Surg Oncol*, **2013**, 11-9.