

RESEARCH ARTICLE

Diet and Colorectal Cancer Risk in Asia - a Systematic Review

Salman Azeem, Syed Wasif Gillani, Ammar Siddiqui, Suresh Babu Jandrajupalli, Vinci Poh, Syed Azhar Syed Sulaiman

Abstract

Diet is one of the major factors that can exert a majorly influence on colorectal cancer risk. This systematic review aimed to find correlations between various diet types, food or nutrients and colorectal cancer risk among Asian populations. Search limitations include Asian populations residing in Asia, being published from the year 2008 till present, and written in the English language. A total of 16 articles were included in this systematic review. We found that red meats, processed meats, preserved foods, saturated/animal fats, cholesterol, high sugar foods, spicy foods, tubers or refined carbohydrates have been found by most studies to have a positive association with colorectal cancer risk. Inversely, calcium/dairy foods, vitamin D, general vegetable/fruit/fiber consumption, cruciferous vegetables, soy bean/soy products, selenium, vitamins C,E and B12, lycophene, alpha-carotene, beta-carotene, folic acid and many other vitamins and minerals play a protective role against colorectal cancer risk. Associations of fish and seafood consumption with colorectal cancer risk are still inconclusive due to many varying findings, and require further more detailed studies to pinpoint the actual correlation. There is either a positive or no association for total meat consumption or white meats, however their influence is not as strong as with red and processed meats.

Keywords: Colorectal cancer - diet - food - Asia - risk - correlation

Asian Pac J Cancer Prev, 16 (13), 5389-5396

Introduction

Cancer is one of the major causes of morbidity and death globally, and colorectal cancer is among the top 5 cancers most commonly diagnosed in both men and women alike (WHO, 2015; Talaiezhadah et al., 2013). In 2008 alone, there has been more than 1.2 million newly diagnosed cases and 608,700 deaths caused by colorectal cancer. There is a higher incidence of colorectal cancer in economically developed countries compared to economically developing countries, and comparatively, more males develop colorectal cancer compared to the fairer sex (Barouni et al., 2012; Morrison et al., 2013; WHO, 2015).

While there are many factors that can alter the risk of getting colorectal cancer, dietary factors are known to be closely correlated with any type of cancer, and its link with colorectal cancer in particular is extremely profound (Jemal et al., 2011, Xue-Yan and Xin-Em Huang, 2015; Wu, Huang, 2015; WHO, 2015). Numerous studies over the years have studied and analyzed different dietary aspects and their link to colorectal cancer to try and find the correlation and reasoning behinds these links, and generally have found that intake of red meats and processed meats, consuming meats cooked at high temperatures, high dietary animal fats and high dietary

sugars potentially increase risk, while high fruit and vegetable consumption, fish consumption and high intake of certain nutrients such as selenium, calcium, vitamin D and folate may have a protective effect against colorectal cancer (Jemal et al., 2011; Magaji et al., 2014; ACS, 2015; CR-UK, 2015; WHO, 2015). However, a majority of these studies were done in countries where colorectal cancer was historically prevalent, mostly Western countries. In Asia, where colorectal cancer has comparatively recently become a more prominent problem, less studies have been conducted on the impact of diet on colorectal cancer among Asians. While it is true that many Asian countries are experiencing increasing 'Westernization' in their dietary patterns (Yusoff, 2012; Suh et al., 2013; Tashan, Andsoy, Iyugun, 2013; Lai et al., 2014; WHO, 2015), the amount, patterns, variations and cooking methods of their diet still differ substantially, and there may be notable difference in bodily response of the Asian population towards different diet components compared to the Western population (CDC, 2014; WHO, 2015; Li Li et al., 2015; Ersin Ozaslan et al., 2015).

Of the studies available that were done on the link of diet and colorectal cancer among the Asian population, there are varied conclusions to the effect of these dietary components towards colorectal cancer risk. Hence, this systematic review aims to compile the available data to

Table 1. Study Findings on Various Diets/Foods/Nutrients

Diet/Food/Nutrient	Conclusion	Articles	Remarks
Fish/Seafood	Positive association	Lee et al (2009)	-positive association for cholesterol rich fish (eel, shrimp, shellfish)
		Ganesh et al (2009)	-positive association for 'dried fish' among men, due to salting, drying and preserving
	Inverse association	Ganesh et al (2009)	-inverse association for fresh fish
		Mahfouz et al (2014)	-inverse association for seafood
	No association	Pham et al (2013)	-no association in cohort studies
			-weak inverse association in case-control studies
		Sugawara et al (2009)	-may be due to comparatively high daily fish consumption
		Lee et al (2009)	-no association for total fish intake
			-lack of inverse association may be caused by raising seafood in polluted waters
	Nayak et al (2009)	-inverse but insignificant association for daily fish consumption	
Total meat	Positive association	Takachi et al (2011)	-positively associated in men (distal colon cancer)
		Ganesh et al (2009)	-positive association in women
	Inverse association	-	
	No association	Lee et al (2009)	-may be due to overall low meat consumption
		Butler et al (2008)	-no association of meat-based diet pattern
-results may be affected by cooking methods			
Red meat	Positive association	Mahfouz et al (2014)	
		Pham et al (2014)	
		Takachi et al (2011)	-positively associated among women (proximal colon cancer)
		Ramadas et al (2009)	-associated with high temperature cooking method
			-increases colorectal adenoma risk but not colorectal carcinoma risk
		Nayak et al (2009)	-beef
			-positive association regardless of consumption quantity
		Nashar et al (2008)	
	Arafa et al (2011)		
	Inverse association	-	
No association	Takachi et al (2011)	-No association for rectal cancer	
Processed meats	Positive association	Mahfouz et al (2014)	
		Pham et al (2014)	
		Nashar et al (2008)	
	Inverse association	-	
No association	Takachi et al (2011)		
White meat	Positive association	-	
	Inverse association	-	
	No association	Lee et al (2009)	
Ramadas et al (2009)		-slight but insignificant positive association	

attempt to reach a consensus on their influence towards colorectal cancer, and for the studies that report otherwise, to attempt to explain what lead to their findings.

Materials and Methods

Literature search was done on the influence of various

Cooking practices	Positive association	Lee et al (2009)	-‘smoking’ method of Chinese cooking
		Takachi et al (2011)	-cooking muscle meats at high temperatures
	Inverse association	-	
Sodium/Preserved foods	Positive association	Mahfouz et al (2014)	
		Takachi et al (2009)	-positive association of preserved foods with high salt content
		Ganesh et al (2009)	-positive association for salted fish roe and dried fish, not due to amount of salt per se but the presence of chemical carcinogens
	Inverse association		
	No association	Takachi et al (2009)	-no association of overall high sodium consumption
Saturated/Animal fats, cholesterol	Positive association	Mahfouz et al (2014)	-positive association with fast foods
		Lee et al (2009)	-positive association for high cholesterol intake
			-positive association for high egg intake (cholesterol rich food)
		Ramadas et al (2010)	-positive association of total cholesterol and LDL with colorectal adenoma risk
	Nashar et al (2008)	-positive association for high fat meats, whole fat dairy products and fried eggs with colorectal adenoma risk	
	Inverse association	-	
No association	Lee et al (2009)	-no association for fat consumption	
High sugar foods Spices	Positive association	Mahfouz et al (2014)	-positive association with artificial sweeteners and soft drinks
		Nayak et al (2009)	-positive association with heavy sugar consumption
	Inverse association	-	
	No association	-	
	Positive association	Mahfouz et al (2014)	-positive association with spicy foods
		Nayak et al (2009)	-positive association with pungent spices
	Inverse association	-	
No association	-		

dietary components, food types or nutrients on colorectal risk. No study design limitations were implemented. Search limitations were confined to studies done on Asian populations residing in Asia, being published from the year 2008 till present, and written in the English language. However, systematic reviews and meta-analyses done in Asia but did not limit their article search to Asian populations residing in Asia were excluded. Search keywords used were ‘*diet*’, ‘*nutrition*’, ‘*food*’, ‘*meat*’, ‘*fish*’, ‘*vegetables*’, ‘*fruit*’, ‘*colorectal cancer*’, ‘*risk*’, ‘*impact*’ and ‘*Asia (or individual Asian countries)*’; words with similar meaning were also attempted. Search terms were linked using the Boolean operator ‘and’. Primary search was done using databases and individual journal websites, like Pubmed, ResearchGate, Google Scholar, Wiley Online Library, British Journal of Cancer, Cancer Epidemiology, The American Journal of Clinical Nutrition, Asian Pacific Journal of Cancer Prevention and

so forth. After primary search, a secondary search was done on the reference lists of the articles found.

Primary screening focused mainly on a quick scan of the title, abstract and discussion sections. Articles that passed primary screening are then read fully in detail as secondary screening.

Results

A total of 16 articles matched the aforementioned criteria and were included for review. 7 articles were from Japan, 2 each from Malaysia and India, and 1 each from Egypt, China, Singapore, Jordan and Saudi Arabia. As seen, a majority of the studies were conducted in Japan, which is not surprising as incidences in Japan has been increasing drastically and have exceeded peak rates of countries that were previously among those with the highest incidence rates, such as the United States of

Calcium/Dairies	Positive association	Mizoue et al (2008)	-positive association for dairy products (excluding milk)
		Nashar et al (2008)	-positive association for whole fat dairy products
	Inverse association	Mahfouz et al (2014)	-inverse association with calcium rich diet
		Lee et al (2009)	-inverse association for milk
		Ishihara et al (2008)	-inverse association of high dietary calcium in men
	Mizoue et al (2008)	-inverse association only for those with high vitamin D intake/exposure	
No association			
Vitamin D	Positive association		
	Inverse association	Mizoue et al (2008)	-Inverse association of high dietary Vitamin D for those with low sunlight exposure
	No association	Ishihara et al (2008)	-no significant association of dietary Vitamin D in both genders
-however may potentiate calcium's effects on risk			
Vegetables/Fruits /Fibers (general)	Positive association		
	Inverse association	Mahfouz et al (2014)	-inverse association for high fruit and vegetable consumption, high fiber bread, fruit juices
		Ramadas et al (2009)	-inverse association for high vegetable and fruit intake
			-protective role of high-fruit low-meat diet
			-vegetables may have stronger preventive role for risk of adenoma to carcinoma conversion, compared to initial adenoma formation
		Ramadas et al (2010)	-inverse association for crude fiber intake
		Arafa et al (2011)	-inverse association for fruits, vegetables and fiber
	Nayak et al (2009)	-inverse association for fruits and vegetables	
	Nashar et al (2008)	-inverse association for fiber, vegetables, fruits, whole wheat products	
No association	Butler et al (2008)	-no association for primarily vegetable diet	
	Ganesh et al (2009)	-no association for non-cruciferous vegetables	
Cruciferous vegetables	Positive association	-	
	Inverse association	Mahfouz et al (2014)	
		Ganesh et al (2009)	-cabbage, sprouts
No association	-		
Tubers	Positive association	Ramadas et al (2009)	
		Nayak et al (2009)	-tapioca
	Inverse association	-	
No association	-		
Refined carbohydrates	Positive association	Nayak et al (2009)	
	Inverse association	-	
	No association	-	

America, Australia and New Zealand (Center et al., 2009; Cetin et al., 2012; Bozkurt et al., 2014).

Some articles studied on a single type of diet whereas some tackled multiple diet areas. The table below summarizes the findings of the articles on each diet type. A positive association means an increase in colorectal cancer risk, whereas an inverse association means a decrease in

colorectal cancer.

In summary, most studies found that fish and seafood consumption has no association. There is either a positive or no association for total meat consumption or white meats. There is a positive association for red meats, processed meats, preserved foods, saturated/ animal fats, cholesterol, high sugar foods, spicy foods,

Soy bean/products	Positive association	-	
	Inverse association	Ramadas et al (2009)	
	No association	-	
Specific nutrients	Positive association	-	
	Inverse association	Ramadas et al (2010)	-inverse association of alpha carotene, beta-carotene, vitamin C, vitamin D, vitamin E and lycophene with colorectal adenoma risk
		Arafa et al (2011)	-antioxidants (selenium and vitamin C), dietary folic acid and vitamin B12
		Nashar et al (2008)	-selenium and folic acid
No association	-		

tubers or refined carbohydrates with colorectal cancer risk. Calcium/dairies, vitamin D (by aiding calcium absorption), general consumption of vegetables/fruits/fiber, cruciferous vegetables, soy bean/products, selenium, vitamin C, vitamin E, vitamin B12, lycophene, alpha carotene, beta-carotene or folic acid have a protective role against colorectal cancer risk. Influence of cooking methods depend individually.

Discussion

Fish and seafoods are well-known for having high n-3 fatty acid content, and it has been long believed that n-3 fatty acids, more popularly known as omega-3 fatty acids, are capable of preventing carcinogenesis via multiple pathways (Pham et al., 2013). According to Cockbain et al, omega-3 fatty acids are capable of reducing mucosal epithelial cell proliferation and have anti-inflammatory properties (Cockbain et al., 2012). However, more and more studies show conflicting findings on the efficacy of omega-3 fatty acids towards cancer prevention, and a re-analysis of past research suggested that there may not be any reduction of cancer risk after all (ACS, 2013; Dirican et al., 2014). Then again, interpretation of these studies have been difficult due to numerous differences in study design and lack of control of variables. Type of fish, whether the fish consumed is fresh or processed, method of preparation and differences of unit of reporting outcome measures have made it challenging to come up with accurate results (cockbain et al., 2012). A meta-analysis that attempted to navigate these variables managed to find that consumption of fish does inhibit colorectal carcinogenesis (Geelen et al., 2007; He et al., 2014). Other than omega-3 fatty acids, fish are also known to contain vitamin D and selenium, which are also known to have anti-cancer properties (Pham et al., 2013). The World Cancer Research Fund and American Institute for Cancer Research (WCRF/AICR) also voiced agreement in the role of fish in colorectal cancer prevention, though they did clarify that their findings are based on limited evidence (CR-UK, 2015).

Of the 6 articles included in this review that discussed the link between fish or seafood with colorectal cancer, 4 have concluded that there is no association. The systematic review by Pham et al. (2013) found no association in the cohort studies but a weak inverse association in case-control studies, but also noted that the case-control studies

are prone to recall and selection bias due to a tendency of selecting controls among those who have undergone a health screening, and did not adjust for meat consumption and other potential confounding factors. This being said, the 2 studies that reported that fish has a protective effect against colorectal cancer were both case-control studies, also having picked their controls from an oncology center and hospital respectively. Of the 2 articles that reported an increase in colorectal risk by fish consumption, one was reporting on 'dry fish', and hence the increase in risk may be attributable to other factors such as the fish being a preserved food item, which is found to increase colorectal cancer risk (Ganesh et al., 2009; Mahfouz et al., 2014). The other paper by Lee et al. (2009) found positive association in cholesterol rich fish, and cholesterol is also found to increase the risk of colorectal cancer. The other 3 papers that reported no association each attempted to explain possible factors that lead to a difference in results of their findings compared to those that reported otherwise, with Sugawara et al stating that it may be due to the comparatively high fish consumption of their study cohort and the different species of fish consumed (Sugawara et al., 2009); Lee et al. (2009) explained on the possible effects of consuming seafood raised in polluted waters in Shanghai, where the study was conducted; Nayak SP et al. (2009) reported on the formation of heterocyclic amines (HCA) on fish meat when cooked which can be carcinogenic, though also stating that fish consumption is safe but does not significantly reduce colorectal cancer risk. Hence, it may be said that instead of having no association at all with colorectal cancer risk, it may be said that results are inconclusive, and further more detailed studies are required to pinpoint the actual association.

Regarding the effect of meats towards colorectal cancer risk, less studies are done on total meat consumption and white meats as they are not identified to be major risks of colorectal cancer, and this is mirrored in our findings. Most studies focus more on red meats and processed meats, which are well known for a long time to play an important role in increasing colorectal cancer risk. Numerous large scale studies conducted in countries with high red meat and processed meat consumption have confirmed this link (Harvard Medical School, 2008). The same has been found in this systematic review, with 7 out of 9 studies supporting the positive association of red meats and colorectal cancer risk and 3 out of 4 studies supporting the link regarding processed meats.

Red meat has high amounts of heme iron, which can not only damage DNA, but also catalyze genotoxic and cytotoxic aldehyde formation and increase multisite carcinogens known as N-nitroso compounds (NOCs). NOCs are also formed in the making of processed meats. Meats are also commonly cooked at high temperature, and this leads to the formation of HCAs and polycyclic aromatic hydrocarbons (PAHs) that are carcinogens as well (Takachi et al., 2011; Pham et al., 2014). In the study done by Lee et al. (2009) which studies on the effects of Chinese cooking methods towards colorectal cancer risk, only the 'smoking' method, which exposes the meat to heat for a long period of time, showed a link. NOCs, PAHs and HCAs are capable to interact with our genes and alter the risk of forming colorectal adenomas. Nayak et al. (2009) explained that beef, a commonly consumed red meat, can increase colorectal cancer risk irrespective of HCA content, meaning that HCA is not concentration dependent and just the involvement of beef in one's diet can increase risk of getting colorectal cancer. They also stated that meat consumption can increase bile acid production which can promote mitosis. Interestingly, Ramadas et al. (2009) reported that red meat can increase the risk of developing colorectal adenomas but does not affect carcinoma risk. Nevertheless, this is still of concern as carcinomas can easily develop from existing adenomas.

4 out of 5 studies confirmed a positive association of saturated or animal fats and cholesterol with colorectal cancer risk, though none gave detailed explanations on the mechanism. However Lee et al. (2009) did mention that cholesterol plays a role as co-carcinogen in cancer development, though a brief search revealed that most studies reporting this finding are animal studies. A number of other mechanisms have also been proposed, including the link between a high fat diet, bile acid and cancer. High fat diets lead to higher bile acid production, and this leads to a higher generation of reactive nitrogen species (RNS) and reactive oxygen species (ROS). Exposure of colonic cells to RNS and ROS can lead to DNA damage and cell mutation. Additionally, long term exposure to RNS and ROS can reduce cell apoptosis capabilities, increasing mutagenesis risk (Bernstein et al., 2009; Ozdemir et al., 2014).

Takachi et al. (2009) studied on the effects of sodium and preserved foods on colorectal cancer risk. They found that while overall high sodium consumption does not have an association, preserved foods with high salt content does have a positive link. Being a Japanese study they singled out salted fish roe and dried fish as examples. They explained that it was not primarily due to the salt content that caused the link, but due to the curing process by the salt that lead to carcinogen formation similar to that mentioned above regarding the cooking and curing of meats. Ganesh et al. (2009) also reported the increase in colorectal cancer risk by 'dry fish' consumption, due to the involvement of salting and drying of the fish for preservation purposes.

Both studies that studied on the effects of sugar consumption to colorectal cancer risk found a positive association, with Mahfouz et al specifically mentioning artificial sweeteners and soft drinks, while Nayak et al.

(2009) mentioned overall heavy sugar consumption. Nayak et al. (2009) also stated a positive association of refined carbohydrates with the risk (Mahfouz et al., 2014). High carbohydrate and sugar content can lead to excess insulin production, and not only so, such a diet can lead to accumulation of visceral adipose tissue and hence obesity, and this in turn is also associated with high insulin production. Insulin, being a growth factor in the colon, can enhance colonic cell proliferation and increase the risk of development of colorectal cancer (Schioen et al., 1999; Levy, 2012).

Spices are commonly included in various Asian cuisine, so not surprisingly there are a couple of studies that try to find the link between spicy food and colorectal cancer. Both studies found a positive link, with Nayak et al. (2009) specifying on pungent spices, such as chilies. There seems to be conflicting findings on the effects of capsaicin, the alkaloid that gives chilies their pungency, on cancer; some reported it to be anti-carcinogenic, while others reported it to be otherwise. There are no human studies so far. But findings in their study alone found that long term consumption of pungent spices can lead to colorectal cancer. Mahfouz et al. (2014) also mentioned similar findings and quoted Nayak et al. (2009) in their study.

Calcium has been frequently mentioned as a protective nutrient against colorectal cancer. Calcium is capable of acting at multiple tiers of colonic cell organization by manipulating a complex series of signaling events, preventing colorectal carcinogenesis. Proliferation of colonic epithelial cells is reduced, and recurrence of adenomas is also likewise reduced (Cho et al., 2004). Calcium can also enhance colonic cell apoptosis (Miller, 2005). Vitamin D also plays a role in colorectal cancer prevention as not only does it aid in the absorption of calcium, it itself also exerts anti-cancer properties by regulating apoptosis, proliferation and differentiation of cells and inhibits angiogenesis (Mizoue et al., 2008). 4 studies in our review support the inverse association of calcium and 1 study supports the role of vitamin D. Some studies also mentioned the role of other nutrients in dairy products that also have anti-carcinogenic properties, such as conjugated linoleic acid (Mizoue et al., 2008). However, Mizoue et al. (2008) has noted that only milk exerts this protective effect, and that other dairy products actually can increase colorectal cancer risk. Different dairy products contain different compositions and are produced differently; hence more studies need to be conducted to explain the association. Findings by Nahsar et al. (2008) on the positive correlation between whole fat dairy products and colorectal cancer risk may be attributed to the fat contents of the product, which mentioned previously is positively linked to colorectal cancer risk. As for Ishihara et al that does not find an association of dietary vitamin D with colorectal cancer risk, it may be due to the relatively low calcium intake among the study population, therefore without sufficient calcium vitamin D cannot exert its influence on calcium fully, and hence the association may not be so pronounced (Ishihara et al., 2008).

High vegetable and fruit intake has always been viewed as beneficial to health and an important element in cancer

prevention, due to rich fiber content, vitamins, minerals and antioxidants (Nayak et al., 2009). This is mirrored in our findings where 6 out of 8 studies supported the inverse association towards colorectal cancer risk, and none reported positive associations. Nashar et al. (2008) stated several potential protective mechanisms by vegetables and fruits, including shortening of feces transit time by fiber, which reduces contact time with the colonic wall and hence less exposure to potential carcinogens; production of short chain fatty acids when fiber is fermented in the colon, which has apoptotic, antiproliferative and differentiating properties (Hinnebusch et al., 2002); the action of selenium, which is present in cereals, which acts as a cofactor for glutathione peroxidase that prevents oxidative tissue damage and hence suppresses cancer cell proliferation; folic acid from fruits, which not only helps in DNA nucleotide synthesis, which is important for DNA repair and replication, but also aids in production of S-adenosylmethionine (SAM), which is important for DNA methylation and can act in the suppression of viral repetitive DNA (Ulrich, 2007); and the provision of anticarcinogens such as protease inhibitors, resveratrol, carotenoids, flavonoids, vitamin C, isothiocyanates and organosulfides. Ramadas et al. (2010) has noted a stronger preventive role by vegetable consumption for the prevention of adenoma to carcinoma conversion, compared to the initial formation of adenomas. A family of vegetables, known as cruciferous vegetables, have been of particular interest as they are not only rich in nutrients such as folate, minerals, various vitamins and carotenoids, they also contain glucosinolates that are broken down during food preparation and consumption to biologically active nitriles, indoles, isothiocyanates and thiocyanates. Isothiocyanates and indoles exhibit anticancer properties, such as inducing apoptosis, anti-inflammatory effects, inactivate carcinogens and protect from DNA damage, among others (NCI, 2012).

Tubers, despite also being rich in fibers, have been found to have a positive correlation with colorectal cancer risk, as found by Ramadas et al. (2009) and Nayak et al. (2009). Nayak et al. (2009) also mentioned tapioca in particular as this is a staple food among their study population in Kerala, South India. They attributed the increase in colorectal cancer link to the presence of toxins such as cyanide derivatives and linamarin. These toxins require processing to be removed, but tapioca was regularly eaten unprocessed and simply boiled. The exact mechanism of risk increment is unknown, but in the study it has caused a 4-fold increase in risk, and it is suspected that it could be due to the direct contact of toxins to the colonic mucosa.

Consumption of soy beans or soy products, which are highly consumed in certain parts of Asia, have been found by Ramadas et al to be inversely associated to colorectal cancer risk. This may be attributable to soy isoflavones, and genistein in particular promotes apoptosis and inhibits proliferation of cancerous colonic cells. Also known as a phytoestrogen, it has the ability to target androgen and estrogen mediated signaling pathways in carcinogenesis. Its antioxidant properties allows it to inhibit metastasis and angiogenesis (Sarkar and Li, 2003). However, Ramadas et

al. (2009) also noted that there may be an underestimation of total soy food intake due to recall bias.

Red meats, processed meats, preserved foods, saturated/animal fats, cholesterol, high sugar foods, spicy foods, tubers or refined carbohydrates have been found by most studies to have a positive association with colorectal cancer risk. Inversely, calcium/dairy foods, vitamin D, general vegetable/fruit/fiber consumption, cruciferous vegetables, soy bean/soy products, selenium, vitamins C, E and B12, lycopene, alpha-carotene, beta-carotene, folic acid and many other vitamins and minerals play a protective role against colorectal cancer risk. Association of fish and seafood consumption with colorectal cancer risk is still inconclusive due to many varying findings, and requires further more detailed studies to pinpoint the actual correlation. There is either a positive or no association for total meat consumption or white meats, however their influence is not as strong as red and processed meats. Several diet/food types require more or further investigation on their influence, particularly foods that are frequently consumed in the Asian population. Studies that focus more on the influence of such foods towards the Asian genetics and physique in particular are scarce, and more studies in this area may unearth new findings that can be helpful towards the prevention of colorectal cancer in Asia.

References

- American Cancer Society (ACR) (2015). What are the risk factors for colorectal cancer?
- American Cancer Society (ACR) (2013). Omega-3 Fatty Acids.
- Arafa MA, Waly MI, Jriesat S, et al (2011). Dietary and lifestyle characteristics of colorectal cancer in Jordan: a case-control study. *Asian Pac J Cancer Prev*, **12**, 1931-6.
- Barouni M, Larizadeh MH, Sabermahani A, et al (2012). Markov's modeling for screening strategies for colorectal cancer. *Asian Pac J Cancer Prev*, **13**, 5125-9.
- Bozkurt O, Inanc M, Turkmen E, et al (2014). Clinicopathological characteristics and prognosis of patients according to recurrence time after curative resection for colorectal cancer. *Asian Pac J Cancer Prev*, **15**, 9277-81.
- Butler LM, Wang R, Koh WP, et al (2008). Prospective study of dietary patterns and colorectal cancer among Singapore Chinese. *Br J Cancer*, **99**, 1511-6.
- Bernstein H, Bernstein C, Payne CM, et al (2009). Bile acids as endogenous etiologic agents in gastrointestinal cancer. *World J Gastroenterol*, **15**, 3329-40.
- Cockbain AJ, Toogood GJ, Hull MA (2012). Omega-3 polyunsaturated fatty acids for the treatment and prevention of colorectal cancer. *Gut*, **61**, 135-49.
- Cancer Research UK (2015). Bowel cancer risk factors.
- Centers For Disease Control And Prevention (CDC) (2014). What are the risk factors for colorectal cancer?
- Center MM, Jemal A, Smith RA, et al (2009). Worldwide Variations in Colorectal Cancer. *CA Cancer J Clin*, **59**, 366-78.
- Cetin B, Kaplan MA, Berk V, et al (2012). Prognostic factors for overall survival in patients with metastatic colorectal carcinoma treated with vascular endothelial growth factor-targeting agents. *Asian Pac J Cancer Prev*, **13**, 1059-63.
- Cho E, Smith-Warner SA, Spiegelman D, et al (2004). Dairy Foods, Calcium, and Colorectal Cancer: A Pooled Analysis of 10 Cohort Studies. *J National Cancer Inst*, **96**, 1015-22.

- Dirican A, Varol U, Kucukzeybek Y, et al (2014). Treatment of metastatic colorectal cancer with or without bevacizumab: can the neutrophil/lymphocyte ratio predict the efficiency of bevacizumab? *Asian Pac J Cancer Prev*, **15**, 4781-4786.
- Ersin Ozaslan, Ayse Ocak Duran, Oktay Bozkurt, et al (2015). Analyses of multiple factors for determination of "selected patients" who should receive rechallenged treatment in metastatic colorectal cancer: a retrospective study from Turkey. *Asian Pac J Cancer Prev*, **16**, 2833-8.
- Ganesh B, Talole SD, Dikshit R (2009). A case-control study on diet and colorectal cancer from Mumbai, India. *Cancer Epidemiol*, **33**, 189-93.
- Geelen A, Schouten JM, Kamphuis C, et al (2007). Fish consumption, n-3 fatty acids, and colorectal cancer: a meta-analysis of prospective cohort studies. *Am J Epidemiol*, **166**, 1116-25.
- Harvard Medical School (2008). Red meat and colon cancer: Harvard Health Publications; 2005.
- Hinnebusch BF, Meng S, Wu JT, et al (2002). The effects of short-chain fatty acids on human colon cancer cell phenotype are associated with histone hyperacetylation. *J Nutr*, **132**, 1012-7.
- He WZ, Jiang C, Yin CX, et al (2014). Prognostic model built on blood-based biomarkers in patients with metastatic colorectal cancer. *Asian Pac J Cancer Prev*, **15**, 7327-31.
- Ishihara J, Inoue M, Iwasaki M, et al (2008). Dietary calcium, vitamin D, and the risk of colorectal cancer. *Am J Clin Nutr*, **88**, 1576-83.
- Jemal A, Bray F, Center MM, et al (2011). Global cancer statistics. *A Cancer J Clin*, **61**, 69-90.
- Lai JH, Zhou YJ, Bin D, Qiangchen, Wang SY (2014). Clinical significance of detecting lymphatic and blood vessel invasion in stage II colon cancer using markers D2-40 and CD34 in combination. *Asia Pac J Cancer Prev*, **15**, 1363-70.
- Li li, Shi Chen, Ke Wang, et al (2015). Diagnostic value of endorectal ultrasound in preoperative assessment of lymph node involvement in colorectal cancer: a meta-analysis. *Asian Pac J Cancer Prev*, **16**, 3485-91.
- Levy R (2012). Insulin and colon cancer linked. Harvard Gazette 2012 (col. Science and Health)
- Lee S-A, Shu XO, Yang G, et al (2009). Animal Origin Foods and Colorectal Cancer Risk: A Report From the Shanghai Women's Health Study. *Nutrit Cancer*, **61**, 194-205.
- Lee JE, McLerran DF, Rolland B, et al (2013). Meat intake and cause-specific mortality: a pooled analysis of Asian prospective cohort studies. *Am J Clin Nutr*, **98**, 1032-41.
- Mahfouz EM, Sadek RR, Abdel-Latif WM, et al (2014). The role of dietary and lifestyle factors in the development of colorectal cancer: case control study in Minia, Egypt. *Cent Eur J Public Health*, **22**, 215-22.
- Magaji BA, Moy FM, Roslani AC, et al (2014). Descriptive epidemiology of colorectal cancer in University Malaya Medical Centre, 2001 to 2010. *Asian Pac J Cancer Prev*, **15**, 6059-64.
- Morrison DS, Parr CL, Lam TH, et al (2013). Behavioural and metabolic risk factors for mortality from colon and rectum cancer: analysis of data from the Asia-Pacific cohort studies collaboration. *Asian Pac J Cancer Prev*, **14**, 1083-7.
- Mizoue T, Kimura Y, Toyomura K, et al., (2008) Calcium, dairy foods, vitamin d, and colorectal cancer risk: the fukuoka colorectal cancer study. *Cancer Epidemiol Biomark Prev*, **17**, 2800-7.
- Miller EA, Keku TO, Satia JA, et al., (2005). Calcium, Vitamin D, and apoptosis in the rectal epithelium. *Cancer Epidemiol Biomark Prev*, **14**, 525-8.
- National Cancer Institute (NCI) (2012). Cruciferous vegetables and cancer prevention.
- Nayak SP, Sasi MP, Sreejayan MP, et al., (2009). A case-control study of roles of diet in colorectal carcinoma in a South Indian population. *Asian Pac J Cancer Prev*, **10**, 565-8.
- Nashar RM, Almurshed KS (2008). Colorectal cancer: a case control study of dietary factors, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia. *J Family Community Med*, **15**, 57-64.
- Ozdemir Y, Akin ML, Sucullu I, Balta AZ, Yucel E (2014). Pretreatment neutrophil/lymphocyte ratio as a prognostic aid in colorectal cancer. *Asia Pac J Cancer Prev*, **15**, 2647-50.
- Pham NM, Mizoue T, Tanaka K, et al., (2013). Fish consumption and colorectal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. *Jpn J Clin Oncol*, **43**, 935-41.
- Pham NM, Mizoue T, Tanaka K, et al (2014). Meat consumption and colorectal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. *Jpn J Clin Oncol*, **44**, 641-50.
- Ramadas A, Kandiah M (2009). Food intake and colorectal adenomas: a case-control study in Malaysia. *Asian Pac J Cancer Prev*, **10**, 925-32.
- Ramadas A, Kandiah M (2010). Nutritional status and the risk for colorectal adenomas: A case-control study in Hospital Kuala Lumpur, Malaysia. *Pakistan J Nutrit*, **9**, 269-78.
- Suh M, Choi KS, Lee YY, et al (2013). Cancer screening in Korea, 2012: results from the Korean National cancer screening survey. *Asian Pac J Cancer Prev*, **14**, 6459-6463.
- Sugawara Y, Kuriyama S, Kakizaki M, et al (2009). Fish consumption and the risk of colorectal cancer: the Ohsaki cohort study. *Br J Cancer*, **101**, 849-54.
- Sarkar FH, Li Y (2003). Soy isoflavones and cancer prevention. *Cancer Invest*, **21**, 744-57.
- Schoen RE, Tangen CM, Kuller LH, et al (1999). Increased blood glucose and insulin, body size, and incident colorectal cancer. *J National Cancer Inst*, **91**, 1147-54.
- Talaiezhadeh A, Tabesh H, Sattari A, Ebrahimi S (2013). Cancer incidence of southwest of Iran: first report from Khuzestan population-based cancer registry, 2002-2009. *Asia Pac J Cancer Prev*, **14**, 7517-22.
- Takachi R, Tsubono Y, Baba K, et al (2011). Red meat intake may increase the risk of colon cancer in Japanese, a population with relatively low red meat consumption. *Asia Pac J Clin Nutr*, **20**, 603-12.
- Takachi R, Inoue M, Shimazu T, et al (2009). Consumption of sodium and salted foods in relation to cancer and cardiovascular disease: the Japan public health center-based prospective study. *Am J Clin Nutr*, **91**, 456-64.
- Tastan S, Andsoy, II, Iyigun E (2013). Evaluation of the knowledge, behavior and health beliefs of individuals over 50 regarding colorectal cancer screening. *Asian Pac J Cancer Prev*, **14**, 5157-63.
- Ulrich CM (2007). Folate and cancer prevention: a closer look at a complex picture. *Am J Clin Nutr*, **86**, 271-3.
- World Health Organization (WHO) (2015). Cancer.
- Wu XY, Huang XE (2015). Screening for patients with non-small cell lung cancer who could survive long term chemotherapy. *Asian Pac J Cancer Prev*, **16**, 647-652.
- Xue-Yan Wu, Xin-En Huang (2015). Clinical application of serum tumor abnormal protein (TAP) in Colorectal cancer patients. *Asian Pac J Cancer Prev*, **16**, 3425-3428.
- Yusoff HM, Daud N, Noor NM, et al (2012). Participation and barriers to colorectal cancer screening in Malaysia. *Asian Pac J Cancer Prev*, **13**, 3983-7.