

RESEARCH ARTICLE

Protein Profiles Associated with Anoikis Resistance of Metastatic MDA-MB-231 Breast Cancer Cells

Chareeporn Akekawatchai^{1*}, Sittiruk Roytrakul², Suthathip Kittisenachai², Patcharee Isarankura-Na-Ayudhya¹, Sarawut Jitrapakdee³

Abstract

Resistance to anoikis, a cell-detachment induced apoptosis, is one of the malignant phenotypes which support tumor metastasis. Molecular mechanisms underlying the establishment of this phenotype require further investigation. This study aimed at exploring protein expression profiles associated with anoikis resistance of a metastatic breast cancer cell. Cell survival of suspension cultures of non-metastatic MCF-7 and metastatic MDA-MB-231 cells were compared with their adherent cultures. Trypan blue exclusion assays demonstrated a significantly higher percentage of viable cells in MDA-MB-231 than MCF-7 cell cultures, consistent with analysis of annexin V-7-AAD stained cells indicating that MDA-MB-231 possess anti-apoptotic ability 1.7 fold higher than MCF-7 cells. GeLC-MS/MS analysis of protein lysates of MDA-MB-231 and MCF-7 cells grown under both culture conditions identified 925 proteins which are differentially expressed, 54 of which were expressed only in suspended and adherent MDA-MB-231 but not in MCF-7 cells. These proteins have been implicated in various cellular processes, including DNA replication and repair, transcription, translation, protein modification, cytoskeleton, transport and cell signaling. Analysis based on the STITCH database predicted the interaction of phospholipases, PLC and PLD, and 14-3-3 beta/alpha, YWHAB, with the intrinsic and extrinsic apoptotic signaling network, suggesting putative roles in controlling anti-anoikis ability. MDA-MB-231 cells grown in the presence of inhibitors of phospholipase C, U73122, and phospholipase D, FIPI, demonstrated reduced ability to survive in suspension culture, indicating functional roles of PLC and PLD in the process of anti-anoikis. Our study identified intracellular mediators potentially associated with establishment of anoikis resistance of metastatic cells. These proteins require further clarification as prognostic and therapeutic targets for advanced breast cancer.

Keywords: Anoikis resistance - MDA-MB-231 - MCF-7 - proteomics - GeLC-MS/MS - STITCH

Asian Pac J Cancer Prev, 17 (2), 581-590

Introduction

Metastasis and invasion are major causes of death from cancers. They are multi-step processes including dissociation of cancer cells from a primary site, survival in the circulatory and lymphatic systems, and colonization in the distant organs. Generally, an epithelial or endothelial cell undergoes a particular type of apoptosis, termed anoikis, in the absence of attachment to extracellular matrix (ECM) or neighboring cells. Cancer cells acquire anoikis resistance to survive after the detachment from ECM and metastasizes to the target organs (Paoli et al., 2013). Previous studies have shown that an acquisition of anoikis resistance is a critical prerequisite for cancer metastasis, and there is a link between an anchorage-independent growth and survival of circulatory tumor cells, and metastatic ability of the cells in orthotopic cancer models (Glinskii et al., 2003; Kim et al., 2012; Malin et al., 2015). The molecular and functional analyses of anokis resistant cancer cells are essential for the understanding

of the cellular basis of cancer metastasis and invasion.

Growing evidence have indicated that cancer cells achieve resistance to anoikis by several mechanisms including alternations of integrin repertoire, constitutive activation of pro-survival and proliferative signaling, epithelial mesenchymal transition (EMT) and metabolic reprogramming (Kim et al., 2012; Paoli et al., 2013). These events are mediated by the activation of different intracellular signaling networks, which in turn lead to anoikis resistance (Paoli et al., 2013). Following ECM detachment, the hyperactivation of receptor tyrosine kinase growth factor receptors, changes in integrin expression pattern and down-regulation of PTEN result in the activation of pro-survival phosphoinositide 3-kinase (PI3K)/AKT signaling systems. This subsequently inhibits apoptotic process, while deregulation of E-cadherins during EMT elicits the migration of β -integrin to the nucleus, where it stimulates expression of target genes responsible for cell motility and invasion (Paoli et al., 2013). Tumor cells reprogram metabolic pathways by oxidizing glucose

¹Department of Medical Technology, Faculty of Allied Health Sciences, Thammasat University, ²National Center for Genetic Engineering and Biotechnology, National Science and Technology Development Agency, Pathumtani, ³Department of Biochemistry, Faculty of Sciences, Mahidol University, Bangkok, Thailand *For correspondence: cakekawatchai@gmail.com, ejarepo@tu.ac.th

via anaerobic glycolysis known as Warburg effect. This Warburg effect is regulated via activation of PI3K/Akt signaling cascade, hypoxia-inducible factor-1 (HIF-1), p53 and AMP-activated protein kinase (AMPK)-liver kinase B-1 (LKB-1) pathways (Kim et al., 2012; Buchheit et al., 2014). Although, various signaling pathways underlying anoikis resistance have been reported, mechanistic players which serve as therapeutic targets for cancer metastasis are still very limited.

Breast cancer cell lines, the estrogen receptor-positive (ER⁺) non-metastatic MCF-7 and ER⁻ metastatic MDA-MB-231 breast cancer cell lines have been well-characterized for their aggressiveness by both *in vitro* and *in vivo* assays (Thompson et al., 1992; Culty et al., 1994; Holliday and Speirs, 2011). Several studies have demonstrated that the malignant phenotypes in the two cell lines are associated with different molecular characters (Eckert et al., 2004; Akekawatchai et al., 2005; Holliday and Speirs, 2011; Niu et al., 2013; Hooshmand et al., 2014). Particularly, identification of biomarkers for tumorigenesis and invasiveness of breast cancer has been studied in these two cell lines (Nagaraja et al., 2006; Lai et al., 2010). In order to explore protein expression profiles responsible for the establishment of anoikis resistance in breast cancer cells, the anchorage-independent survival of these two cell lines was the subject of this study. A comparative proteomic analysis was then conducted to identify protein candidates that are required for anti-anoikis property of breast cancer cells. The proteins as well as their predicted function found in this study provided more evidence to understand the molecular mechanisms involved in anoikis resistance of the metastatic cancer cells.

Materials and Methods

Cell lines and maintenance

The non-metastatic breast cancer MCF-7 and metastatic MDA-MB-231 cell lines (obtained from the American Type Culture Collection) were kindly provided by Professor Shaun R. McColl (The University of Adelaide, South Australia) (Akekawatchai et al., 2005; Niu et al., 2013). The two cell lines were cultured in Roswell Park Memorial Institute-1640 (RPMI-1640) medium (GibcoBRL, NY) supplemented with 10% (v/v) fetal bovine serum (GibcoBRL, NY), 10 mM HEPES, 100 units penicillin and 100 µg/ml streptomycin, at 37°C in a 5% CO₂ atmosphere.

Anchorage-independent culture of MCF-7 and MDA-MB-231 cell lines, and treatments

Anchorage-independent cell growth was performed by culturing the cells on anti-adhesive polymer, poly-2-hydroxyethyl methacrylate (polyHEMA)-coated 6-well plates containing RPMI-1640 medium supplemented with 10% FBS. Briefly, culture plates were prepared by adding 0.5 ml of 5 mg/ml polyHEMA (Sigma-Aldrich) in 95% ethanol into 6-well plates and dried for 2 days with lids in place followed by extensive washes with PBS (Fukazawa and Uehara, 2000). Cells were seeded at a density of 5x10⁵ per well and cultured in RPMI-1640 medium supplemented with 10% FBS for 24, 48 and 96 hours. At

specific time points, the adherent cells were trypsinized and suspended in medium whereas the detached cells were collected. The cell suspension was then subjected to assess viability by trypan blue exclusion assay and cell death by the annexin V and dead cell assay.

To determine an involvement of phospholipase activity in survival of suspended cells, the cells were cultured in the medium supplemented with various concentrations of specific inhibitors of phospholipase C (PLC), U-73122 (Santa Cruz Biotechnology, CA), or phospholipase D (PLD), 4-Fluoro-N-(2-(4-(5-fluoro-1H-indol-1-yl) piperidin-1-yl) ethyl) benzamide, 5-Fluoro-2-indolyl des-chlorohalopemide (FIPI) hydrochloride hydrate (FIPI) (Sigma Aldrich) compared with control cultures supplemented with dimethyl sulfoxide (DMSO) and DMSO:methanol (MeOH), respectively for 24 hours and then subjected to the annexin V and dead cell analysis. A percentage change of apoptotic cells in inhibitor treated cultures were calculated relative to that in control cultures.

Detection of cell viability by trypan blue exclusion assay

The cell suspension was stained with 0.4% trypan blue in PBS (ratio of 1:1) for 5 minutes before examined under a microscope. A viable cell incapable of excluding the dye shows blue stain while a non-viable cell shows a clear cytoplasm. The percentage of viable cells in anchorage-independent culture was determined in relative to adherent culture.

Cell death analysis by the annexin V and dead cell assay

The cell suspension was analyzed by a Muse™ Annexin V and Dead Cell kit (Merk Millipore Co., Darmstadt, Germany) according to the manufacturer's protocol. Briefly, the cells were washed twice in ice-cold phosphate buffer saline (PBS) and resuspended at a concentration of 5 x 10⁵ cells/ml in 100 µl binding buffer (10 mM HEPES/NaOH (pH 7.4), 140 mM NaCl, 2.5 mM CaCl₂). The cells were then incubated with 100 µl of the Muse™ Annexin V and Dead Cell reagent (7-aminoactinomycin D; 7-AAD) at room temperature for 20 minutes in the dark before subjected to a Muse™ cell analyzer. The survival cells was determined as the percentage of annexin V⁻/7-AAD⁻ cells whereas that of apoptotic cells included the percentages of early (annexin V⁺/7-AAD⁻) and late apoptotic (annexin V⁺/7-AAD⁺) cells.

Protein extraction for GeLC-MS/MS

The cultured cells at 24 hours were harvested for protein extraction. The adherent cells were scraped into 2 ml of cold PBS whereas suspended cells were collected. Cells were centrifuged at 3,000 rpm for 5 minutes, and the pellet was washed once in 1 ml of PBS. The cell pellet was stored at -80°C until use. Subsequently, the frozen cells were thawed and centrifuged at 9,200 x g for 10 minutes, and the pellet was resuspended in 0.5% (w/v) sodium dodecylsulfate (SDS) in water. Protein concentration was measured by Lowry's method and 50 µg of protein samples were subjected to 12.5% SDS-polyacrylamide gel before staining with colloidal Coomassie blue G. Each lane was sliced in to 1 x 1 x 1 mm³ sized gel plugs.




Figure 1. A higher degree of anti-apoptotic ability of MDA-MB-231 than that of MCF-7 cells in the anchorage-independent culture. The survival of cells grown under adherent and anchorage-independent conditions was examined A, representative photographs showing morphologies of the suspended and adherent MCF-7 and MDA-MB-231 cells cultured for 48 hours (x100). The viability of cells cultured for 24, 48 and 94 hours was analyzed by typan blue inclusion assay (C) and the survival of cells cultured at 24 hour cell cultures was also determined by flow cytometry of annexin V-7-AAD double stained cells (B and D). Data are presented as the mean $\pm$ S.E.M of % cells relative to adherent culture; n=5 (C) and n=4-5 (D) experiments each performed in duplicate. Asterisks indicate statistical difference between the percentages of viable or apoptotic MCF-7 and MDA-MB-231 cells (Student's unpaired t test) where *, $p < 0.05$ and **, $p < 0.005$




Figure 2. Venn diagram and heatmap analysis of proteins differentially expressed in anchorage-independent MCF-7 and MDA-MB-231 compared with the adherent cells. The extracts of suspended and adherent cells cultured for 24 hours were subject to GeLC-MS-MS analysis. A and B show the diagrams of 925 proteins differentially expressed in suspended MCF-7 and MDA-MB-231 cells compared with the adherent cells. A, numbers of proteins expressed in suspended or adherent MCF-7 and MDA-MB-231 cells are shown. B, column 1 and 2 represent protein expression in adherent and suspended MCF-7 whereas column 3 and 4 represent that in adherent and suspended MDA-MB-231 cells respectively. Green, black and red colors represent proteins with low, average and high levels of expression respectively

In-gel digestion

The gel plugs were placed in 96 well-plate and washed with 200 μ l/well of sterile distilled water. The plate was shaken for 5 minutes at room temperature and the gel plugs were destained with 25 mM ammonium bicarbonate (NH_4HCO_3) in 50% methanol until they become transparent. After the removal of destaining

solution and three washes with sterile distilled water, samples were dehydrated with 100% acetonitrile (ACN), reduced with 10 mM dithiothreitol (DTT) in 10 mM NH_4HCO_3 at room temperature for 1 hour and alkylated with 100 mM iodoacetamide (IAA) in 10 mM NH_4HCO_3 at room temperature for 1 hour in the dark. The gel slices were then washed twice with 100% ACN for 5 minutes.

Table 1. A list of 21 proteins expressed in suspended and adherent MDA-MB-231, but not in MCF-7 cells

Database	Protein name	Protein ID score	Peptides	Biological process	Subcellular distribution
ID no.					
gil386730457	cib gene product	15.56	AADMLAEYERR	cytoskeleton	membrane
gil24417711	nesprin-2	16.31	KAQDLTSLK	cytoskeleton	Membrane/ cytoskeleton
gil323498894	4-alpha-glucanotransferase	17.65	KFENAAVQK	Metabolism	cytoplasm
gil7662018	PHD finger protein 3	28.98	KIVAAK	Transcription	nucleus
gil42490753	transcriptional enhancer factor TEF-5	10.18	TIASSRLR	Transcription	nucleus
gil56204959	cleavage and polyadenylation specific factor 3-like	10.46	IAVDK	Transcription	cytoplasm
gil323302928	Tuf1p	16.13	LLTRTAFKAXGKLLR	Translation	Cytoplasm
gil188591432	transposase, IS6 family	23.28	TPRTAPEQVQR	Epigenic transcription/ transposition	nucleus
gil334117440	glycosyltransferase family 2	12.32	AFKEDLSMALLR	Protein modification	Golgi/ER
gil330808175	Phospholipase	12.92	LGLSLSDDMR	Signal transduction	cytoplasm
gil337729402	phage related protein	0.2	INMADK	Signal transduction	cytoplasm
gil51094943	Piccolo protein (Aczonin)	7.1	RDSSSSSLRLK	Signal transduction	membrane/ cell junction
gil52317138	olfactory receptor 5R1	12.93	MMVNFVVER	Signal transduction	membrane
gil386019860	acrB gene product	10.52	KMPKRTDIK	Signal transduction	membrane
gil57167934	amino acid ABC transporter, ATP-binding protein	10.32	IFFMNK	Transport	membrane
gil126362955	sodium channel protein type 3 subunit alpha isoform 3	6.83	NSNGTTTETEVRK	Transport	membrane
gil157388921	transmembrane and coiled-coil domain-containing protein 3 precursor	19.65	AAITR	Transport	membrane
gil333929654	outer membrane autotransporter barrel domain-containing protein	23.13	ASAEVAVNK	Transport	membrane
gil300869897	carB gene product	12.14	VADVASVGLGRAPR	Stress response	mitochondria
gil14133207	KIAA0556 protein	8.54	PEPPPLGMDGQTLR	Unknown	unknown
gil327249992	phage tail fibre repeat family protein	14.93	GRALLAHDTAEAMR	Unknown	unknown

In-gel digestion was performed by adding 10 ng/ml trypsin in 50% ACN/10 mM NH_4HCO_3 to the gels and incubated at room temperature for 20 minutes. The gels were kept immersed throughout the digestion by adding 30% ACN before further incubated at 37°C overnight. To extract the digested peptides, 50% ACN in 0.1% formic acid (FA) was added into the gels and incubated for 10 minutes in a shaker. The peptide products were collected, vacuum dried and stored at -80°C for further mass spectrometric analysis.

LC-MS/MS

The tryptic peptides were analyzed using a HCTultra PTM Discovery System (BrukerDaltonics Ltd, U.K./Germany) coupled to an UltiMate 3000 LC System (Dionex




Figure 3. Subcellular distribution and functional annotation of proteins expressed in adherent and suspended MDA-MB-231 cells. Subcellular localization (A) and function (B) of 54 proteins expressed in adherent and suspended MDA-MB-231, but not MCF-7 cells, were obtained from the online Gene Ontology (Go) cat, Uniprot and STITCH 4.0 databases.

(<http://uniprot.org>), STITCH 4.0 (<http://stitch.embl.de/>) databases. Protein-protein interaction was analyzed according to STITCH 4.0 database (<http://stitch.embl.de/>).

Statistic analysis

Data were analyzed by unpaired student's t-test using Graphpad Prism Software (San Diego, CA, USA). In all analyses, *p* values less than 0.05 were considered statistically significant.

Results

MDA-MB-231 cells grown under an anchorage-independent condition show anoikis property higher than MCF-7 cells.

The ability of MDA-MB-231 and MCF-7 cells to grow in an anchorage-independent condition was initially compared. As shown in Figure 1, attachment prevention of the two cell lines altered their morphologies observed under a microscope. Although MCF-7 cells formed a large cluster of spheroid cells within 24 hours (Data not shown), the poorly formed aggregates of dead or dying cells were observed in 48 hours later (Figure 1A). In contrast, MDA-MB-231 cells rapidly formed multicellular spheroids which were more sustainable as shown in Figure 1A. Trypan blue exclusion assay showed that the percentage of viable MCF-7 cells was dramatically decreased within 24 hours of culture whereas that of MDA-MB-21 cells was gradually decreased. The significantly higher percentages of viable MDA-MB-231 cells were observed in 24 and 48 hours of the culture (Figure 1C). Flow cytometric

analysis of annexin V-7-AAD double stained cells at 24 hours of culture shown in Figure 1B and 1D demonstrated attachment prevention-induced apoptosis in the two cell lines and a 1.7 fold lower percentage of apoptotic cells relative to adherent culture in MDA-MB-231 than that in MCF-7 cell cultures. These data suggested a significant higher degree of anchorage-independent survival ability of MDA-MB-231 cells.

Protein profiles of MDA-MB-231 and MCF-7 cells grown under anchorage-independent and adherent conditions analyzed by GeLC-MS/MS

In order to explore protein expression profiles involving in the anoikis resistance property of MDA-MB-231 breast cancer cells, the suspended and adherent cells cultured at 24 hours was analyzed by GeLC-MS/MS due to the highest degree of difference between the viability of MDA-MB-231 and MCF-7 cells. A two-way hierarchical clustering analysis shown in Figure 2B indicated that 925 proteins were differentially expressed in the two cell lines grown under suspended or adherent conditions. A Venn diagram showed numbers of proteins that were expressed in the two culture conditions of MCF-7 and MDA-MB-231 cells (Figure 2A). Of particular, there were 54 proteins which were expressed only in MDA-MB-231, but not in MCF-7 cells, 33 proteins were found only in the adherent cells whereas 21 proteins remained expressed in the detached cells. Thus, further analysis for their subcellular distribution and functional annotation was performed by the Gene Ontology (Go) cat, Uniprot and STITCH 4.0 as shown in Figure 3 A

Table 2. A list of 33 Proteins Expressed in Adherent MDA-MB-231, but not in MCF-7 cells, and Down-regulated in Detached MDA-MB-231 Cells

Database ID no.	Protein name	Protein ID score	Peptides	Biological process	Subcellular distribution
gil18088719	beta-tubulin	59.84	ISVYYNEATGGK	cytoskeleton	cytoplasm
gil29789403	PDZ domain-containing protein 8	21.7	EGGLDDSVFIAVK	cytoskeleton	cytoplasm
gil4758178	cytoplasmic dynein 1 intermediate chain 1 isoform a	12.35	SHR RTPVQR	cytoskeleton, cell movement, migration	cytoplasm
gil119596348	junctophilin 2, isoform CRA_c	2.21	RAAMGR	cytoskeleton	membrane
gil21245106	lipase member H precursor	8.87	VAMVLK	lipid metabolism	membrane
gil2906146	malate dehydrogenase precursor	36.51	AGAGSATLSMAYAGAR	metabolism	cytoplasm/ mitochondria
gil321313747	dihydropteroate synthase	8.22	QIARMAKMM DAMLNK	metabolism	cytoplasm
gil301017356	glycosyltransferase, group 1 family	12.48	TINLSSERK	metabolism	membrane
gil297620850	vesicle-fusing ATPase	14.82	GGSVNK	Transport	Cytoplasm
gil329938632	DNA gyrase subunit A	5.21	VGGIADVR	DNA replication	nucleus
gil320178239	ATP-dependent helicase hrpA	11.31	KGRSGR	DNA replication	Nucleus
gil76496497	protein artemis isoform a	10.7	GLRAPTLKRR	DNA repair	Nucleus
gil116052873	cbrB gene product	5.56	SGASADS	Transcription	nucleus

Database ID no.	Protein name	Protein ID score	Peptides	Biological process	Subcellular distribution
gil119597574	AT hook containing transcription factor 1, isoform CRA_d	1.05	LIRTY	Transcription	Nucleus
gil187761317	methylcytosine dioxygenase TET2 isoform a	2.79	LPPNLSNPNDYK	Transcription	nucleus
gil333985858	restriction modification system DNA specificity domain-containing protein	22.44	KRKLVELLK	Transcription	nucleus
gil154744870	zinc finger protein 574	13.26	TFVNLTK	Transcription	Neucleus
gil333396626	transcriptional regulator CtsR	11.79	LYDDQVIERK	Transcription	Neucleus
gil300253409	heat shock protein 70-p1	19.14	TTPSYVAFTESER	Transcription	nucleus/ cytoplasm
gil306836586	cse1 family CRISPR-associated protein	4.83	EGVAHA	transcription	Nucleus
gil4506609	60S ribosomal protein L19	9.95	KGTANAR	Translation	Cytoplasm
gil337751588	histidinol-phosphate aminotransferase	11.76	KNGGPA	Translation	Cytoplasm
gil133777378	NMT1 protein	7.77	ALNSS	Protein modification	Cytoplasm
gil300245991	carboxylase-like protein	18.84	ESGGAASYLVI-SIRNYGAAGLAK	Protein modification	plasma mem- brane
gil4504237	semaphorin-7A isoform 1 preproprotein	12.78	AAPSAPR	Signal transduction	membrane
gil8923394	transmembrane protein 161A isoform 1 precursor	8.54	ALAGKPRPR	Signal transduction	membrane
gil386316270	pyrG gene product	31.38	ADMTEWSAMVDK	Signal transduction	membrane
gil1072378	PEP-19	9.05	QGAGPTNGKDK	Signal transduction	Cytoplasm/ Neucleus
gil4507949	14-3-3 protein beta/alpha	34.12	NLLSVAYK	Signal transduction	cytoplasm
gil307544419	bclI gene product	10.76	IGGVLSIGPK	Signal transduction	Cytoplasm
gil119623684	FLJ45422 protein, isoform CRA_b	5.11	MGVMAPR	Signal transduction	nucleus
gil333369681	cytochrome c, class I	30.77	AQKGMPALVESVR	Signal transduction	cytoplasm
gil1469874	KIAA0146	10.99	TPENSAKKK	Unknown	uncharacterised

and B, respectively. These proteins were involved in various cellular processes, including cell signaling (24%), transcription (22%), metabolism (9%), cytoskeleton/structure (9%), transport (9%), protein modification (6%), translation (5%), stress response (4%), DNA replication and repair (4%) and unknown function (6%) and they were located in cytoplasm (33%), membrane (30%), nucleus (28%), golgi apparatus and endoplasmic reticulum (2%), mitochondria (2%) and unidentified location (5%). A full list of 21 proteins which are expressed both in the suspended and adherent MDA-MB-231 cells is shown in Table 1 whereas 33 proteins which were found only in the adherent cells are shown in Table 2. Our data demonstrated the differential protein expression of the two cell lines in response to attachment prevention and there were 54

proteins found in detached and adherent MDA-MB-231, but not in MCF-7, cells that potentially involved in the higher degree of anti-anoikis of the highly metastatic MDA-MB-231 cells.

STITCH analysis for an association of candidate proteins with the apoptotic signaling network in the highly metastatic MDA-MB-231 cells

To identify signal transduction pathways relevant to the anoikis resistance phenotype of the MDA-MB-231 cells, a total of 54 proteins expressed only in the suspended and adherent MDA-MB-231, were analyzed for their interaction with the classical extrinsic and intrinsic apoptotic signaling molecules, TNF receptor superfamily, member 6 (Fas), BCL-2-associated X protein (Bax),

B-cell lymphoma 2 (BCL2) and apoptosis-related cysteine peptidases (caspases) according to online STITCH 4.0 database (Kuhn et al., 2014). The analysis was based on the following parameters: species, Homo sapiens; confidence score, medium; active prediction methods, all and using a list of multiple protein names as input. As shown in Figure 4, the 12 candidate proteins including domain family member (TEAD3), zinc finger protein 574 (ZNF574), tubulin tyrosine ligase-like family, member 1 (TTLL1), PHD finger protein 3 (PHF3), amylo-1, 6-glucosidase, 4-alpha-glucanotransferase (AGL), DNA repair-scaffolding protein KIAA0146 (KIAA0146), olfactory receptor, family 5, subfamily R, member 1 (OR5R1), PDZ domain containing 8 (PDZD8), zinc finger protein 280D (ZNF280D), terminal uridylyl transferase 1 (TUT1), N-ethylmaleimide-sensitive factor (NSF) and ribosomal protein L19 (RPL19) did not show any interaction network with apoptotic pathways. However, the analysis predicted contribution of phospholipases, PLCG1, PLCG2, PLD2 and 14-3-3 beta/alpha, YWHAB, to the apoptotic network. Apparently, the phospholipase isoforms, PLC and PLD, remained expressed in the detached cells (Table 1) and were predicted to be downstream of epidermal growth factor receptor (EGFR) that are negative regulator of the extrinsic Fas-FasL and intrinsic Bax-related signaling pathways. Another protein, 14-3-3 beta/alpha, YWHAB, which is down-regulated in response to cell detachment (Table 2) was found to be associated with BH3 interacting domain death agonist (BID) and BCL2-associated agonist of cell death (BAD). The STITCH analysis suggests that constitutive expression of phospholipase isoenzymes, and the down-regulation of 14-3-3 beta/alpha, YWHAB, potentially play roles in the regulation of apoptotic signaling pathways in response to the detachment of MDA-MB-231 cells.

Phospholipases is essential for the anchorage-independent survival of MDA-MB-231 and MCF-7 cells

Previous studies have clearly demonstrated higher expression levels and activity of PLC and PLD isoforms in MDA-MB-231 than those in MCF-7 cells (Zhong et al., 2003; Abalsamo et al., 2012). To validate our proteomics data and determine roles of PLC and PLD activity in supporting anoikis resistant property, both cell lines were grown under an anchorage-independent culture condition in the presence of, U-73122 or FIPI, which are inhibitors of PLC or PLD, respectively (Kumar et al., 2012; Henkels et al., 2013). The percentages of apoptotic cells represented by annexin V⁺/7-AAD⁻ and annexin V⁺/7-AAD⁺ cells were measured. Figure 5 A and B show percentage changes of the annexin V⁺/7-AAD⁻ and annexin V⁺/7-AAD⁺ cells in U-73122 and FIPI-treated cultures of MDA-MB-231 and MCF-7 cells relative to their adherent culture conditions. The data indicated that both U-73122 and FIPI treatment increased percent apoptotic cells in a dose-dependent manner in both cell lines. We also observed a higher percentage increase of apoptotic cells in the anchorage-independent culture conditions of specific PLC and PLD inhibitor-treated MDA-MB-231 than that of MCF-7 cells. These data suggest a stronger contribution of PLC and PLD activity in supporting anchorage-independent

survival of MDA-MB-231 than MCF-7 cells.

Discussion

The non-metastatic MCF-7 and metastatic MDA-MB-231 breast cancer cell lines have long been widely used in breast cancer research (Holliday and Speirs, 2011). *In vitro* and *in vivo* studies have shown considerable differences in the invasive and metastatic abilities (Thompson et al., 1992; Culty et al., 1994). The significant difference in their cellular and molecular characteristics has also been demonstrated in many studies (Eckert et al., 2004; Akekawatchai et al., 2005; Holliday and Speirs, 2011; Niu et al., 2013; Hooshmand et al., 2014). The two cell lines have been shown to be resistant to anoikis (Mahadev et al., 2002; Bharadwaj et al., 2005; Kim et al., 2010). However, the studies demonstrating a comparison of their anchorage-independent survival are still very limited. Our data showed the significantly higher survival ability of MDA-MB-231 than that of MCF-7 cells in response to attachment prevention, which is consistent with the invasive and metastatic abilities (Thompson et al., 1992; Culty et al., 1994). Previous studies also indicated the differential gene expression and proteomic profiles of the two cell lines and identified proteins which can serve as potential markers for tumorigenesis and invasiveness of breast cancer (Nagaraja et al., 2006; Lai et al., 2010). Therefore, the two breast cancer cell lines were used to comparatively analyze putative mediators involving anoikis resistance in response to ECM detachment.

As proteomics is a potential tool widely used in searching for proteins as well as their functionality in different cellular systems (Chesor et al., 2014; Kuhn et al., 2014; Laothumthut et al., 2015), the expression of the two cell lines in response to cell detachment was analyzed by the GeLC-MS/MS technique. Our data revealed that 925 proteins were expressed differentially in suspended or adherent cultures of MCF-7 and MDA-MB-231 cells, indicating their involvement in the attachment prevention. Of particular, 54 out of 925 proteins found only in the detached and adherent MDA-MB-231 cells, of which 33 proteins are down-regulated whereas 21 proteins remained expressed in response to the attachment prevention. These proteins were identified to be associated with various functions including cytoskeleton/structure, cell metabolisms, DNA replication and repair, transcription and translation, protein modification, cell signaling, transport and stress response, all of which potentially support ability of the cell to resist anoikis.

Previous studies indicated that cancer cells may employ different strategies to counteract the apoptotic pathways in response to the detachment from ECM, leading to anchorage-independent growth (Paoli et al., 2013). Thus, an association of the proteins identified in this study with the classical apoptotic signaling pathways was examined by STITCH 4.0. This demonstrates that there was no interaction of the 12 candidate proteins including the remaining expressed proteins, TTLL1, AGL, PHF3, TEAD3, OR5R1, and the down-expressed proteins, ZNF574, TUT1, KIAA0146, PDZD8, ZNF280D, NSF and RPL19 to the apoptotic signaling network in

MDA-MB-231 cells. There are possible explanations for the findings. Some of these proteins have not been well-characterized and, therefore, no data available in the database. Additionally, although these proteins may have no direct association with anoikis ability of the cells, their involvement in other activities supporting the anchorage-independent survival could not be ruled out. In contrast, our data indicated the interaction of phospholipase isoforms, PLC and PLD, which remained expressed in MDA-MB-231 cells, and 14-3-3 beta/alpha, YWHAB, which was down-expressed in response to detachment of the cells, with the intrinsic and extrinsic apoptotic signaling network. Whereas the expression of phospholipases potentially participate in the down-regulation of apoptotic signal transduction supporting the cells to overcome anoikis in response to the attachment prevention, the down-regulation of the 14-3-3 beta/alpha may support a certain degree of apoptotic cell death observed in the MDA-MB-231 cells. It is a very first time to specifically implicate these signaling proteins in anoikis resistance of the highly metastatic MDA-MB-231 cells.

Several previous studies have characterized expression and function of PLC and PLD isoforms in MCF-7 and MDA-MB-231 cells (Zhong et al., 2003; Abalsamo et al., 2012). Our proteomics analysis demonstrated that there was a constitutive expression of phospholipases observed only in MDA-MB-231, but not in MCF-7 cells and remained expressed in response to the attachment prevention. Inhibitor experiments also suggested the involvement of PLC and PLD in anoikis resistance of the two cell lines and a stronger contribution of PLC and PLD activity in anti-anoikis of MDA-MB-231 than those in MCF-7 cells. These data are consistent with several previous studies demonstrating the higher expression levels and activity of different phospholipase isoforms including phosphatidylcholine-PLC (PC-PLC) and PLD in MDA-MB-231 than those in MCF-7 cells. The activities of these isoforms were apparently essential for maintaining the mesenchymal-like phenotype, anti-apoptosis induced by serum withdrawal and, migration and invasion of the cells (Zhong et al., 2003; Zheng et al., 2006; Abalsamo et al., 2012).

Generally, PLC and PLD are phospholipid-hydrolyzing isoforms which can generate many intracellular bioactive lipid mediators which in turn regulate various cellular processes, leading to tumorigenesis, migration and invasion (Park et al., 2012). Both isoforms can be activated in response to the stimulation of mitogenic signals including epidermal growth factors (EGF) through their respective receptor tyrosine kinase EGFRs (Roskoski, 2014). It appears that the interaction of SH-2 domain of phospholipase C gamma (PLCG) with the phosphotyrosine residues on the activated RTKs can activate PLCG-mediated proliferative signaling via the protein kinase B-induced gene expression (Park et al., 2012; Roskoski, 2014). The growth factor signaling also increases the activity of phospholipase D (PLD) isoforms that generate phosphatidic acid and recruit RAF to plasma membrane to activate the mitogen-activated protein kinase (MAPK) signaling pathway (Rizzo et al., 1999). In this study, analysis based on STITCH database

predicted that PLC and PLD are potentially downstream of epidermal growth factor receptor (EGFR) that are negative regulator of the extrinsic Fas-FasL and intrinsic Bax-related signaling pathways (Figure 4), consistent with the evidence demonstrating the EGFR expression and activity of Fas-FasL-mediated apoptotic signaling in the MDA-MB-231 cells (Pledge-Tracy et al., 2007; Cui et al., 2013). Thus, it is highly possible that the metastatic MDA-MB-231 cells utilize growth factor signaling through PLC and PLD to counteract the apoptotic signaling pathways in response to the ECM detachment. While accumulating studies indicated that the PI3K signaling system plays a central role in anti-anoikis of a cell (Park et al., 2012), our data have shown a potential involvement of the phospholipase-dependent signaling system in overcoming anoikis of this cell type.

In conclusion, this study has demonstrated the significantly higher degree of anchorage-independent survival in the metastatic MDA-MB-231 than that in non-metastatic MCF-7 cells. The comparative proteomic analysis indicated the differential protein expression profiles in the two cell lines in response to attachment prevention. In particular, there were 54 identified proteins expressed only in adherent and detached MDA-MB-231, but not in MCF-7 cells, that potentially contribute to the high level of anti-apoptotic ability of the MDA-MB-231 cells. Analysis of these identified proteins according to STITCH database together with the functional experiments suggested that this cell type potentially exploits the growth factor signaling through phospholipases, PLC or PLD, to counteract the apoptotic process induced in response to the ECM detachment. In contrast, STITCH analysis indicated that the up-regulation of apoptotic signaling pathways in response to detachment of the cells may be promoted by the down-regulation of 14-3-3 beta/alpha, YWHAB, regulatory protein. These signaling mediators as well as other identified proteins reported in this study require more clarification in terms of their expression and function in the establishment of anoikis resistance of the metastatic breast cancer cells and may serve as biomarkers and therapeutic targets for the advanced breast cancer.

Acknowledgements

This study was financially supported by Thailand Research Fund (TRF) and Commission on Higher Education (CHE), MRG5280163 and Thammasat University (research fund academic year of 2009). The authors declare that they have no conflict of interest.

References

- Abalsamo L, Spadaro F, Bozzuto G, et al (2012). Inhibition of phosphatidylcholine-specific phospholipase C results in loss of mesenchymal traits in metastatic breast cancer cells. *Breast Cancer Res*, **14**, 50.
- Akekawatchai C, Holland JD, Kochetkova M, et al (2005). Transactivation of CXCR4 by the insulin-like growth factor-1 receptor (IGF-1R) in human MDA-MB-231 breast cancer epithelial cells. *J Biol Chem*, **280**, 39701-8.
- Bharadwaj S, Thanawala R, Bon G, et al (2005). Resensitization of breast cancer cells to anoikis by tropomyosin-1: role

- of Rho kinase-dependent cytoskeleton and adhesion. *Oncogene*, **24**, 8291-303.
- Buchheit CL, Weigel KJ, Schafer ZT (2014). Cancer cell survival during detachment from the ECM: multiple barriers to tumour progression. *Nat Rev Cancer*, **14**, 632-41.
- Chesor M, Roytrakul S, Graidist P, et al (2014). Proteomics analysis of siRNA-mediated silencing of Wilms' tumor 1 in the MDA-MB-468 breast cancer cell line. *Oncol Rep*, **31**, 1754-60.
- Cui W, Zhang S, Shan C, et al (2013). microRNA-133a regulates the cell cycle and proliferation of breast cancer cells by targeting epidermal growth factor receptor through the EGFR/Akt signaling pathway. *FEBS J*, **280**, 3962-74.
- Culty M, Shizari M, Thompson EW, et al (1994). Binding and degradation of hyaluronan by human breast cancer cell lines expressing different forms of CD44: correlation with invasive potential. *J Cell Physiol*, **160**, 275-86.
- Eckert LB, Repasky GA, Ulku AS, et al (2004). Involvement of Ras activation in human breast cancer cell signaling, invasion, and anoikis. *Cancer Res*, **64**, 4585-92.
- Fukazawa H, Uehara Y (2000). U0126 reverses Ki-ras-mediated transformation by blocking both mitogen-activated protein kinase and p70 S6 kinase pathways. *Cancer Res*, **60**, 2104-7.
- Glinskii AB, Smith BA, Jiang P, et al (2003). Viable circulating metastatic cells produced in orthotopic but not ectopic prostate cancer models. *Cancer Res*, **63**, 4239-43.
- Henkels KM, Boivin GP, Dudley ES, et al (2013). Phospholipase D (PLD) drives cell invasion, tumor growth and metastasis in a human breast cancer xenograph model. *Oncogene*, **32**, 5551-62.
- Holliday DL, Speirs V (2011). Choosing the right cell line for breast cancer research. *Breast Cancer Res*, **13**, 215.
- Hooshmand S, Ghaderi A, Yusoff K, et al (2014). Differentially expressed proteins in ER+ MCF7 and ER- MDA-MB-231 human breast cancer cells by RhoGDI-alpha silencing and overexpression. *Asian Pac J Cancer Prev*, **15**, 3311-7.
- Kim JB, Yu JH, Ko E, et al (2010). The alkaloid Berberine inhibits the growth of Anoikis-resistant MCF-7 and MDA-MB-231 breast cancer cell lines by inducing cell cycle arrest. *Phytomedicine*, **17**, 436-40.
- Kim YN, Koo KH, Sung JY, et al (2012). Anoikis resistance: an essential prerequisite for tumor metastasis. *Int J Cell Biol*, **2012**, 306879.
- Kuhn M, Szklarczyk D, Pletscher-Frankild S, et al (2014). STITCH 4: integration of protein-chemical interactions with user data. *Nucleic Acids Res*, **42**, 401-7.
- Kumar B, Kumar A, Ghosh S, et al (2012). Diospyrin derivative, an anticancer quinonoid, regulates apoptosis at endoplasmic reticulum as well as mitochondria by modulating cytosolic calcium in human breast carcinoma cells. *Biochem Biophys Res Commun*, **417**, 903-9.
- Lai TC, Chou HC, Chen YW, et al (2010). Secretomic and proteomic analysis of potential breast cancer markers by two-dimensional differential gel electrophoresis. *J Proteom Res*, **9**, 1302-22.
- Laothumthut T, Jantarat J, Paemane A, et al (2015). Shotgun proteomics analysis of proliferating STRO-1-positive human dental pulp cell after exposure to nacreous water-soluble matrix. *Clin Oral Invest*, **19**, 261-70.
- Mahadev K, Raval G, Bharadwaj S, et al (2002). Suppression of the transformed phenotype of breast cancer by tropomyosin-1. *Exp Cell Res*, **279**, 40-51.
- Malin D, Strekalova E, Petrovic V, et al (2015). ERK-regulated alphaB-crystallin induction by matrix detachment inhibits anoikis and promotes lung metastasis *in vivo*. *Oncogene*, **34**, 5626-34.
- Nagaraja GM, Othman M, Fox BP, et al (2006). Gene expression signatures and biomarkers of noninvasive and invasive breast cancer cells: comprehensive profiles by representational difference analysis, microarrays and proteomics. *Oncogene*, **25**, 2328-38.
- Niu M, Klingler-Hoffmann M, Brazzatti JA, et al (2013). Comparative proteomic analysis implicates eEF2 as a novel target of PI3Kgamma in the MDA-MB-231 metastatic breast cancer cell line. *Proteome Sci*, **11**, 4.
- Paoli P, Giannoni E, Chiarugi P (2013). Anoikis molecular pathways and its role in cancer progression. *Biochim Biophys Acta*, **1833**, 3481-98.
- Park JB, Lee CS, Jang JH, et al (2012). Phospholipase signalling networks in cancer. *Nat Rev Cancer*, **12**, 782-92.
- Pledge-Tracey A, Sobolewski MD, Davidson NE (2007). Sulforaphane induces cell type-specific apoptosis in human breast cancer cell lines. *Mol Cancer Ther*, **6**, 1013-21.
- Rizzo MA, Shome K, Vasudevan C, et al (1999). Phospholipase D and its product, phosphatidic acid, mediate agonist-dependent raf-1 translocation to the plasma membrane and the activation of the mitogen-activated protein kinase pathway. *J Biol Chem*, **274**, 1131-9.
- Roskoski R, Jr. (2014). The ErbB/HER family of protein-tyrosine kinases and cancer. *Pharmacol Res*, **79**, 34-74.
- Thompson EW, Paik S, Brunner N, et al (1992). Association of increased basement membrane invasiveness with absence of estrogen receptor and expression of vimentin in human breast cancer cell lines. *J Cell Physiol*, **150**, 534-44.
- Zheng Y, Rodrik V, Toschi A, et al (2006). Phospholipase D couples survival and migration signals in stress response of human cancer cells. *J Biol Chem*, **281**, 15862-8.
- Zhong M, Shen Y, Zheng Y, et al (2003). Phospholipase D prevents apoptosis in v-Src-transformed rat fibroblasts and MDA-MB-231 breast cancer cells. *Biochem Biophys Res Commun*, **302**, 615-9.