

## RESEARCH ARTICLE

# Analysis of Knowledge Level in Brazilian Students about Human Papillomavirus Infection and Development of Penile Cancer

George Kemil Abdalla<sup>1</sup>, Emanuella Francisco Fajardo<sup>1</sup>, Bruno Belmonte Martinelli Gomes<sup>2</sup>, Thiago Mantello Bianco<sup>3</sup>, Ana Karina Marques Salge<sup>4</sup>, Eduardo Elias Vieira de Carvalho<sup>1</sup>, Marlize Moura Dos Reis<sup>1</sup>, Dayana Pousa Siqueira Abrahão<sup>1</sup>, Douglas Reis Abdalla<sup>1\*</sup>

### Abstract

**Introduction:** Human papillomavirus (HPV), belonging to the Papovavirida family, is the most prevalent sexually transmitted disease (STD) agent worldwide. In Brazil, it is estimated that there are 3-6 million people infected with HPV. **Aim:** The aim of this study was to evaluate the knowledge of young male students about penis cancer related to HPV infection. **Methods:** This exploratory and quantitative study was conducted to analyze answers of 242 male students attending a private college located in Uberaba city, Minas Gerais state, Brazil, during 2015. **Results:** Most of the 242 participants (88.8%) affirmed having started sexual life very early, the majority (79.3%) were currently married and 69.8% had a single sexual partner. Regardless of their knowledge about HPV virus and its relationship with penis cancer, our data showed a general lack of awareness of the participants. **Conclusion:** Our results suggest that despite efforts to propagate information about HPV infection and its relation to penis cancer, the level of knowledge of students is low. Because of that, it is important to improve the information spread by media, emphasizing prevention and treatment of HPV infection in men.

**Keywords:** Papillomavirus infection- penile cancer- knowledge- university students

*Asian Pac J Cancer Prev*, **18** (5), 1371-1376

### Introduction

Human papillomavirus (HPV) belongs to the gender Papillomavirus, family Papillomaviridae and it presents circular, double-stranded and *non-enveloped DNA* (Xavier et al., 2007).

HPV is a world widely virus, which can be transmitted by direct or indirect contact with the injury carrier. HPV is mostly transmitted by sexual contact, the infection can be acquired after a single sexual relationship with an infected partner. After inoculation, the incubation time is within three weeks to eight months (Silva et al., 2013).

HPV is the most prevalent sexually transmitted disease (STD). There are 3-6 million people infected in Brazil. Around 3-5% of Brazilian people with active sexual life present HPV as a disease. (Queiroz et al., 2005).

Brazil is one of the countries with highest prevalence of HPV infection, registering around 137,000 new cases per year (Santos et al., 2011).

HPV infections are not always macroscopically visible, and they are classified in: I) Latent infection, which can be diagnosed by molecular biology tools. 'It probably occurs', e seguir com o singular daí em diante because of an exposure to the virus, with further virus penetration in the host cell, without virus replication or even without complete virus maturation. This is the most common sexual disease in Brazil. II) Subclinical, with no clinical symptoms, but with alterations that can be detected by diagnostic methods such as: peniscopy, colpocitology, colposcopy, biopsy and clinically, when there are visible lesions (Xavier et al., 2007). The virus is also being associated to other cancer types, besides the cervical, anal and penis cancers.

There is correlation between HPV and epithelium, larynges and esophagus cancers, and it is already clear the correlation between HPV and head/neck cancers. The main transmission and propagation agent is the man, indirectly contributing to the high number of cervical

<sup>1</sup>Department of Health Science, Faculty of Human Talent, Uberaba, Minas Gerais, Zip Code: 38100-000, <sup>2</sup>Department of Celular and Molecular Biology, <sup>3</sup>Department of Clinical Oncology, Stem Cells and Celular Therapy, Medicine Faculty of Ribeirão Preto, Ribeirão Preto, São Paulo, Zip code: 05508-220, <sup>4</sup>Nursing School, Federal University of Goiás, Goiânia, Goiás, Zip Code: 74605-080, Brazil. \*For Correspondence: drabdalla@facthus.edu.br

neoplasias. (Santos et al., 2011).

The main risk factor to the penis cancer is HPV and, in both genders, it is also associated to other neoplasia, causing benign lesions in the skin (warts) and in mucosal membranes (condyloma). In addition, HPV is associated to malignant lesions in the anus, vagina, oropharynx, larynx and bronchia. Esse dado de que foi encontrado no Brasil em 1986 quer dizer que foi a primeira vez no MUNDO foi no Brasil em 1986 ou no mundo já tinha acontecido antes e no Brasil foi só em 1986? De qualquer forma, acredito que esse dado ficaria melhor no parágrafo que antecede o anterior (aquele que termina com a Ref Santos et al, 2011)., recent data from research using molecular markers show that around 50% of penis carcinoma presents HPV DNA (Cubilla et al., 2010). However, data are not totally clear yet, some authors suggest that HPV detection in penis carcinoma can be correlated with 22,2-82,9% of cancer cases. These variable values might be related to problems in diagnostic methodologies or even lack of knowledge within the relation between HPV and penis cancer (Kirrander et al., 2011).

Diagnostic tracking and early treatment have presented high efficacy, being used as strategy approaches to change a prevention model based in lesions detections to biomolecular virus identification. The concept is to use a more sensitive test to detect early infections that can be severe before they achieve malignant stage (Santos et al., 2011).

Men difficulty to look for medical attendance is a crucial factor to the increase in the number of infections of infections related to HPV, the lack of knowledge causes absence of important information to a healthy life, because it avoids prevention. The vulnerability of these men group causes suffering to themselves and to their families as well and, besides, the public expenses with medical attendances and treatments may collaborate to the financial burden of public system of attention to health (Brasil, 2008).

Because of all that was exposed and because of the fast increase of infections by HPV and its close relation with penis cancer, the aim of this study was to evaluate the knowledge among male undergraduate students about penis cancer and its relation to HPV virus.

## Materials and Methods

### Study Design

This study is a research with quantitative interest, with the aim of evaluating the knowledge among undergraduate students from Faculdade de Talentos Humanos (FACTHUS), located at Uberaba/MG city, during the year of 2015. We interviewed 242 undergrads (around 3,000 students in total) belonged to diferents areas: Exacts, Humans, and Healthy. The students were divided in basic or intermediate-final groups, considering as basic the students who were between the first and second years of college, and intermediate-final the students who were between the third and fifth year.

We removed from the study all the female students and all the students that were attending to classes belonged to both groups, basic and intermediate-final.

Data was collected through a quiz with questions

about HPV, preventive exam and penis cancer, as well as questions about social/demographic aspects. This in a developed and modified way to estimate the level of knowledge about the theme.

Volunteers were well directed about the questions by professionals. The participants of the study received the quiz with the instructions and orientation, and they did not have limited time to answer it; the doubts from the participants were immediately answered by the professional who applied the quiz.

### Ethical aspects

This study was approved by the Ethics Committee In Research from the Faculty of Human Talent, through the protocol number 0058/2015. All the students accepted to participate of the study as volunteers and they agreed with it by signing the Agreement Term.

### Statistical Analyses

Continuous variables were described as median (confidence of 95%) and nominal variables were number of occurrence and percentage. Statistical analyses were done using the software SPSS 20. Logistic regression test was used to test the influence of the quiz variables to the knowledge about HPV infection and penis cancer. Statistical results were considered significant when  $p < 0.05$ .

## Results

We interviewed 242 students attending to differents courses of FACTHUS. Their age were between 17 and 49 years old (median of 23), being 79.3% single and 20.7% married. Most of the interviewed (81.4%) do not have children; most of them are attending the first to the fifth college semester (71.9%). About parents educational level, most of them have complete high school (50.0% of paternal and 51.2% maternal education level).

About social behavior, data indicates that, independently of study area, most of the students (90.5%) does not have smoking habit. By the other hand, they initiated sexual life early (88.8%), before 20 years old; 69.8% have single sexual partner, being the partner a woman in 95% of the cases. Within the students 175 use condoms, being 72.3%. All these data can be visualized at Table 1.

Data presented at Table 2 show that 166 students belong to the Exact area, 59 belong to Human and 17 to the Health area. We can see that most of them are single (Exact 84.9%, Human 67.8% and Healthy 64.7%), with  $p = 0.006$ . Most of the Exact area has been completed the basic formation of the college (80.1%), Human 52.5% and Health 58,8%, being  $p = 0.0001$ . Independently of the area, the students do not smoke ( $p = 0.008$ ), initiated sexual life early ( $p = 0.045$ ), have single partner ( $p = 0.001$ ) and use condom ( $p = 0.019$ ). We can notice that the results presented at Table 1 and Table 2 do not change after study area split.

About the penis cancer knowledge (Table 3), putting together factors as knowledge about HPV infection, prevention methods, beginning of sexual life, condom use and single sexual partner, marital status and age, the results indicate that when the person knows about HPV infection

Table 1. Data about General Population. Note: \* students from 1st up to 5th period; \*\* students from 6th to 10th period

Variable	Volunteer n = 242
Age	Median (Range) 23 (17-49)
Marital Status	n (%)
Single	192 (79.3)
Married	50 (20.7)
Formation Period	n (%)
Basic*	174 (71.9)
Intermediate-Final**	68 (28.1)
Parent's Education	
Mother	n (%)
Elementary School Incomplete	43 (17.8)
Complete primary education	39 (16.1)
Complete High School	124 (51.2)
Full Higher Education	36 (14.9)
Father	n (%)
Elementary School Incomplete	48 (19.8)
Complete primary education	43 (17.8)
Complete High School	121 (50.0)
Full Higher Education	30 (12.4)
Smokes	n (%)
Yes	9 (3.7)
No	219 (90.5)
Sometimes	14 (5.8)
Parenthood	n (%)
Yes	45 (18.6)
No	197 (81.4)
Sexual Activity	n (%)
Not started yet	10 (4.1)
Before 20 years	215 (88.8)
Before 25 years	17 (7.0)
After 25 years	0 (0.0)
Steady partner	n (%)
Yes	169 (69.8)
No	73 (30.2)
Partner's Gender	n (%)
Male	8 (3.3)
Female	230 (95.0)
Both	4 (1.7)
Use of Condom	n (%)
Yes	175 (72.3)
No	67 (27.7)

\* Basic Cycle Students between 1st and 5th; \*\* Intermediate-final Cycle Students between 6th and 10th

there is 2.13 times more chance of knowing about penis cancer ( $p=0.034$ ), and when the person is aware about prevention methods there is an increase of 4.4 times more chance of knowing about penis cancer ( $p=0.0001$ ).

Data also indicates that single people has 2.37 times

more chance of knowing what is penis cancer ( $p=0.073$ ). People who study Health area presents between 2.77-2.56 times more chance of knowing what is penis cancer than people who study Exact or Human. Besides that, older students present between 2.38-3.4 times more chance of cancer knowledge than younger ones. Another factor that collaborates to the penis cancer knowledge is the beginning of sexual life, showing that people who already initiated the sexual life have 2.35 times more chance of knowing what penis cancer is. However, the educational level (inside the college), single sexual partner, and condom use do not influence about the penis cancer knowledge.

## Discussion

HPV infection has been described as a risk factor to the development of cancers in the male genital tract, including penis cancer (Wilkin and Chiasson, 2004). This way, this study was made to show the knowledge of young undergrads about HPV infections and its relation with penis cancer.

Our study indicates that the participants were young, with an average age of 23 years old. Thinking about that, we need to take in consideration that penis cancer caused by HPV infection has been observed mostly in young men (Micali et al., 2006). Thus, the information received by people about this theme and manners of prevention may be an important tool to fight against penis cancer caused by HPV infection. Our study showed that many men are exposed to HPV infection because most of them began their sexual life very early, however, part of them affirm to use condom, which is the main way of preventing sexual diseases. Considering that HPV is a sexual transmitted disease (STD), the premature beginning of sexual life may be a risk factor to the disease development, so this study data corroborates with Costa and Goldenberg (2013) that shows the first sexual relationship happens before 20 years old. These data are reinforcing the data of Silva et al., (2013) that indicated that most of the participants were single and that their relationship with several partners could contribute to increase the risk of HPV infection. Besides, the study of Conti et al., (2006) showed that within the interviewed, more than 60% used condom during sexual activities. Other data from national and international literature, indicate an increase of condom use by young population, which is happening because of several awareness campaigns, about STDs and pregnancy prevention (Veira et al., 2004).

According to Pinto and collaborators (2011), the sociocultural, political and economical aspects are important criteria to be related to the vulnerability, such as educational level, life habits, access to health services and prevention of STDs. It is important to mention that the participants of this study were undergrad students, which allow us to think that they have access to quality information and that they have basic knowledge about diseases prevention, such as HPV infection. This way, we notice that in spite of they have access to information about penis cancer prevention, the knowledge level of the students about this theme was different among them, indicating that the students who have known about

Table 2. Population Data Separated by Training Areas. Note: \* Students from 1st to 5th period; \*\* students from the 6th to the 10th period

Variable	Volunteer Exact Courses n = 166	Volunteer Human Courses n = 59	Volunteer Health Courses n = 17	P Value
Age	Median (Range) 21 (17-48)	Median (Range) 26 (18-49)	Median (Range) 26 (18-45)	NS
Marital Status	n (%)	n (%)	n (%)	
Single	141 (84.9)	40 (67.8)	11 (64.7)	0.006
Married	25 (15.1)	19 (32.2)	6 (35.3)	
Formation Period	n (%)	n (%)	n (%)	
Basic*	133 (80.1)	31 (52.5)	10 (58.8)	0.0001
Intermediate-Final**	33 (19.9)	28 (47.5)	7 (41.2)	
Parent's Education				
Mother	n (%)	n (%)	n (%)	
Elementary School Incomplete	27 (16.3)	11 (18.6)	5 (29.4)	NS
Complete primary education	27 (16.3)	9 (15.3)	3 (17.6)	
Complete High School	90 (54.2)	26 (44.1)	8 (47.1)	
Full Higher Education	22 (13.3)	13 (22.0)	1 (5.9)	
Father	n (%)	n (%)	n (%)	
Elementary School Incomplete	32 (19.3)	9 (15.3)	7 (41.2)	NS
Complete primary education	29 (17.5)	12 (20.3)	2 (11.8)	
Complete High School	87 (52.4)	27 (45.8)	7 (41.2)	
Full Higher Education	18 (10.8)	11 (18.6)	1 (5.9)	
Smokes	n (%)	n (%)	n (%)	
Yes	2 (1.2)	5 (8.5)	2 (11.8)	0.008
No	154 (92.8)	52 (88.1)	13 (76.5)	
Sometimes	10 (6.0)	2 (3.4)	2 (11.8)	
Parenthood	n (%)	n (%)	n (%)	
Yes	22 (13.3)	19 (32.2)	4 (23.5)	0.005
No	144 (86.7)	40 (67.8)	13 (76.5)	
Sexual Activity	n (%)	n (%)	n (%)	
Not started yet	9 (5.4)	1 (1.7)	0 (0.0)	0.045
Before 20 years	9 (5.4)	4 (6.8)	4 (23.5)	
Before 25 years	148 (89.2)	54 (91.5)	13 (76.5)	
After 25 years	0 (0.0)	0 (0.0)	0 (0.0)	
Steady partner	n (%)	n (%)	n (%)	
Yes	110 (66.3)	45 (76.3)	14 (82.4)	NS
No	56 (33.7)	14 (23.7)	3 (17.6)	
Partner's Gender	n (%)	n (%)	n (%)	
Male	162 (97.6)	56 (94.6)	12 (70.6)	0.001
Female	4 (2.4)	1 (1.7)	3 (17.6)	
Both	0 (0.0)	2 (3.4)	2 (11.8)	
Use of Codon	n (%)	n (%)	n (%)	
Yes	129 (77.7)	35 (59.3)	11 (64.7)	0.019
No	37 (22.3)	24 (40.7)	6 (35.3)	

HPV already know better about penis cancer prevention. However, most of the participants reported that they do not know about HPV.

The results about knowing what HPV is and how to prevent its infection are according to the results found by Sepúlveda-Carrillo et al., (2014), who did a literature

review about sexuality, knowledge, preventional methods and vulnerability to HPV infection, and when they analysed 766 papers, they found out that at least 1% of men understand the severity of HPV infection, and this study also revealed that most of men relate HPV only to women infections. The study of Silva et al., (2013)

Table 3. Students' Knowledge about Penile Cancer Related to Sociodemographic and Behavioral Factors (Multivariate Analysis by Linear Regression). Note: \* students from 1st to 5th period; \*\* students from the 6th to the 10th period

Variable	Volunteer n = 242	Knowledge About Penile Cancer	P Value
Knowledge about HPV	n (%)	OR (95%IC)	
Yes	81 (33.5)	2.13 (1.06-4.27)	0.034
No	161 (66.5)	1	
Knowledge about HPV prevention	n (%)	OR (95%IC)	
Yes	122 (50.4)	4.41 (2.20-8.82)	0.0001
No	120 (49.6)	1	
Marital Status	n (%)	OR (95%IC)	
Single	192 (79.3)	2.37 (0.92-6.07)	0.073
Married	50 (20.7)	1	
Age	n (%)	OR (95%IC)	
Under 18 years	69 (28.5)	0.29 (0.07-1.20)	0.088
19-30 years	125 (51.7)	0.42 (0.17-1.02)	0.056
Over 30 years	48 (19.8)	1	
Formation Period	n (%)	OR (95%IC)	
Basic*	174 (71.9)	1.15 (0.55-2.41)	NS
Intermediate-Final**	68 (28.1)	1	
Formation Field	n (%)	OR (95%IC)	
Exact	166 (68.6)	0.36 (0.09-1.35)	NS
Human	59 (24.4)	0.39 (0.09-1.59)	NS
Health	17 (7.0)	1	
Sexual Activity	n (%)	OR (95%IC)	
Not started yet	10 (4.1)	1	
Before 20 years	215 (88.8)	2.22 (0.35-14.23)	NS
Before 25 years	17 (7.0)	2.58 (0.26-25.32)	NS
Steady partner	n (%)	OR (95%IC)	
Yes	169 (69.8)	1.10 (0.53-2.21)	NS
No	73 (30.2)	1	
Use of Condom	n (%)	OR (95%IC)	
Yes	175 (72.3)	1.37 (0.61-3.07)	NS
No	67 (27.7)	1	

showed that 58% of interviewed men did not have any knowledge about HPV. These data reveal that there is an important need of actions with the aim of getting to those people who is sexual active and can be spreading infection just because of lack of knowledge.

Presenting the ways of viral transmission and infection prevention should guarantee the reduction of men infection and consequently reduce the infection in their partners. Grigol and Facio-Junior (2013) showed higher incidence of HPV infection within men between 17-30 years old (61.7%), what shows the importance of awareness campaigns directed to this age group, because this group is vulnerable to infection, since they have poor knowledge and no perception about HPV infection.

Pelucchi et al., (2010) reveals the need of more health services related to identification of damages related to HPV infection, because modification on people behavior would happen only by information passed directly to target population. Besides, prevention methods such as vaccines and condoms should lead to drop in 90-70% the number of genital and cervical warts in women, and

drop around 40% the number of men with genital warts (Einstein et al., 2011).

It is worrying the level of knowledge about penis cancer and its prevention. Because of the early beginning of sexual life and the lack of knowledge about HPV infection, young people have been the most part of infected people, being the penis cancer more notable among them.

The results obtained in this study are in agreement with the results found by Reis (2010), that conclude 64.5% of undergrads do not recognize the relationship between HPV and penis cancer. These data, as ours, indicate that there is carelessness and vulnerability by young men about the HPV infection and possible complications, such as penis cancer. It is worth pointing out that the participants of our study attend to a private college and have access to different kinds of media, which shows us that prevention campaigns and information strategies are still inefficient.

Our results suggest that besides of all the effort to propagate information about HPV infection and its relation to penis cancer, the level of knowledge of the students is low. This is a worrying fact, since the participants are part

of superior education and have full access to information. Because of that, it is important to improve the information spread by media, emphasizing prevention and treatment of HPV infection in men. Besides, it is important to create strategies for health professionals to be trained to inform people, contributing to reduce the number of infections.

## Acknowledgements

The present study was supported by grants from the Faculty of Human Talent, Uberaba, Minas Gerais.

## References

- Brasil MS (2008). National policy for comprehensive health care for man. Ministry of health.
- Conti FS, Bortolin S, Kulkamp IC (2006). Education and health promotion: behavior and knowledge of adolescents from public and private schools in relation to human papillomavirus. *DST – J bras Doenças Sex Transm*, **18**, 30-5.
- Costa LA, Goldenberg P (2013). Human papillomavirus (HPV) among youth: a warning sign. *Health Risk Soc*, **22**, 249-61.
- Cubilla AL, Lloveras B, Alejo M, et al (2010). The basaloid cell is the best tissue marker for human papillomavirus in invasive penile squamous cell carcinoma: a study of 202 cases from Paraguay. *Am J Surg Pathol*, **34**, 104-14.
- Einstein MH, Baron M, Levin MJ, et al (2011). Comparative immunogenicity and safety of human papillomavirus (HPV)-16/18 vaccine and HPV-6/11/16/18 vaccine. Follow-up from months 12-24 in a Phase III. Randomized study of healthy women aged 18-45 years. *Hum Vaccine*, **7**, 1343-58.
- Grigol FN, Facio Junior N (2013). Characterization of men with condylomathosis at a University Hospital. *Health Sci Arch*, **20**, 118-22.
- Kirrandar P, Kolaric A, Helenius G, et al (2011). Human papillomavirus prevalence, distribution and correlation to histopathological parameters in a large Swedish cohort of men with penile carcinoma. *BJU Int*, **108**, 355-9.
- Micali G, Nasca MR, Innocenzi D, Schwartz RA (2006). Penile Cancer. *JAAD*, **54**, 369-91.
- Pelucchi C, Esposito S, Galeone C, et al (2010). Knowledge of human papillomavirus infection and its prevention among adolescents and parents in the greater Milan area, Northern Italy. *BMC Public Health*, **10**, 1-12.
- Pinto DS, Fuzii HT, Quaresma JAS (2011). Prevalence of genital HPV infection in urban and rural women in the Eastern Brazilian Amazon. *Public Health Note*, **27**, 769-78.
- Reis AAS (2010). Human papillomavirus and public health: cervical cancer prevention. *Cien Saude Colet*, **15**, 1055-60.
- Santos IM, Maioral MF, Hass P (2001). HPV infection in men: Importance in transmission, treatment and prevention of the virus. *Est Biol*, **32**, 111-18.
- Sepúlveda-Carrillo GJ, Goldenberg P (2014). Knowledge and Practices of young people related to human papillomavirus infection – new updated. *Rev Colomb Obstet Ginecol*, **65**, 152-61.
- Silva ACN, Sousa GB, Lacerda GAF (2013). Self-perception of university men in relation to human papillomavirus: a problem to be considered in public health. *Interdisciplinary. Eletr J Univar*, **2**, 71 –7.
- Vieira M, Guimarães BEM, Barbosa MA, et al (2004). Factors associated with condom use by female adolescents in Goiânia City. *DST – J bras Doenças Sex Transm*, **16**, 77-83.
- Queiroz DT, Pessoa SMF, Sousa RA (2005). Infection by human papilomavirus (HPV): doubts and challenges. *Acta Paul Enferm*, **18**, 190-6.
- Xavier SD, Filho IB, Carvalho JM, Framil VMS, Castro TMPPG (2007). Frequency of appearance of human papillomavirus (HPV) in oral Mucosa of men with anogenital HPV by a molecular technique. *Arq Int Otolaryngologist*, **11**, 36-44.
- Wilkin T, Chiasson MA (2004) Sexually transmitted infections in men and women. in 'principles of gender-specific medicine, Legato MJ. Elsevier, pp 966-77.