

RESEARCH ARTICLE

Prevalence of Human Papillomavirus subtypes 16 and 18 among Yemeni Patients with Cervical Cancer

Hussain Gadelkarim Ahmed^{1,2*}, Saleh Hussein Bensumaidea³, Fawaz D Alshammari⁴, Fahaad Saleh H Alenazi¹, Bassam Ahmed ALmutlaq¹, Majid Zannon Alturkstani⁵, Ihdaa Ali Aladani⁵

Abstract

Background: The Human Papillomavirus (HPV) is a DNA tumor virus that causes epithelial proliferation. There are more than 100 HPV subtypes, of which 13 subtypes are regarded as high risk subtypes that can cause cancers of epithelial mucosal surfaces. High risk human papilloma viruses (HR-HPV) subtypes 16 and 18 plays a major role in the etiology of cervical cancer worldwide. Therefore, the aim of this study was to screen for the existence of HPV16 and HPV18 among Yemeni women with cervical lesions. **Methodology:** Formalin fixed paraffin wax processed tissue blocks were retrieved for 200 patients (150 were previously diagnosed with cervical cancer and the remaining 50 were diagnosed with different benign conditions). **Results:** Of the 200 cervical cancer tissue specimens, HR-HPV 16 was identified in 74/200 (37%) samples and couldn't be recognized in 126/200(63%) tissue samples. HR-HPV 18 was identified in 32/200 (16%) specimens and couldn't be recognized in 168/200(84%) tissue specimens. **Conclusion:** HR-HPV subtypes were prevalent among Yemeni women with cervical cancer, with significant increase of HR-HPV subtype 16 over the HR-HPV subtype 18.

Keywords: HPV16- HPV18- Yemeni- cervical cancer- prevalence

Asian Pac J Cancer Prev, **18** (6), 1543-1548

Introduction

Human papillomavirus (HPV) is one of the most common sexually transmitted diseases worldwide, which is strongly involved in the pathogenesis of genital cancers such as cervical cancer (Zandberg et al., 2013). The majority of individuals who involve in multi-sexual activity becoming infected at some stages of their lifetime (Baseman and Koutsky, 2005). More than 130 HPV types have been well-known and categorized into low- or high-risk groups according to their potential for oncogenesis depending on persistent infection (Zur Hausen, 2009). According to the International Agency for Research on Cancer (IARC), the HPV types 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59 and 66, regarded as high risk Human Papillomaviruses (HR-HPV). The HPV types 6, 11, 40, 42, 43, 44, -53, 54, 61, 72, and 81, regarded as low-risk Human Papillomaviruses (LR-HPV) (Steben and Duarte-Franco, 2007). HPV genome encodes for early structural genes (E1 to E8) and late structural genes (L1 and L2). The late-coding regions yield structural proteins whereas the early-coding regions, mainly E6 and E7, are responsible for malignant transformation

(Psyri et al., 2011). E6 and E7 proteins of HR-HPVs have transforming activity that leads to autonomous or synergistic immortalization of the cell whereas; low-risk LR-HPVs have weak immortalization activity. This difference is thought to correlate with the ability of E7 proteins to make the degradation of the Rb gene rather than just their affinity for the Rb gene. HR-HPV E6 and E7 proteins can also induce mitotic abnormalities through mitotic spindle checkpoints, whereas, this is not realized in LR-HPV (Jemal et al., 2011).

A strong association has previously been built for the responsibility of HPV in oncogenesis of cervical cancer, and it is involved in other genital cancers (Parkin and Bray, 2006). Moreover, the influence of HPV is more far-reaching than initially supposed, with a strong link between infection and the development of other cancers such as, oral, pharynx and esophageal cancers (Zandberg et al., 2013).

Cervical cancer is the second greatest females' cancer in developing countries with an estimated 445,000 new cases in 2012 (84% of the new cases world wide). Approximately 270,000 women died from cervical cancer in 2012; more than 85% of these deaths cases occurred

¹Department of Pathology College of Medicine, ⁴Department of Clinical Laboratory, College of Applied Medical Science, University of Hail, ⁵Batterjee Medical College, Jeddah, Kingdom of Saudi Arabia (KSA), ²Department of Histopathology and Cytology, FMLS, University of Khartoum, Sudan, ³Department of Pathology, College of Medicine, Hadhramout University of Science and Technology, Yemen. *For Correspondence: hussaingad1972@yahoo.com

in less developed countries. HPV types (16 and 18) are known to be responsible of 70% of cervical cancers and precancerous cervical lesions (WHO, 2015). HR-HPV types 16 and 18 are frequently associated with invasive cervical cancer than other types. This indicates that HR-HPV types 16 and 18 are more carcinogenic than other HPV types. Moreover, HR-HPV type 16-related cancers occur at a younger age than other cervical cancers (Pinkowish, 2009).

However, there is relatively complete paucity of data from Yemen regarding the association between HPV and cervical cancer. The few previous reports in this context have investigated HPV types other than HR-HPV types 16 and 18. Therefore, the aim of this study was to find out the prevalence of HR-HPV types 16 and 18, which is essential for any future strategies in term of control, prevention and vaccination.

Materials and Methods

In the present study 200 formalin fixed paraffin embedded tissue (PET) blocks were obtained from 150 patients previously diagnosed with cervical cancer and the remaining 50 with benign cervical lesions, their ages ranging from 21 to 75 years with a mean age of 47 years. The sample size represents a full coverage of patients whom their samples were referred to histopathology laboratories which provide diagnostic services for 6 provinces. All samples were confirmed by histopathologic diagnosis regardless to age and histological type. Samples were sequentially recruited from 6 Yemeni provinces (Aden, Azaal, Al-Janal, Tohama, Hadhramout and Saba'a) between March 2013 and July 2014. Specimens of patients with a history of neo-adjuvant treatments or another cancer or cancer that metastasized from other organs were excluded. Personal and demographic data as well as, clinico-pathological characteristics were obtained from each patient's file.

DNA Extraction

Three cuts of 3-µm-thickness sections of PET were cut and put into 1.5 ml micro-tubes. Positive controls (HPV 16 infected SiHa cells and HPV 18 infected HeLa cells embedded in paraffin wax) and negative controls (blank paraffin wax block) were also used for quality control. Total DNA from PET was extracted from each specimen using DNA extraction kit purchased from Sacace

biotechnologies-Casera –Italy. The 268 bp fragment of the β-globin gene amplification was used to assess the quality of DNA in PETs. The primers were GH20 (5' GAA GAG CCA AGG ACA GGT AC3') and PC04 (5'-CAA CTT CAT CCA CGT TCA CC-3'). Positive specimens for β-globin gene were used and indicated that the samples were accessible for the subsequent analysis.

Polymerase chain reaction (PCR) for detection of HPV

PCR was applied to amplify the HPV E6/E7 gene of HPV 16 and 18 using type-specific primers (type-specific PCR, TS PCR). Below were primer sequences (Wang, et al., 2008) : HPV16 E6: forward 5'-CTG CAA GCA ACA GTT ACT GCG ACG-3', reverse 5'-CAT ACA TCG ACC GGT CCA CC-3', product of 315 bp; HPV 18 E7: forward 5'-GAG CCG AAC CAC AAC GTC AC-3', reverse 5'-GGA TGC ACA CCA CGG ACA CA-3', product of 152 bp.

Gel-electrophoresis

The PCR products were visualized in 2% Agarose gel with 0.5 µg/ml Ethidium bromide. Ten micro liters of 100bp DNA ladder and PCR product was loaded on the gel. Gel-electrophoresis was made at 120V and 36 mA for 60 minutes. Images were reserved by Gel documentation system (Gel mega, digital camera and software in a computer).

Statistical Analyses

Data management was done by using the Statistical Package for Social Sciences (SPSS version 16; SPSS Inc, Chicago, IL). SPSS was applied for analysis and to perform Fisher exact test for statistical significance (P value < 0.05 was considered significant). The 95% confidence level and confidence intervals were used.

Results

Of the 200 cervical cancer tissue specimens, HR-HPV 16 was identified in 74/200 (37%) samples and couldn't be recognized in 126/200(63%) tissue samples. Out of the 74 infected specimens, 72/74 (97.3%) were found among cervical cancer's patients and 2/74 (2.7%) were found among those with benign cervical lesions. The prevalence of HR-HPV type 16 was 72/150 (48%). With 95% confidence level, the risk associated with relationship between HR-HPV subtypes 16 and 18 and malignant




Figure 1. Description of the Study Population by Positive HR-HPV Subtypes 16 and 18

Table 1 Distribution of the Study Population by Age and HR-HPV Subtypes 16 and 18 Infection

	<35years	36-45	46-55	56+	Total
HR-HPV 16					
Positive	18	27	24	5	74
Negative	13	33	48	32	126
Total	31	60	72	37	200
HR-HPV 18					
Positive	6	9	13	4	32
Negative	25	51	59	33	168
Total	31	60	72	37	200




Figure 2. Description of Age by Positive HR-HPV Subtypes 16 and 18




Figure 3. Description of Positive HR-HPV Subtypes 16 and 18 by Residence

transformation was statistically significant, $P < 0.001$, as indicated in Figure 1.

HR-HPV 18 was identified in 32/200 (16%) specimens and couldn't be recognized in 168/200 (84%) tissue specimens. Out of the 32 infected specimens, 31/32 (96.9%) were found among cervical cancer's patients and 1/32 (3.1%) was identified among those with benign cervical lesions. Moreover, co-infections of both types were identified among 10 patients, representing 10/74 (13.5%) of HR-HPV 16 and 10/32 (31.3%) of HR-HPV 18. The prevalence of HR-HPV type 18 was 31/150 (20.7%).

As shown in Table 1, for HR-HPV subtype 16, the highest frequency of infection rates were identified among age range 36–45 years representing 27/74 (36.5%) followed by age range 46–55, <36 and 56+ years, constituting 24/74 (32.4%), 18/74 (24.3%) and 5/74 (6.8%), respectively. For HR-HPV subtype 18, the highest




Figure 4. Description of Percentage Positive HR-HPV Subtypes 16 and 18 within Entire Residence

Table 2. Distribution of the Study Population by HR-HPV Subtypes 16 and 18 by Diagnosis

HR-HPV	Diagnosis						
HR-HPV16	Benign	CIN1	CIN2	CIN3	SCC	ADC	Others
Positive	2	2	5	11	46	5	1
Negative	48	13	8	5	34	14	3
Total	50	15	13	16	80	19	5
HR-HPV18							
Positive	1	1	3	4	9	12	2
Negative	49	14	10	12	71	7	3
Total	50	15	13	16	80	19	5

frequency of infection rates were identified among age range 46–55 years representing 13/32 (40.6%) followed by age range 36–45, <36 and 56+ years, constituting 9/32 (28%), 6/32 (18.8%) and 4/32 (12.6%), in this order.

On the other hand, when computing the percentage in each group, for HR-HPV subtype 16, the extreme percentage of infection was found in age group <35 years constituting 58% followed by 36–45, 46–55, and 56+ years, demonstrating 45%, 33%, and 14%, respectively. For HR-HPV subtype 18, the great percentage of infection was identified in age group <35 years constituting 19% followed by 46–55, 36–45, and 56+ years, establishing 18%, 15%, and 11%, as indicated in Figure 2.

In respect to the residence, the great majority of HR-HPV subtype 16 positive cases were coming from Aden representing 22/74 (29.7%) followed by Al-Janad, Azaal, and Tohama, constituting 18/74 (24.3%), 14/74 (19%) and 10/74 (13.5%), respectively. For HR-HPV subtype 18, most positive cases were from Al-Janad representing 10/32 (31.3%) followed by Azaal representing 9/32 (28%), as shown in Figure 3.

Nevertheless, when calculating the percentage within each entire residence, for HR-HPV subtype 16, the highest percentage of infection was found in the Aden representing 49% followed by Tohama, both (Azaal and Al-Janad) and Saba'a constituting 42%, 35%, and 33%, in this order. For HR-HPV subtype 18, the great proportion was identified in Azaal representing 22.5% followed by Al-Janad, Tohama and both (Hadhramout and Aden) constituting 19%, 12.5% and 9%, as indicated in Figure 4.

Regarding the relationship between infection with HR-HPV 16 and 18 and diagnosis, most positive cases of HR-HPV 16 were associated with Squamous Cell




Figure 5. Description of Percentage Positive HR-HPV Subtypes 16 and 18 within Entire of Diagnosis

Carcinoma (SCC) constituting 46/74 (62.2%) followed by Cervical Intraepithelial Neoplasia grade 3 (CIN3), CIN2 and Adenocarcinoma (ADC), representing 11/74 (14.9%), 5/74(6.8%), and 5/74(6.8%), respectively. For HR-HPV18, positive cases were linked to ADC, representing 12/32 (37.5%) followed by SCC, CIN3 and CIN2, constituting 9/32 (28%), 4/32 (12.5%) and 3/32(9.4%), in this order, as shown in Table 2.

Nevertheless, in calculating the percentage of the positive cases within each diagnostic group, we found that for HR-HPV16, the highest proportion of infection was identified among CIN3 patients constituted 68.8%, followed by SCC, CIN2 and ADC, representing, 57.5%, 38.5% and 26.3%, respectively. For HR-HPV16, the highest percentage of infection was found in ADC constituted 63.2%, followed by others, CIN3, and CIN2, representing, 40%, 25% and 23%, respectively, as indicated in Figure 5.

Discussion

Cervical cancer is the fourth most common cancer that affects women worldwide (Ferlay et al., 2013). It is believed that more than 95% of cases of cervical cancers are associated with a history of persistent infection by HR-HPV (Walboomers et al., 1999; Elsbali et al., 2012). Striking variations in the incidence of cervical cancer exist between developed and developing countries. Even within developed or developing world, HPV infection rates differ significantly between geographic regions and population clusters.

In the present study we investigated HR-HPV types 16 and 18, since they were the most common cervical cancer associated types, and are believed to be responsible for over 70% of cases of cervical cancer (Koutsky, 1997). This is one of few studies of its kind, which highlights the HR-HPV types 16 and 18 prevalence in cervical cancers in Yemen. There are some studies investigated HPV from Yemen, but devoted to subtypes other than HR-HPV types 16 and 18. In a recent study from Yemen that investigated 84 patients with cervical cancer for the presence of HPV subtypes 16 and 18 applying both immunohistochemistry and molecular techniques; they found that 73.8% were infected with HPV subtypes 16 and 27.4% were infected with HPV subtypes 18 (Muñoz et al., 2004). This study shows relatively inconsistent higher prevalence rates from our findings. Another study from Yemen, found prevalence rates of HPV subtypes 52,56,58,59, and 66, among cases were 0.6%, 0%, 4%, 3.3% and 0% respectively (Bensumaidea et al., 2014a). Another study investigated the presence of HPVs subtypes 31, 33, 35, 39 and 45 among Yemeni patients with cervical cancer; the study identified HPV 31, HPV 33, HPV35, HPV 39 and HPV45 in 10/150 (6.7 %), 6/150 (4 %), 6/150 (4 %), 5/150 (3.3 %) and 10/150 (6.7 %), respectively. The prevalence of these HPV subtypes among Yemeni cervical cancer patients was 24 % (Bensumaidea et al., 2014b).

The findings of this study showed prevalence rates of 48% for HR-HPV type 16 and 20.7% for HR-HPV type 18. These findings show little variation from that reported by the International Association for Research in

Cancer (IARC), in which they report prevalence rates of 53% for HPV subtype 16 and 15% for HPV subtype18 (Ahmed et al., 2015). Studies on the prevalence of HPV genotypes in cervical cancer consistently indicate that infection with HR-HPV type 16 is the most common in a variety of studies from diverse regions (Muñoz, 2000; Zhi et al.,2015; Wang et al., 2015). Then followed by infection with other HR-HPV types including HR-HPV subtype 18 (Siddiqa et al., 2014). In general HPV infection rates differ significantly between geographic regions and population clusters, (Clifford et al., 2005; Koutsky, 1997). Particularly, in the Middle East countries (Elsbali et al., 2012). However, assessment of HR-HPVs in the pathogenesis of cervical cancer and other cancers in Yemen is essential in order to evaluate the status for future plan of prevention strategies including vaccination against HPVs. In this study, we present an additional evidence of the existing epidemiological evidences regarding the presence of HPV in cervical cancers in Yemen and the potential need for vaccination against HR-HPV infections, particularly HR-HPV 16 and 18 and its effect on human health in this conflicting country. Co-infections of both HR-HPV 16 & 18 types were identified in 13.5% of the patients. This percentage is much lower than reports from highly HPV endemic areas which ranging from 30% to 65% (Al Moustafa et al., 2014; Torres-Ibarra et al., 2014).

In regard to the relationship between age and HR-HPV type 16, the highest percentage was found in age range 36–45 years (36.5%), followed by age ranges 46–55 (32.4%) and <36 years (24.3%). Many studies showing that HPV16 positivity was significantly associated with younger age, particularly when early diagnosis is employed, such as in diagnosis of CIN. This increasing infection will decrease with increasing of age (Lebelo et al., 2015). In study investigated 317 women with CIN3, HPV subtype16 was recognized in 70% of those with age range of 16-25, 59% of 26-35, and 48% of >36-year-olds ($P < 0.025$). This link acquired the form of a tendency with decreasing HPV type 16 prevalence with increasing of age ($P < 0.008$). This means that HPV type16 is commoner in younger women with high-grade cervical lesions. This report seems to show some sorts of discrepancies with our findings, which might be attributed to the large number of cervical cancers (69.3%) over the precancerous lesions (30.7%) in our series. A recent study has shown that HPV16 is responsible of absolute risk of \geq CIN3 equally in women aged 25-29 and \geq 30 years (14.2% and 15.1%, respectively) followed by HPV31 (8.0% and 7.9%), HPV52 (6.7% and 4.4%) and HPV18 (2.7% and 9.0%). The positivity increased significantly with disease development for HR-HPV16 and HR-HPV18 which were accountable for 45.6% and 8.4% of \geq CIN3, respectively. Notable, HPV 18 was responsible for 50% of adenocarcinoma in situ and 50% of invasive cancer cases (Baandrup et al., 2012). This also explains our results, since they were dedicated to older population. About 84.5% of the studied patients in the present study were older than the age of 35 year-olds.

In regard to the relationship between age and HR-HPV type 18, the highest percentage was found in age range 46–55 years followed by age ranges 36-

45 and <36, representing, 40.6%, 28%, and 18.8%, respectively. Although, HR-HPV 18 shows relatively similar association in regard to the age, but some studies agree with our findings that it tends to occur at relatively higher age (Monsonogo et al., 2015).

Furthermore, in the present study, HR-HPV16 was frequently identified in SCC and CIN. These findings were in consistency with several studies (Clifford et al., 2005; Newall et al., 2008; Clifford et al., 2003). On the other hand, most of cases of HR-HPV18 infection were found to be related to adenocarcinoma. Several studies have shown that HPV18 is more strongly linked with ADC than SCC (Ciapponi et al., 2011, Vandenbroucke et al., 2013; Chen et al., 2015).

In regard to the relationship between HR-HPV 16 &18 and residence, there is no specific justification for variations in the proportions of the infections, since all social practices and believes are relatively common all over the 6 provinces.

The limitations in this study include non-representativeness of the sample for all Yemen provinces; exclusion of the other High risk and low risk HPV types, although these HPV types were reported elsewhere.

Further studies on the distribution of all HPV subtypes in Yemeni women of all Yemeni provinces are still required.

In conclusions, the rate of infection with HR-HPV 16 and 18 is relatively higher in Yemen with predominance of HR-HPV type 16. This study may provide valuable data for future overall management including; prevention, treatment of HPV infection, cervical cancer and endorse the urgent demand for HPV vaccines.

Ethical consent

This study was approved by concerned laboratories in Yemen and by the ethics board of the Faculty of Medical Laboratory Science, Sudan University for Science and Technology (where the study was carried out). All cervical tissues were obtained as a part of the specimens required for diagnosis.

Competing interests

The authors declare that they have no competing interests.

Authors' contribution

- HGA and FDA: Contributed to the design of the study, data analyses, interpretation of results, and draft and approval of manuscript.

- SHB, FSA, BAM, MZA and IAA: Data Collection and Molecular identification, and draft and approval of manuscript.

“All authors read and approved the final manuscript.”

Acknowledgments

The authors would like to thank people at histopathology Laboratories at all 6 mentioned Yemeni provinces, for helping in the sample collection.

References

- Ahmed HG, Bensumaideia SH, Ashankyty IM (2015). Frequency of Human Papilloma virus (HPV) subtypes 31, 33, 35, 39 and 45 among Yemeni women with cervical cancer. *Infect Agent Cancer*, **10**, 29.
- Al Moustafa A, Al-Awadhi R, Missaoui N, et al (2014). Human papillomaviruses-related cancers Presence and prevention strategies in the Middle East and North African Regions. *Hum Vaccin Immunother*, **10**, 1812–21.
- Baandrup L, Munk C, Andersen KK, et al (2012). HPV16 is associated with younger age in women with cervical intraepithelial neoplasia grade 2 and 3. *Gynecol Oncol*, **124**, 281-5.
- Baseman JG, Koutsky LA (2005). The epidemiology of human papillomavirus infections. *J Clin Virol*, **32**, 16-24.
- Bensumaideia SH, Ahmed HG, Bafakeer SS, Abd-al-Aziz MS, Alshammari FD (2014)a. A correlation of immunohistochemical and molecular detection of human Papilloma Virus subtypes 16 and 18. *IJSR*, **3**, 2487-90.
- Bensumaideia SH, Bafakeer SS, Abd El Aziz MS, Alshammari FD, Ahmed HG (2014)b. Prevalence of Human Papilloma virus subtypes 52, 56, 58, 59 and 66 among Yemeni patients with cervical cancer. *Egypt Acad J Biolog Sci*, **6**, 21-6.
- Chen AA, Gheit T, Franceschi S, et al (2015). Human Papillomavirus type 18 genetic variation and cervical cancer risk worldwide. *J Virol*, **12**, JVI.01747-15.
- Ciapponi A, Bardach A, Glujovsky D, Gibbons L, Picconi MA (2011). Type-specific HPV prevalence in cervical cancer and high-grade lesions in Latin America and the Caribbean: systematic review and meta-analysis. *PLoS One*, **6**, e25493.
- Clifford GM, Rana RK, Franceschi S (2005). Human papillomavirus genotype distribution in low-grade cervical lesions: comparison by geographic region and with cervical cancer. *Cancer Epidemiol Biomarkers Prev*, **14**, 1157-64.
- Clifford GM, Smith JS, Aguado T, Franceschi S (2003). Comparison of HPV type distribution in high-grade cervical lesions and cervical cancer: a meta-analysis. *Br J Cancer*, **89**, 101-5.
- Elasbali AM, Saad El Din AH, Abdallah RAH, Ahmed HG (2012). Cervical and oral screening for HR-HPV types 16 and 18 among Sudanese women cervical lesions. *Infect Agent Cancer*, **7**, 17.
- Ferlay J, Soerjomataram I, Ervik M, et al (2013). Globocan 2012 v1.0, Cancer incidence and mortality worldwide. IARC Cancer Base No.11, 2013, Available from: <http://globocan.iarc.fr>.
- IARC (2006). Monographs on the evaluation of carcinogenic risks to humans, volume 90, human papillomaviruses. Lyon: IARC; 2006. Available at: <http://www.who.int/bulletin/volumes/85/9/06-038414/en/>.
- Jemal A, Bray F, Center MM (2011). Global cancer statistics. *CA Cancer J Clin*, **61**, 69-90.
- Koutsky L (1997). Epidemiology of genital human papillomavirus infection. *Am J Med*, **102**, 3-8.
- Lebelo RL, Bogers JJ, Thys S, et al (2015). Detection, genotyping and quantitation of multiple hpv infections in south african women with cervical squamous cell carcinoma. *J Med Virol*, **87**, 1594-600.
- Monsonogo J, Cox JT, Behrens C, et al (2015). Prevalence of high-risk human papilloma virus genotypes and associated risk of cervical precancerous lesions in a large U.S. screening population: data from the ATHENA trial. *Gynecol Oncol*, **137**, 47-54.
- Muñoz N (2000). Human papillomavirus and cancer: The epidemiological evidence. *J Clin Virol*, **19**, 1–5.
- Muñoz N, Bosch FX, Castellsagué X, et al (2004). Against which

- human papillomavirus types shall we vaccinate and screen? The international perspective. *Int J Cancer*, **111**, 278-85.
- Newall AT, Brotherton JM, Quinn HE, et al (2008). Population seroprevalence of human papillomavirus types 6, 11, 16, and 18 in men, women, and children in Australia. *Clin Infect Dis*, **46**, 1647-55.
- Parkin DM, Bray F (2006). Chapter 2: The burden of HPV-related cancers. *Vaccine*, **24**, S3/11-S3/25.
- Pinkowish MD (2009). Human Papillomavirus genotype distributions inform screening and vaccination policy. *CA Cancer J Clin*, **59**, 280-1.
- Psyrris A, Boutati E, Karageorgopoulou S (2011). Human papillomavirus in head and neck cancers: biology, prognosis, hope of treatment, and vaccines. *Anticancer Drugs*, **22**, 586-90.
- Siddiqi A, Zainab M, Qadri I, Bhatti MF, Parish JL (2014). Prevalence and genotyping of high risk Human Papillomavirus in cervical cancer samples from Punjab, Pakistan. *Viruses*, **6**, 2762-77.
- Steben M, Duarte-Franco E (2007). Human papillomavirus infection: epidemiology and pathophysiology. *Gynecol Oncol*, **107**, 2-5.
- Torres-Ibarra L, Conde-Glez CJ, Salmerón J, et al (2014). Risk factors for anal HPV-16/18 infection in Mexican HIV-infected men who have sex with men. *Prev Med*, **69**, 157-64.
- Vandenbroucke L, Robert AL, Lavoué V, et al (2013). Adenocarcinoma of the uterine cervix: particularities in diagnosis and treatment. *J Gynecol Obstet Biol Reprod (Paris)*, **42**, 207-16.
- Walboomers JM, Jacobs MV, Manos MM, et al (1999). Human papillomavirus is a necessary cause of invasive cervical cancer worldwide. *Pathol*, **189**, 12-9.
- Wang L, Wu B, Li J, Chen L (2015). Prevalence of human papillomavirus and its genotype among 1336 invasive cervical cancer patients in Hunan province, central south China. *J Med Virol*, **87**, 516-21.
- Wang Y, Wang A, Jiang R, et al (2008). Human papillomavirus type 16 and 18 infection is associated with lung cancer patients from the central part of China. *Oncol Rep*, **20**, 333-9.
- WHO (2015). Human papillomavirus (HPV) and cervical cancer 2015. Available at: <http://www.who.int/mediacentre/factsheets/fs380/en/>.
- Zandberg DP, Bhargava R, Badin S, Cullen KJ (2013). The role of human papillomavirus in nongenital cancers. *CA Cancer J Clin*, **63**, 57-81.
- Zhi YF, Cha XX, Li XF, Qiu C, Rong SH (2015). Prevalence and genotype distribution of human papillomavirus in women in the Henan Province. *Genet Mol Res*, **14**, 5452-61.
- Zur Hausen H (2009). Papillomaviruses in the causation of human cancers-a brief historical account. *Virol J*, **384**, 260-5.