

RESEARCH COMMUNICATION

Using Salivary Cotinine to Validate Self-Reports of Tobacco Use by Indian Youth Living in Low-Income Neighborhoods

Poonam Dhavan¹, Shalini Bassi^{2*}, Melissa H Stigler¹, Monika Arora^{2,3}, Vinay K Gupta², Lakshmy Ramakrishnan⁴, Cheryl L Perry¹, K Srinath Reddy^{2,3}

Abstract

Background: Self-reported tobacco use among young people can underestimate the actual prevalence of tobacco use. Biochemical validation of self-reports is particularly recommended for intervention studies where cessation outcomes are to be measured. Literature on biochemical validation of self-reports of multiple forms of tobacco use in India is sparse, particularly among young people. **Methods:** The study was conducted during the baseline household survey of a community-based tobacco prevention and cessation intervention trial for youth (10-19 years old) residing in slum communities in Delhi, India in 2009. Salivary cotinine measurement on 1,224 samples showed that youth were under-reporting use of chewing and smoking tobacco. **Results:** Self-reports had a low sensitivity (36.3%) and a positive predictive value of 72.6%. No statistically significant difference in under-reporting was found between youth in the control and intervention conditions of the trial, which will be taken into consideration in assessing intervention outcomes at a later time point. **Conclusion:** Biochemical validation of self-reported tobacco use should be considered during prevention and cessation studies among youth living in low-income settings in developing countries like India. **Impact:** The future results of biochemical validation from Project ACTIVITY (Advancing Cessation of Tobacco In Vulnerable Indian Tobacco consuming Youth) will be useful to design validation studies in resource-poor settings.

Keywords: Tobacco use - youth - underestimation - saliva - cotinine - India

Asian Pacific J Cancer Prev, 12, 2551-2554

Introduction

Most estimates of youth tobacco use are based on self-reports, either in schools or households (Reddy et al., 2006; Warren et al., 2006). Although generally considered reliable, self-reported estimates of drug use can be less sensitive and reliable among adolescents as compared to adults (Caraballo et al., 2004). This is particularly true for socially undesirable behaviors or when there are laws banning certain behaviors (Gorber et al., 2009). Therefore, several studies, especially in developed countries like the United States, corroborate self-reported tobacco use using actual biochemical validation or a bogus pipeline procedure (Adams et al., 2008). Biochemical validation is particularly recommended in intervention studies, where cessation outcomes need to be assessed and for special population groups like children and pregnant women (Benowitz et al., 2002).

Of the available markers for validation of tobacco use, a cotinine measurement is most commonly used in population studies. Cotinine is a metabolite of nicotine with a long half-life (15-19 hours) that can be

measured in body fluids such as saliva, urine and plasma. Measurement of salivary cotinine is commonly used in population studies, given the ease of sample collection and minimal discomfort to study subjects (Iqbal et al., 2007). Biochemically assessed cotinine levels can be an indicator of active smoking, smokeless tobacco use, exposure to secondhand smoke or use of pharmaceutical nicotine (Post et al., 2005; Jarvis et al., 2008).

In developing countries like India, young people are particularly vulnerable to initiation and addiction to tobacco products (Reddy and Gupta, 2004; Reddy et al., 2006; Stigler et al., 2007; Sinha et al., 2008). Studies have shown that between 4 to 75 % of 13-15 year old students in India report lifetime use of some tobacco product (Reddy and Gupta, 2004). It is also known that persons from lower socioeconomic strata are more likely use tobacco (Subramanian, 2004; Mishra et al., 2005; Mathur et al., 2008). With this background, an intervention trial, Project ACTIVITY (Advancing Cessation of Tobacco In Vulnerable Indian Tobacco consuming Youth) funded by the National Institutes for Health (NIH), was started among disadvantaged youth living in slums in Delhi, India

¹Michael & Susan Dell Center for Healthy Living, University of Texas, School of Public Health, Austin, Texas, USA, ²Health Related Information Dissemination Amongst Youth (HRIDAY), ³Public Health Foundation of India, ⁴All India Institute of Medical Sciences, Ansari Nagar, New Delhi, India *For correspondence: shalini@hriday-shan.org

in 2009 (Arora et al., 2010). The objective of this trial is to test the efficacy of a comprehensive community-based intervention for prevention and cessation of tobacco use among adolescents and young adults (10-19 years old) (Arora et al., 2010). A baseline household survey, including salivary cotinine testing on a sub-sample, was conducted to measure the prevalence of tobacco use in intervention and control communities, before the start of the intervention. The aims of the salivary cotinine testing were (i) to compare estimates of tobacco use prevalence based on self-reports vs. biochemical testing, and (ii) to assess whether misreporting of tobacco use by self-reports, if any, differed by trial condition. Given the study location and sample enrolled in this study, this paper extends the current body of research on biochemical validation of self-reported tobacco use to developing-country settings where tobacco use is on the rise, particularly among youth.

Materials and Methods

Ethical clearances were obtained for Project ACTIVITY from the Independent Ethics Committee in India and the Institutional Review Board at the University of Texas Health Science Center at Houston.

Participants

A total of 4191 households with 8702 youth between 10 to 19 years were randomly selected for baseline household surveys in seven control and seven intervention slum communities in Delhi. Two communities were randomly selected from the intervention and control conditions for biochemical validation. Given budgetary constraints, saliva was only collected from a convenience sample of 1250 youth from four communities (2 intervention and 2 control) for cotinine measurement.

Data collection

Informed consent for interview and salivary sample collection was obtained from the youth and also parents (for 10-17 year old youth). Once the study team reached a sample size of 1250 across all four communities, interviewers stopped requesting for further salivary samples. Confidential subject identification numbers were used on both the interviews and the saliva sample tubes. Self-reported use of smoking tobacco (cigarettes and bidis) and chewing tobacco (Gutkha, Kahini, Zarda etc.) was assessed in the interviews. Current use was defined as use in the past month by asking, "During the past 30 days, did you smoke cigarettes/bidis/chew tobacco?" A dichotomous tobacco use variable was created for each product and assigned a '0' for those who reported never using these products and '1' for those who did. A composite variable for 'any tobacco use' was created with a '0' for those who did not report using any tobacco products and '1' for those who used one or more.

Saliva samples were collected using the kits provided by Salimetrics (Salimetrics, 2009) which contained a piece of chewable cotton stick in a sterile tube. Stimulated saliva samples were collected by asking the subject to chew the cotton stick for two minutes and then placing the cotton back in a sterile tube. Samples were transported to the

laboratory within ten hours of collection. Saliva sample collection took approximately 2-3 minutes per study subject and was done at the end of the interview, which lasted about 30 minutes.

Data analysis

Salivary cotinine was biochemically analyzed using Enzyme-Linked Immuno Sorbent Assay (ELISA) kit from Salimetrics, USA (Dhar, 2004; Salimetrics, 2006). All statistical analyses were conducted using SAS (version 9.1). Cotinine value of 15 ng/ml was used as the cut-point, as recommended by the Society for Nicotine and Tobacco Research (Benowitz et al., 2002). Sensitivity analyses were also conducted using lower (10, 12 ng/ml) and higher (20 ng/ml) cut points recommended in other studies (Jarvis et al., 2008). Kappa statistics were computed to compare overlap between the self-reported and biochemically confirmed use of any tobacco product. Cochran-Mantel-Haenszel test procedure with Breslow-Day test for homogeneity of odds ratio was used to test whether discrepancies between self-reported and biochemically confirmed tobacco use differed by trial condition. A chi-square p-value <0.05 was set as the statistical significance level.

Results

Of the self-reported data and biochemical samples that were collected from 1250 youth, 26 biochemical samples could not be included in this study due to insufficient amount of saliva for testing. The 1224 youth in the final sample for biochemical validation included 611 control subjects (49.9%), 626 boys (50.12%), and 1004 school-going youth (80.38%). 652 (52.20%) of this sample were 10-14 years old. These demographic data were compared by trial condition, and no statistically significant differences were found.

Prevalence of tobacco use

Table 1 presents the prevalence of tobacco use. Self-reported current use of smoking (bidis or cigarettes) was 2.25% and that of chewing tobacco (Gutkha, Khaini, Zarda etc.) was 3.21%. Self-reported current use of any tobacco product was 4.11%. Mean value of salivary cotinine was 76.24 ng/ml for self-reported smokers; 121.74 ng/ml

Table 1. Prevalence of Current Use of Tobacco Products According to Different Measurement Tools (n=1224)

Tools & Classification	N	Prevalence % (95% CI)	Cotinine ng/ml Mean (SD)
Interviewer administered questionnaire (Self-reported current use)			
Cigarettes/Bidis	28	2.3 (1.4 - 3.1)	76.2 (81.5)
Chewing Tobacco	40	3.2 (2.3 - 4.4)	121.7 (85.3)
Any Tobacco	51	4.1 (3.0 - 5.2)	101.2 (87.0)
None	1173	95.8 (94.7 - 97.0)	8.6 (33.1)
Salivary Cotinine			
> 10 ng/ml	109	8.9 (7.3 - 10.5)	121.1 (78.5)
> 12 ng/ml	105	8.6 (7.0 - 10.2)	125.3 (76.9)
> 15 ng/ml	102	8.3 (6.8 - 9.9)	128.6 (75.5)
> 20 ng/ml	97	7.2 (6.4 - 9.4)	134.3 (73.1)

C.I, Confidence Interval; SD, Standard Deviation

Table 2. Agreement and Discrepancy between Self-reported Tobacco use and Salivary Cotinine Concentration for the Overall Sample (n=1224)

Self-reported tobacco use status ^a	CL > 15 ng/ml	CL ≤ 15 ng/ml	Total
	N Column % (95% C.I.)	N Column % (95% C.I.)	
User	37 36.3 (32.9 – 39.7)	14 1.2 (0.2 – 2.3)	51
Non-user	65 63.7 (60.3 – 67.1)	1108 98.8 (97.7 – 99.8)	1173
Total	102	1122	1224

CL, Cotinine level; C.I., Confidence Interval; ^aFor any tobacco product, past 30 days; sensitivity = 37/102 = 36.3%, specificity = 1108/1122 = 98.8%, positive predictive value = 37/51 = 72.6%, negative predictive value = 1108/1173 = 94.5%

Table 3. Agreement and Discrepancy between Self-reported Tobacco use and Salivary Cotinine Concentration by Trial Condition^b (n=1224)

Self-reported tobacco use status	Salivary cotinine (ng/mg)					
	Control (n=611)			Intervention (n=613)		
>15	≤15	Total	>15	≤15	Total	
User	25	7	32	12	7	19
Non-user	34	545	579	31	563	594
Total	59	552	611	43	570	613

^bChi² p-value = 0.37 (Used Breslow-Day test)

for self-reported tobacco chewers and 101.21 ng/ml for subjects who reported either smoking or chewing tobacco. Among youth who reported no tobacco use in the past month at the interviews, mean saliva cotinine was 8.6 ng/ml.

When results of the biochemical tests were reviewed, 8.33% (n=102) of the youth were identified as tobacco users at 15 ng/ml as the cut-point. Changing the cut point to 20, 12 and 10 ng/ml did not make much difference in prevalence rates (7.92%, 8.58% and 8.91%, respectively).

Self-report vs. biochemically confirmed tobacco use

At the cut-point of 15 ng/ml, agreement (the kappa statistic) between the self-reported and biochemically confirmed use of any tobacco product was 0.45 (95% C.I. 0.35 – 0.55). Changing the cut-points to 10, 12 or 20 ng/ml did not affect this finding.

Table 2 compares self-reported tobacco use to biochemical testing. Using the salivary cotinine test as gold standard, sensitivity and specificity of self-reported use of any tobacco product was 36.3% and 98.8% respectively.

Assuming that the prevalence of tobacco use in this sample was representative of that in the population from which the subjects were drawn, the interviewer administered questionnaire had a positive predictive value of 72.6% and negative predictive value of 94.5% for self-reported use of any tobacco product. Sensitivity and specificity of self-reports was similar between the younger subjects (10-14 years old) as compared to the older subjects (15-19 years old).

Self-report vs. biochemically confirmed tobacco use, by trial condition

As shown in Table 3, no statistically significant differences ($p > 0.05$) in the relationship between self-reported and biochemically confirmed prevalence were found between subjects in the two study conditions, at salivary cotinine level of 15 ng/ml. That is, we found no statistical evidence for differential misreporting of tobacco use by study condition at the baseline survey for Project ACTIVITY.

Discussion

Consistent with previous studies, biochemical validation of self-reported tobacco use using salivary cotinine found that youth were under-reporting tobacco use, overall (Behera et al., 2003; Caraballo et al., 2004). The difference of -4.2% is close to the mean difference between self-reported and salivary cotinine-based prevalence estimates found by a 2009 systematic review -4.8 (Gorber et al., 2009). Over 60% of youth who were confirmed to be tobacco users based on salivary cotinine levels self-reported themselves as non-users, although they gave consent at the beginning of the interviews for salivary sample collection.

The sensitivity of self-reported tobacco use in this study sample was lower than some previous studies (Caraballo et al., 2004; Kandel et al., 2006). Sensitivity of self-reports of current smoking in youth maybe lower when the overall prevalence and frequency of smoking is low, as in this study sample (Jarvis et al., 2008). Previous studies have suggested that youth are more likely to underreport socially stigmatized behaviors like tobacco use in household settings due to reduced anonymity with presence of parents, family and neighbors (Kann et al., 2002; Caraballo et al., 2004). In our study, although field workers were instructed to find a secluded spot away from the homes of the youth to conduct the interviews, this was often challenging, given the overcrowded setting in the slums. In addition to privacy concerns, youth may also underreport tobacco use behaviors since sale of tobacco to minors (under 18 years old) is now illegal in India (Tobacco Control Act of India, 2003). Some studies have reported higher sensitivity of self-reported tobacco use for older adolescents, but our results showed no significant differences by age group (Malcon et al., 2008). As was also reported by the SRNT, sensitivity and specificity in this study did not vary by much over a range of cut-points (10-20 ng/ml) (Gorber et al., 2009).

Comparison by trial condition showed that there was no differential misreporting of self-reported tobacco use at baseline between study conditions. This will need to be monitored at subsequent surveys after interventions, and if misreporting differs between study conditions, appropriate statistical correction will be applied in evaluation of Project ACTIVITY (Benowitz et al., 2002).

A few limitations should be taken into consideration here. The communities for participation in the trial were selected using certain eligibility criteria with non-random sampling methods. However, assignment to intervention or control condition, as well as selection of households

with youth for interviewing was randomized (Arora et al., 2010). Tobacco smoking within the seven days preceding sample collection is generally considered to be the cut-point at which cotinine can be detected (Benowitz et al, 2002). However, our baseline surveys only used 'past 30 days use' as current use (Post et al., 2005; Kandel et al., 2006). Also, no information was available regarding exposure to environmental tobacco smoke and nicotine dependence levels, which would help in interpreting discrepancies in self-reports (Molina et al., 2010).

To our knowledge, this study is among the first from India to use salivary cotinine tests in a community-based household survey and future results of biochemical validation from Project ACTIVITY will be useful to design validation studies in resource-poor settings.

In conclusions, biochemical validation of self-reported tobacco use among youth should be used at baseline and follow-up in any tobacco cessation intervention trial to effectively monitor and evaluate the potential impact of the interventions, as well as correct for any differential misreporting between control and intervention groups. Although there is no full substitution for self-reported tobacco use in tobacco surveillance studies, examining the validity of self-reported tobacco use would be beneficial to tobacco surveillance, research and interventions among adolescents in developing country settings like India, where tobacco use among youth is on the rise.

Acknowledgements

We are thankful to our partner NGOs: Prayas Juvenile Aid Centre Society, MANCH, All India Asaya Mahila Sahyog Samiti, Ankur who coordinated and supported us in implementation of the project. We would also like to thank our field staff and the community leaders who made this project successful. This research would not have been possible without the participation of children, young adults and their parents in selected communities of Delhi. We recognize Ms. Poojan Chandaliya and Dr Ruby Gupta for their assistance with laboratory analysis. The research was funded by a grant from Fogarty International Centre, National Institutes of Health, USA (Award Number RO1 TW007933). All contributing authors declare that none of them had any competing interest that would influence the results of this study.

References

- Adams J, Parkinson L, Sanson-Fisher RW, et al (2008). Enhancing self-report of adolescent smoking: The effects of bogus pipeline and anonymity. *Addictive Behavior*, **33**, 1291-6.
- Arora M, Tewari A, Tripathy V, et al (2010). Community-based model for preventing tobacco use among disadvantaged adolescents in urban slums of India. *Health Promot Int*, **25**, 143-52.
- Behera D, Uppal R, Majumdar S (2003). Urinary levels of nicotine & cotinine in tobacco users. *Indian J Med Res*, **118**, 129-33.
- Benowitz NL, Jacob P, Ahijevych K, et al (2002). Biochemical verification of tobacco use and cessation. *Nicotine Tobacco Res*, **4**, 149-59.
- Caraballo RS, Giovino GA, Pechacek TF (2004). Self-reported cigarette smoking vs. serum cotinine among US adolescents. *Nicotine Tobacco Res*, **6**, 19.
- Dhar P (2004). Measuring tobacco smoke exposure: quantifying nicotine/cotinine concentration in biological samples by colorimetry, chromatography and immunoassay methods. *J Pharmaceut Biomed Anal*, **35**, 155-68.
- Gorber SC, Schofield-Hurwitz S, Hardt J, et al (2009). The accuracy of self-reported smoking: A systematic review of the relationship between self-reported and cotinine-assessed smoking status. *Nicotine Tobacco Res*, **11**, 12-24.
- Iqbal SS, Mendoza K, Curran MJ, et al (2007). Salivary cotinine measurement in wave I of the social life health & aging project (NSHAP). Chicago core on biomarkers in population-based aging research.
- Jarvis MJ, Fidler J, Mindell J, et al (2008). Assessing smoking status in children, adolescents and adults: Cotinine cut points revisited. *Addiction*, **103**, 1553-61.
- Kandel DB, Schaffran C, Griesler PC, et al (2006). Salivary cotinine concentration versus self-reported cigarette smoking: Three patterns of inconsistency in adolescence. *Nicotine Tobacco Res*, **8**, 525-37.
- Kann L, Brener ND, Warren CW, et al (2002). An assessment of the effect of data collection setting on the prevalence of health risk behaviors among adolescents. *J Adolesc Hlth*, **31**, 327-35.
- Malcon MC, Menezes AMB, Assunção MCF, et al (2008). Agreement between self-reported smoking and cotinine concentration in adolescents: A validation study in Brazil. *J Adolesc Hlth*, **43**, 226-30.
- Mathur C, Stigler MH, Perry CL, et al (2008). Differences in prevalence of tobacco use among Indian urban youth: The role of socioeconomic status. *Nicotine Tobacco Res*, **10**, 109-16.
- Mishra A, Arora M, Stigler MH, et al (2005). Indian youth speak about tobacco: Results of focus group discussions with school students. *Hlth Educ Behavior*, **32**, 363.
- Molina AJ, Fernández D, Delgado M, et al (2010). Sensitivity and specificity of a self-administered questionnaire of tobacco use; including the fagerström test. *International J of Nursing Studies*, **47**, 181-9.
- Post A, Gilljam H, Rosendahl I, et al (2005). Validity of self reports in a cohort of Swedish adolescent smokers and smokeless tobacco (snus) users. *Tobacco Control*, **14**, 114.
- Reddy KS, Gupta PC (2004). Report on Tobacco Control in India. New Delhi: Ministry of Health and Family Welfare, Government of India.
- Reddy KS, Perry CL, Stigler MH, et al (2006). Differences in tobacco use among young people in urban India by sex, socioeconomic status, age, and school grade: Assessment of baseline survey data. *Lancet*, **367**, 589-94.
- Salimetrics (2006). Salivary DHEA Enzyme Immunoassay Kit. State College, PA.
- Salimetrics L (2009). High sensitivity salivary cotinine quantitative enzyme immunoassay kit.
- Sinha DN, Gupta PC, Reddy KS, et al (2008). Linking global youth tobacco survey 2003 and 2006 data to tobacco control policy in India. *J School Health*, **78**, 368-73.
- Stigler MH, Perry CL, Arora M, et al (2007). Intermediate outcomes from project MYTRI: Mobilizing Youth for Tobacco-Related Initiatives in India. *Cancer Epidemiology Biomarkers Prev*, **16**, 1050.
- Subramanian S, Nandy S, Kelly M, et al (2004). Patterns and distribution of tobacco consumption in India: Cross-sectional multi-level evidence from the 1998-9 National Family Health Survey. *BMJ*, **328**, 801.
- Warren C, Jones N, Eriksen M, et al (2006). Patterns of global tobacco use in young people and implications for future chronic disease burden in adults. *Lancet*, **367**, 749-53.
- Tobacco Control Act of India (2003). Available from: http://mohfw.nic.in/Tobacco_Control_Act-2003.pdf (accessed on August 6, 2011).