

## RESEARCH ARTICLE

# Apoptosis Induction in Human Leukemic Promyelocytic HL-60 and Monocytic U937 Cell Lines by Goniotalamin

Ploingarm Petsophonsakul<sup>1</sup>, Wilart Pompimon<sup>2</sup>, Ratana Banjerdpongchai<sup>1\*</sup>

### Abstract

Goniotalamin is an active compound extracted from *Goniotalamus griffithii*, a local plant found in northern Thailand. Goniotalamin inhibits cancer cell growth but is also toxic to normal cells. The aims of this study were to identify the cytotoxic effect of goniotalamin and the mechanism of cell death in human HL-60 and U937 cells. Cytotoxicity was determined by MTT assay and cell cycle profiles were demonstrated by staining with propidium iodide (PI) and flow cytometry. Apoptosis was confirmed by staining with annexin V-FITC/propidium iodide (PI) and flow cytometry. Reduction of mitochondrial transmembrane potential was determined by staining with dihexyloxycarbocyanine iodide and flow cytometry and expression of Smac, caspase-8 and -9 was demonstrated by Western blotting. Goniotalamin inhibited growth of HL-60 and U937 cell lines. An increase of SubG1 phase was found in their cell cycle profiles, indicating apoptosis as the mode of cell death. Apoptosis was confirmed by the flip-flop of phosphatidylserine using annexin V-FITC/PI assay in HL60 and U937 cells in a dose response manner. Furthermore, reduction of mitochondrial transmembrane potential was found in both cell types while expression of caspase-8, -9 and Smac/Diablo was increased in HL-60 cells. Taken together, our results indicate that goniotalamin-treated human leukemic cells undergo apoptosis via intrinsic and extrinsic pathways.

**Key words:** Goniotalamin - apoptosis - *Goniotalamus griffithii* – caspase - human leukemic cells

*Asian Pacific J Cancer Prev*, 14 (5), 2885-2889

### Introduction


Among constituents from the stems of *Goniotalamus griffithii* is goniotalamin (GTN), a plant bioactive styryl-lactone (Mu et al., 2003). GTN can also be extracted from several other species of *Goniotalamus* such as *Goniotalamus macrophyllus* (Alabsi et al., 2012), *Goniotalamus tapisoides* Mat Salleh (Kim et al., 2012) and *Goniotalamus andersonii* (Inayat-Hussain et al., 1999).

It is cytotoxic to and induces apoptosis in cancer cells such as cervical cancer (HeLa), breast carcinoma MCF-7 cell lines and colon cancer HT29 cells (Alabsi et al., 2012); Ca9-22 oral cancer cells (Yen et al., 2012); hepatoblastoma HepG2 cells (Al-Qubaisi et al., 2011); human lung cancer cells (Chiu et al., 2011); Jurkat T leukemia cells (Inayat-Hussain et al., 2010); and invasive breast carcinoma MDA-MB231 cell lines (Chen et al., 2005). It is also toxic to normal mouse fibroblast 3T3 cells (Alabsi et al., 2012), coronary artery smooth muscle cells (Chan et al., 2010), vascular smooth muscle cells (Chan et al., 2006). GTN induces apoptosis and autophagy in renal cancer cells and inhibits the nitric oxide synthase (NOS) activity and expression

(de Fátima et al., 2008). Furthermore, goniotalamin induces genotoxicity in Chinese ovary hamster (CHO) cells by inducing chromosome aberration (Umar-Tsafe et al., 2004). GTN, a plant styrylpyrone derivative, induces apoptosis in Jurkat T-cells as assessed by the externalization of phosphatidylserine via caspase-3 and -7 activation (Inayat-Hussain et al., 1999). It induces leukemic HL-60 cell apoptosis via the loss of mitochondrial transmembrane potential and caspase-9 activation (Inayat-Hussain et al., 2003). Even though there are several reports of the apoptotic induction of GTN in many cancer cell lines, the involvement of extrinsic pathway remains elusive.

The aims of this study were to determine the cytotoxic effect of GTN on HL-60 and U937 leukemic cell lines and the mechanisms of cell death involving both mitochondrial (intrinsic) and death receptor (extrinsic) pathways. The differences between the U937 and HL60 subclones are pronounced, the latter expressing fucose residues, which might be part of the CD15 cell adhesion molecules. The differences of the carbohydrate residues between the two cell lines can attribute to their differentiation within the myelomonocytic cell lineage (Schumacher et al., 1996).

<sup>1</sup>Department of Biochemistry, Faculty of Medicine, Chiang Mai University, Chiang Mai, <sup>2</sup>Faculty of Science, Lampang Rajabhat University, Lampang, Thailand \*For correspondence: [ratana.b@cmu.ac.th](mailto:ratana.b@cmu.ac.th)



**Figure 1. The Structure of Goniotalamin**

## Materials and Methods

### Chemicals

Goniotalamin was purified from the stem of *Goniotalamus griffithii* as previously reported (Mu et al., 2003) and the structure is shown in Figure 1. RPMI-1640 medium was obtained from Invitrogen, USA. MTT (3-(4,5-dimethyl)-2,5-diphenyl tetrazolium bromide), propidium iodide (PI) and 3,3'-dihexyloxycarbocyanine iodide (DiOC<sub>6</sub>) were obtained from Sigma-Aldrich (St. Louis, MO, USA). Rabbit polyclonal antibody to caspase-8, rabbit monoclonal antibody to caspase-9 and rabbit monoclonal antibody to Smac/Diablo and horseradish peroxidase (HRP) conjugated secondary antibody were purchased from Abcam, Cambridge, UK. SuperSignal West Pico Chemiluminescent Substrate was obtained from Pierce, Rockford, IL, USA. Complete mini protease inhibitor cocktail and annexin V-fluos staining kit were purchased from Roche, Indianapolis, IN, USA.

### Cell culture

Human leukemic promyelocytic HL-60 and human leukemic monocytic U937 cells were gifts from Dr. Sukhathida Ubol and Dr. Watchara Kasinroek, respectively. The cells were cultured in RPMI-1640 medium supplemented with 10% fetal bovine serum, penicillin G (100 units/ml) and streptomycin (100 µg/ml) at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub>. Goniotalamin was dissolved in dimethyl sulfoxide (DMSO) as a vehicle and the maximal volume used did not exceed 10 µl/ml of media. The human leukemic cells (1x10<sup>6</sup>) were treated with goniotalamin at indicated concentration and duration.

### MTT assay for cytotoxicity

HL-60 and U937 (3x10<sup>5</sup> cells/ml) were cultured and incubated with goniotalamin (0, 10, 20, 40 and 80 µM) at 37°C in 5% CO<sub>2</sub> atmosphere for 24 h. The cell viability was determined by using MTT assay (Banjerdpongchai et al., 2011). Briefly, MTT dye solution was added and incubated in CO<sub>2</sub> incubator for 4 h. Then, 100 µl of DMSO was added to dissolve the violet dye crystals. Absorbance was measured by using a microtiter plate reader (Biotek, USA) at 570 nm. The percentage of cell viability was calculated and 10, 20 and 50% inhibitory concentrations (IC<sub>10</sub>, IC<sub>20</sub> and IC<sub>50</sub>) were determined and used for further experiments. Since HL-60 cells were more sensitive to GTN than U937, the doses of GTN

were varied lower as follows: 0, 2, 4, 6, 8, 10 µM.

### Determination of cell cycle distribution

For flow cytometric assessment of DNA fragmentation and cell cycle distribution, 1x10<sup>6</sup> cells were harvested and re-suspended in a solution containing PI (50 µg/ml), 0.1% Triton X-100 and 0.1% sodium citrate in PBS. Cells then were analyzed in a FACScan equipped with a 488 nm argon laser using CellQuest software (Becton-Dickinson, USA) (Banjerdpongchai et al., 2010). Data were depicted as histograms and the percentage of cells displaying hypodiploid DNA content was indicated.

### Determination of phosphatidylserine externalization in apoptotic cells

Goniotalamin-treated cells were washed once in phosphate-buffered saline solution, centrifuged at 200 x g and the cell pellet was suspended in 100 µl of binding buffer provided by the annexin V-fluos staining kit. Annexin V-FITC (20 µl) and PI (10 µl) were added and the cell suspension was left at room temperature for 15 min in the dark. Finally, 970 µl of binding buffer were added. Analysis was conducted using FACScan (Becton Dickinson, USA). Cells that were stained with annexin V-FITC, and annexin V-FITC together with PI, were designated as early and late apoptotic cells, respectively (Pronmaban et al., 2012).

### Determination of mitochondrial transmembrane potential (MTP)

For MTP determination, 5x10<sup>5</sup> cells were treated with the GTN at IC<sub>10</sub>, IC<sub>20</sub> and IC<sub>50</sub> for 24 h, harvested and re-suspended in a PBS containing 40 nM of DiOC<sub>6</sub> (Wudtiwai et al., 2011). The cells were incubated for 15 min at 37°C and then subjected to flow cytometer (Becton Dickinson, USA).

### Western blot analysis

The goniotalamin-treated cells were washed once in ice cold PBS and incubated at 4°C for 10 min with ice-cold cell lysis buffer (250 mM sucrose, 70 mM KCl, 0.25% Triton X-100 in PBS containing complete mini protease inhibitor cocktail). Following centrifugation at 20,000 x g for 20 min, supernatant (50 µg, determined by Bradford method) was separated by 10% SDS-PAGE and 4% SDS-PAGE for stacking gel and then transferred onto nitrocellulose membrane. After treating with 5% non-fat milk in PBS containing 0.2% Tween-20, membrane was incubated with rabbit polyclonal antibody to caspase-8, rabbit monoclonal antibody to caspase-9 or rabbit monoclonal antibody to Smac/Diablo, followed by appropriate horseradish peroxidase (HRP)-conjugated secondary antibodies (1:20,000). Protein bands were visualized on X-ray film with SuperSignal West Pico Chemiluminescent Substrate (Pronmaban et al., 2012).


### Statistical analysis

Results are expressed as mean±S.D. Statistical


**Table 1. Inhibitory Concentrations at 10, 20 and 50 Percent (IC<sub>10</sub>, IC<sub>20</sub> and IC<sub>50</sub>) of Goniotalamin on Human Leukemic Cell Lines**

Cell types	HL-60	U937
IC <sub>10</sub> (μM)	1.95±1.4	1.15±0.1
IC <sub>20</sub> (μM)	2.94±1.7	2.71±0.1
IC <sub>50</sub> (μM)	5.67±2.0	11.5±1.1

\*The cytotoxicity effects of goniotalamin on human leukemic HL-60 and U937 cells were determined by MTT assay




**Figure 2. Cytotoxic Effect of Goniotalamin on Human Leukemic HL-60 and U937 Cell Lines.** HL-60 or U937 cells were treated with GTN at various concentrations for 24 h, then the cell viability was determined by the MTT assay. \*p<0.05, compared to control



**Figure 3. Histograms and Analysis of Percent Cells in Subdiploid Areas of Goniotalamin-treated Human Leukemic Cells.** HL-60 (A, B) and U937 (C, D) cells were treated with GTN for 24 h, stained with propidium iodide (PI) and processed by flow cytometry. Representative histograms and data are shown from three independent experiments. \*p<0.01, compared to control

difference between control and treated group was determined by one-way ANOVA (Kruskal Wallis analysis) at limit of p<0.05 from 3 independent experiments conducted in triplicate. For comparison between two groups, data were analyzed using Student's



**Figure 4. Dot Plots and Percent Analysis of Goniotalamin-treated Human Leukemic Cells for Phosphatidylserine Externalization of the Apoptotic Cells.** HL-60 (A, B) and U937 (C, D) cells were treated with GTN for 24 h and stained with annexin V-FITC and PI and processed by flow cytometer as described in Materials and Methods. Representative dot plots and data are shown from three independent experiments. \*p<0.05, compared to control t-test.

## Results

### Cytotoxicity of goniotalamin


GTN inhibited human HL-60 and U937 leukemic cell growth in a dose response manner. HL-60 cells were sensitive to GTN more than U937 cells with lower IC<sub>50</sub> (5.67 μM and 11.5 μM, respectively) as shown in Figure 2 and Table 1. Goniotalamin at IC<sub>10</sub>, IC<sub>20</sub> and IC<sub>50</sub> levels of each cell line were calculated and selected for further experiments.

### Cell cycle distribution


The cells were induced to die with the characteristic DNA content as subdiploid (subG1) or hypodiploid which is less than 2n (Figure 3A and 3C), shown as cells under P2 area. The percentage of cells with hypodiploid DNA content increased in a dose-dependent manner both in HL-60 and U937 cells (p<0.01) as shown in Figure 3B and 3D. The subG1 is a characteristic hallmark of apoptotic cells.

### Phosphatidylserine externalization and apoptotic induction

When HL-60 and U937 cells were treated with GTN at IC<sub>10</sub>, IC<sub>20</sub> and IC<sub>50</sub> levels for 24 h, and the flip-flop of phosphatidylserine (PS) was determined by using annexin V fluos staining kit. As shown in Figure 4A and 4B, the early apoptotic cell population (right lower



**Figure 5. Reduction of Mitochondrial Transmembrane Potential in Goniiothalamin-treated HL-60 and U937 Cells.** Histograms and data analysis of HL-60 (A, B) and U937 (C, D) cells represents the percentage of cells with loss of mitochondrial transmembrane potential (MTP) which are under P4 area. Representative histograms and data are shown from three independent experiments. \*p<0.05, compared to control



**Figure 6. Effect of Goniiothalamin on Initiation Caspase-8 and -9 and Apoptosis-related Smac/Diablo Protein Expressions in Human HL-60 Cells.** Immunoblots of caspase-8 and -9 and Smac/Diablo were determined in GTN-treated cells at various concentrations, viz. IC<sub>10</sub>, IC<sub>20</sub> and IC<sub>50</sub> levels. Representative bands (A) and band intensity analysis of caspase-8 (B), caspase-9 (C) and Smac/Diablo (D) are obtained from three independent experiments

quadrant) increased in a dose response manner in HL-60 cell lines. U937 cells were also induced to die via apoptosis with PS flip-flop out to the outer layers (Figure 4C and 4D). Both HL-60 and U937 cells responded to GTN in a concentration-dependent manner significantly at doses of IC<sub>10</sub>, IC<sub>20</sub> and IC<sub>50</sub> levels for HL-60 cells (p<0.01), whereas for U937 cells at doses of IC<sub>20</sub> and

IC<sub>50</sub> levels (p<0.01) (Figure 4B and 4D).

*Reduction of mitochondrial transmembrane potential (MTP)*

The loss of mitochondrial transmembrane potential was found in both HL-60 and U937 cells. The percentage of cells with reduction or loss of mitochondrial transmembrane potential (cells under P4 area) increased in a dose-dependent manner (p<0.05) as shown in Figure 5A, 5B for HL60 and Figure 5C, 5D for U937 cells.

*The expression of Smac/Diablo, caspase-8 and -9*

The human leukemic HL-60 and U937 cells were incubated with GTN at various concentrations for 24 h and the immunoblots of Smac/Diablo, caspase-8 and -9 were performed. The expression of Smac/Diablo, cleaved caspase-8 and cleaved caspase-9 increased in a concentration-dependent manner as shown in Figure 6A-6D. The pro-caspase-8 band intensity decreased and the cleaved form increased dose-dependently (Figure 6A, 6B), whereas the band intensity of cleaved caspase-9 (Figure 6A, 6C) and Smac/Diablo (Figure 6A, 6D) also increased in a concentration-dependent manner.

**Discussion**

Goniiothalamin is cytotoxic towards both cancerous (HGC-27, MCF-7, PANC-1, HeLa) and non-cancerous (3T3) cells but these cells die via necrotic cell death (Ali et al., 1997). However, many cancer cells, such as leukemic HL-60 cells, Jurkat T cells, colon HT29 cells were induced to undergo apoptosis by GTN as well (Alabsi et al., 2012; Inayat-Hussain et al., 2003; Inayat-Hussain et al., 2010; Inayat-Hussain et al., 1999). Even though there are reports of mechanisms of cancer cell apoptosis induced by GTN to be mitochondrial pathway, the cross-link between the extrinsic and intrinsic pathways remains elusive. The present study compared the characteristic of HL-60 and U937 apoptotic cell death, which underwent the same mechanism. IC<sub>50</sub> levels of GTN towards HL-60 and U937 cells were 5.67±2 and 11.5±1.1 μM, respectively. It has been shown that U937 cells were less sensitive to goniiothalamin than HL-60 cells.

The apoptosis of HL-60 and U937 cells was confirmed by the cell cycle profile demonstrating of the increase of subG1 population of both cells in concentration-dependent manner. The flip-flop of PS to the outer layer of cell membrane also occurred in HL-60 and U937 cells induced by GTN. The loss of mitochondrial transmembrane potential indicated the involvement of the mitochondrial pathway of apoptosis in both cell lines. The release of Smac/Diablo, a protein in the intermembranous space of mitochondria, into the cytosol increased as shown by immunoblot. Smac/Diablo acts as a negative regulator or an antagonist of inhibitors of apoptotic protein (IAPs), so it acts as a stimulator of apoptosis and is proposed to be of use in anti-cancer

treatment (de Almagro and Vucic, 2012).

The expression of caspase-9 and Smac/Diablo together with the reduction of MTP indicated the mitochondrial pathway of apoptotic cell death in HL-60 cells. However, the increase of cleaved caspase-8 expression was related to the extrinsic pathway. This suggests the cross-talk between the intrinsic and extrinsic pathways of apoptosis induced by goniiothalamine in human leukemic cell lines. The combined treatment of GTN and conventional chemotherapeutic drugs may be helpful in reducing the adverse effects of the chemotherapy and increase the apoptosis induction effect of such combination. The differences of the cell surface carbohydrate residues between HL-60 and U937 cells can be attributed to their differentiation within the myelomonocytic cell lineage (Schumacher et al., 1996). There are also differences in protein expression of angiopoietins and kindlins in human leukemic cells (Wu et al., 2012), HtrA2 and WT1 in acute myeloid leukemia (Li et al., 2012). Nevertheless, the mode of cell death induced by goniiothalamine in both cell lines showed a similar mode and mechanism of cell death as proven in the recent study but the extensiveness of apoptosis induction may be different according to the characteristics and sensitivity of each cell line. In conclusion, goniiothalamine induced human leukemic cells to die via apoptosis involving mitochondrial (caspase-9 and Smac) and death receptor (caspase-8) pathways.

## Acknowledgement

The authors thank National Research Council of Thailand (NRCT), Thailand Research Fund and the Commission of Higher Education, grant No. RMU5080003 for financial support.

## References

- Al-Qubaisi M, Rozita R, Yeap SK, et al (2011). Selective cytotoxicity of goniiothalamine against hepatoblastoma HepG2 cells. *Molecules*, **16**, 2944-59.
- Alabsi AM, Ali R, Ali AM, et al (2012). Apoptosis induction, cell cycle arrest and in vitro anticancer activity of goniiothalamine in a cancer cell lines. *Asian Pac J Cancer Prev*, **13**, 5131-6.
- Ali AM, Mackeen MM, Hamid M, et al (1997). Cytotoxicity and electron microscopy of cell death induced by goniiothalamine. *Planta Med*, **63**, 81-3.
- Banjerpongchai R, Kongtawelert P, Khantamat O, et al (2010). Mitochondrial and endoplasmic reticulum stress pathways cooperate in zearalenone-induced apoptosis of human leukemic cells. *J Hematol Oncol*, **3**, 50-65.
- Banjerpongchai R, Punyati P, Nakrob A, et al (2011). 4'-Hydroxycinnamaldehyde from *Alpinia galanga* (Linn.) induces human leukemic cell apoptosis via mitochondrial and endoplasmic reticulum stress pathways. *Asian Pac J Cancer Prev*, **12**, 593-8.
- Chan KM, Rajab NF, Ishak MH, et al (2006). Goniiothalamine induces apoptosis in vascular smooth muscle cells. *Chem Biol Interact*, **159**, 129-40.
- Chan KM, Rajab NF, Siegel D, et al (2010). Goniiothalamine induces coronary artery smooth muscle cells apoptosis: the p53-dependent caspase-2 activation pathway. *Toxicol Sci*, **116**, 533-48.
- Chen WY, Wu CC, Lan YH, et al (2005). Goniiothalamine induces cell cycle-specific apoptosis by modulating the redox status in MDA-MB-231 cells. *Eur J Pharmacol*, **522**, 20-9.
- Chiu CC, Liu PL, Huang KJ, et al (2011). Goniiothalamine inhibits growth of human lung cancer cells through DNA damage, apoptosis, and reduced migration ability. *J Agric Food Chem*, **59**, 4288-93.
- de Almagro MC, Vucic D (2012). The inhibitor of apoptosis (IAP) proteins are critical regulators of signaling pathways and targets for anti-cancer therapy. *Exp Oncol*, **34**, 200-11.
- de Fátima A, Zambuzzi WF, Modolo LV, et al (2008). Cytotoxicity of goniiothalamine enantiomers in renal cancer cells: involvement of nitric oxide, apoptosis and autophagy. *Chem Biol Interact*, **176**, 143-50.
- Inayat-Hussain SH, Annuar BO, Din LB, et al (2003). Loss of mitochondrial transmembrane potential and caspase-9 activation during apoptosis induced by the novel styryl-lactone goniiothalamine in HL-60 leukemia cells. *Toxicol In Vitro*, **17**, 433-9.
- Inayat-Hussain SH, Chan KM, Rajab NF, et al (2010). Goniiothalamine-induced oxidative stress, DNA damage and apoptosis via caspase-2 independent and Bcl-2 independent pathways in Jurkat T-cells. *Toxicol Lett*, **193**, 108-14.
- Inayat-Hussain SH, Osman AB, Din LB, et al (1999). Caspases-3 and -7 are activated in goniiothalamine-induced apoptosis in human Jurkat T-cells. *FEBS Lett*, **456**, 379-83.
- Kim RP, Bihud V, Bin Mohamad K, et al (2012). Cytotoxic and antioxidant compounds from the stem bark of *Goniiothalamus tapisoides* Mat Salleh. *Molecules*, **18**, 128-39.
- Li XY, Zhang Q, Li Y, et al (2012). Expression of HtrA2 and WT1 genes in acute myeloid leukemia. *Zhongguo Shi Yan Xue Ye Xue Za Zhi*, **20**, 1-6.
- Mu Q, Tang WD, Liu RY, et al (2003). Constituents from the stems of *Goniiothalamus griffithii*. *Planta Med*, **69**, 826-30.
- Prommaban A, Kodchakorn K, Kongtawelert P, et al (2012). *Houttuynia cordata* Thunb fraction induces human leukemic Molt-4 cell apoptosis through the endoplasmic reticulum stress pathway. *Asian Pac J Cancer Prev*, **13**, 1977-81.
- Schumacher U, Gräfin von Armansperg N, Kreipe H, Welsch U (1996). Lectin binding and uptake in human (myelo) monocytic cell lines: HL60 and U937. *Ultrastruct Pathol*, **20**, 463-71.
- Umar-Tsafe N, Mohamed-Said MS, Rosli R, et al (2004). Genotoxicity of goniiothalamine in CHO cell line. *Mutat Res*, **562**, 91-102.
- Wu WB, Zhang Q, Li Y, et al (2012). Expression of Kindlins and angiopoietins in acute myeloid leukemia. *Zhongguo Shi Yan Xue Ye Xue Za Zhi*, **20**, 7-11.
- Wudtiwai B, Sripanidkulchai B, Kongtawelert P, et al (2011). Methoxyflavone derivatives modulate the effect of TRAIL-induced apoptosis in human leukemic cell lines. *J Hematol Oncol*, **4**, 52-65.
- Yen CY, Chiu CC, Haung RW, et al (2012). Antiproliferative effects of goniiothalamine on Ca9-22 oral cancer cells through apoptosis, DNA damage and ROS induction. *Mutat Res*, **747**, 253-8.