

RESEARCH ARTICLE

The *XRCC1* Arg399Gln Gene Polymorphism and Risk of Colorectal Cancer: a Study in Kashmir

Nighat Parveen Khan¹, Arshad Ahmad Pandith², Adfar Yousuf¹, Nuzhat Shaheen Khan³, Mosin Saleem Khan¹, Imtiyaz Ahmad Bhat², Zahoor Wani Nazir¹, Khursheed Alam Wani⁴, Mahboob Ul Hussain⁵, Syed Mudassar^{1*}

Abstract

Background: The DNA repair gene *XRCC1* Arg399Gln gene polymorphism has been found to be implicated in the development of various cancers, including colorectal cancer (CRC), in different populations. We aimed to determine any association of this polymorphism with the risk of CRC in Kashmir. **Materials and Methods:** A total of 120 confirmed cases of CRC and 146 healthy cancer free controls from the Kashmiri population were included in this study. Genotyping was carried out by the polymerase chain reaction- restriction fragment length polymorphism (PCR-RFLP) method. **Results:** Genotype frequencies of *XRCC1* Arg399Gln observed in controls were 34.2%, 42.5% and 23.3% for GG (Arg/Arg), GA (Arg/Gln), AA (Gln/Gln), respectively, and 28.3%, 66.7% and 5% in cases, with an odds ratio (OR)=5.7 and 95% confidence interval (CI) =2.3-14.1 (p=0.0001). No significant association of Arg399Gln SNP with any clinicopathological parameters of CRC was found. **Conclusions:** We found the protective role of 399Gln allele against risk to the development of CRC. The *XRCC1* heterozygote status appears to be a strong risk factor for CRC development in the Kashmiri population.

Keywords: X-ray cross complementing gene - SNP - DNA repair - colorectal cancer - Kashmiri population

Asian Pac J Cancer Prev, 14 (11), 6779-6782

Introduction

Colorectal cancer (CRC) is the major cause of mortality and morbidity in western countries. It is third most common cancer in both men and women (Hisamuddin et al., 2006). As per cancer fact and figures for African-Americans an estimated 16,650 cases and 7,050 deaths from CRC occurred among African Americans in 2011. In Kashmir CRC ranks 4th common cancer among other cancers (Arshad et al. accepted for publication). Kashmiri population has a peculiar dietary habits and mode of living. Food of this region has been found to contain nitrosoamines, which are carcinogenic (Khuroo et al., 1992). Exposure to mutagenic and carcinogenic aromatic amines via environmental factors, well-cooked food and other sources can form DNA adducts in vivo and thus lead to DNA damage which if uncorrected can ultimately lead to tumorigenesis. However, integrity of damaged DNA is maintained by DNA repairing enzymes. *XRCC1*, *XRCC3*, and *XPD* are polymorphic genes belonging to 3 of the major DNA repair pathways. *XRCC1* [X-ray repair cross complementary] gene located on 19q13.31 chromosome encodes protein involved in base excision repair (BER) and the repair of single-strand breaks. The *XRCC1* gene

product plays an important role in the pathway by acting as a scaffold for other DNA repair proteins, such as DNA polymerase (Kubota et al., 1996) and DNA ligase III (Caldecott et al., 1994). The protein also has a BRCA1 C-terminus (BRCT) domain, which is characteristic of proteins involved in the recognition of and response to DNA damage. *XRCC1* interacts with poly (ADP-ribose) polymerase (PARP) by means of the BRCT domain to enable the recognition and subsequent repair of single-strand breaks (Caldecott et al., 1996; Masson et al., 1998).

DNA damage caused by a variety of internal and external factors, including ionizing radiation, alkylating agents, and oxidation, requires repair by the BER pathway. *XRCC1* lacking mutant hamster ovary cell lines are hypersensitive to ionizing radiation, hydrogen peroxide and alkylating agents, and are more susceptible to chromosomal aberrations compared to the parental cells. The importance of *XRCC1* to genetic stability is further indicated by that mice lacking *XRCC1* die in early embryogenesis (Tebbs et al., 1999).

DNA damage or reduced DNA repair capacity is viewed as an important mechanism in genetic instability and carcinogenesis caused by ionizing radiation and environmental chemical agents (Lei et al., 2013).

¹Biochemistry Department, ²Immunology and Molecular Medicine Department, ⁴Surgery Department, Sher-I-Kashmir Institute of Medical Sciences, ³Biochemistry Department, ⁵Biotechnology Department, University of Kashmir, Kashmir, India *For correspondence: parveennighat24@gmail.com, syed.mudassar@skims.ac.in

Polymorphisms in *XRCC1* gene that cause amino acid substitutions may impair the interaction of *XRCC1* with the other enzymatic proteins and consequently alter DNA strand break repair. Genetic polymorphisms in DNA repair genes may influence individual variation in DNA repair capacity, which may be associated with risk of developing cancer (Skjelbred et al., 2006; Wang et al., 2009).

Many studies have reported a link between Arg399Gln *XRCC1* SNP and increased risk of colorectal cancer (Krupa et al., 2004; Mariana et al., 2007) and also with other cancers like prostate, gastric, breast cancer, Hepatocellular Carcinoma, Glioma (Moullan et al., 2003, Rybicki et al., 2004; Liu et al., 2007; Wei-Hong et al., 2012). These studies have reported that AA genotype is associated with increased risk of colorectal cancer. Also GA genotype was found as a risk factor for progression of tumor growth. In contrast to this view some of the studies have reported protective role of AA allele (Olshan et al., 2002; Seedhouse et al., 2002; Skjelbred et al., 2006).

We aim to conduct a case-control study for *XRCC1* Arg399Gln gene polymorphism and to see its association with the risk of colorectal cancer in Kashmiri population.

Materials and Methods

A total of 120 blood samples of CRC patients were collected in the General surgery ward and clinics at Sher-I-Kashmir Institute of Medical Sciences [SKIMS] between May 2009 and January 2011. All the cases were histologically confirmed to be CRC. A pool of 146 control subjects were recruited from the same hospital and belonged to the same geographic area, ethnic background, and were of matching sex and age group. The controls did not have a previous diagnosis of any type of cancer. A written informed consent was obtained from each recruited subject, and the study was approved by the SKIMS ethical committee.

DNA extraction and polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP)

Genomic DNA was isolated by Phenol chloroform method from whole-blood samples of both cases and controls. Previously reported primers (Chih-Ching et al., 2005) were used for the amplification of the target regions of 615bp of the *XRCC1* Arg399Gln polymorphisms. PCR was carried out in a final volume of 25ul containing 50 ng genomic DNA template, 1x PCR buffer (Biotools, B&M Labs, Madrid, Spain) with 2mmol/L MgCl₂, 0.4mmol/L of each primer (Genescript, Piscataway, NJ), 50mmol/L dNTPs (Biotools, B&M Labs), and 1.25U Taq polymerase (Biotools, B&M Labs). For PCR amplification, the standard protocol was used as follows: one initial denaturation step at 94°C for 7 minutes, followed by 35 denaturation cycles of 30 seconds at 94°C, 30 seconds of annealing at 58°C, and 30 seconds of extension at 72°C, followed by a final elongation cycle at 72°C for 5 minutes. For RFLP, the PCR products of *XRCC1* SNP were digested with MspI (Fermentas, Inc., Glen Burnie, MD) 1U at 37°C for 4 hours. Digested PCR product was run on 3% Agarose gel and were visualized using ethidium bromide. GG allele revealed 375bp and 239bp fragments whilst the AA allele

was not digested by MspI and thus was visualized by 615bp single product. AG (heterozygote) allele revealed three bands of 615bp, 376 and 239bp (Figure 1).

Data analysis

The cases and controls were compared using the chi-square test for categorical variables, such as sex and smoking status, and other demographic variables. Crude odds ratios (OR) were used as estimates of the relative risk, and 95% confidence intervals (CI) were calculated to estimate the association between certain genotypes or other related risk factors of CRC.

Results

In this study 120 confirmed CRC cases and 146 healthy cancer free controls were included. In cases 63 (53%) were males and 57 (47%) females and the controls consisted of 77 (52.7%) male and 69 (47.26%) females. Of the total cases 54 (45%) were non-smokers and 66 (55%) were smoker. 52 (43%) of the cases were in the age group of less than 50 and 68 (57%) of the cases were in the age group of greater than 50. Among controls 66 (45%) were in less than fifty and 80 (55%) were in greater than fifty age group respectively (Table 1). *XRCC1* genotype distributions of wild AA and variant-type alleles GG among different clinico-pathologic characteristics are shown among the Kashmiri population for the cases. Patients with age group ≥ 50 years had marginally higher risk (OR=1.5) than cases with ≤ 50 years of age, but the difference is observed to be




Figure 1. A) Lane M 100bp DNA ladder. Lane 1-7 amplified 609 bp PCR product; and **B) Lane M 100 bp DNA Ladder.** Lane 2 AA, 1 & 3 GG, Lane 4, 5 GA genotype

Table 1. Comparison of Clinico-pathological Characteristics of Controls and Cases

Variables		Cases (n=120)	Controls (n=146)	OR (95%CI)	p value
Age group	<50	52(43%)	66(45%)	0.90(0.6-1.6)	0.75
	≥ 50	68(57%)	80(55%)		
Gender	Male	63(53%)	77(52.7%)	0.99(0.6-1.6)	0.96
	Female	57(47%)	69(47.26%)		
Smoking status	Never	54(45%)	60(34%)	1.17(0.7-1.8)	0.5
	Ever	66(55%)	86(66%)		
Residence	Rural	64(53.3%)	75(51.36%)	1.00(0.6-1.6)	0.7
	Urban	56(47%)	71(48.63%)		
Tumor site	Colon	72(60%)			
	Rectum	48(40%)			
Grade	WD	77(64%)			
	MD/PD	43(36%)			

Table 2. Association between XRCCI 399Arg/Gln Phenotypes and Clinic-pathological Characteristics of Cases

Variables	Cases (n=120)	Arg/Arg (GG)	Arg/Gln (GA)	Gln/Gln (AA)	OR (95%CI)	p value
		34(28.3%)	80(66.7%)	6(5%)		
Age group	<50	14	36	2	Reference	0.8
	≥50	20	44	4	0.9(0.63-1.2)	
Gender	Male	16	37	4	Reference	0.13
	Female	18	43	2	0.9(0.6-1.2)	
Smoking status	Never	15	34	3	Reference	0.9
	Ever	19	46	3	1.0(0.75-1.3)	
Residence	Rural	19	47	4	Reference	0.7
	Urban	15	33	2	0.9(0.7-1.2)	
Tumorsite	Colon	21	41	5	Reference	0.4
	Rectum	13	39	1	1.4(1.04-1.9)	
Grade	WD	21	49	5	Reference	0.9
	MD/PD	13	31	1	1.0(0.7-1.4)	

Table 3. Genotype Frequency of XRCCI Gene Polymorphism in Cases and Controls

Characteristic	Controls (N=146)	Cases (N=120)	OR (95%CI)	p value	
XRCCI polymorphism					
Overall	G/G	50	34	Reference	0.5
	A/A+G/A	96	86	1.3(0.6-2.0)	
Age <50 years	G/G	22	14	Reference	0.4
	A/A+G/A	43	38	1.4(0.6-3.1)	
≥50 years	G/G	28	20	Reference	0.5
	A/A+G/A	53	48	1.3(0.7-2.5)	
Male	G/G	31	16	Reference	0.3
	A/A+G/A	57	41	1.4(0.7-2.8)	
Female	G/G	19	18	Reference	0.6
	A/A+G/A	39	45	1.2(0.6-2.5)	
Never smoker	G/G	18	15	Reference	0.7
	A/A+G/A	38	37	1.2(0.5-2.8)	
Ever smoker	G/G	32	19	Reference	0.3
	A/A+G/A	58	49	1.4(0.7-2.7)	
Rural	G/G	26	19	Reference	0.5
	A/A+G/A	56	51	1.2(0.6-2.3)	
Urban	G/G	24	15	Reference	0.4
	A/A+G/A	40	35	1.4(0.56-3.08)	
Allele typed	G	162(55%)	148(62%)		0.15
	A	130(45%)	92(38%)		

statistically insignificant (Table 2).

The lesser frequency of AA genotype and increased frequency of GG/GA genotype in cases clearly show that AA act as protective genotypes while as GG/GA increases the risk of CRC. The distribution of genotype frequency of GG, GA and AA among controls were 34.2%, 42.5% and 23.3%, while in cases it was 28.3%, 66.7% and 5% respectively with (odds ratio OR=1.3 and 95% Confidence Interval: CI=0.6-2.0) (Table 2). The present study shows the higher frequency of GA genotype in cases than in controls (Table 3). The total frequency of allele G and A observed in our study in cases is 0.62 and 0.38 respectively, whereas in controls it is 0.55 and 0.45 respectively. *XRCCI* AA (399Gln) genotype frequency is higher in controls than in cases suggesting its protective role against the development of CRC. We did not find any significant association of any of the clinico-epidemiological characteristics of CRC with *XRCCI* Arg399Gln gene polymorphism (Table 3).

Discussion

XRCCI is an essential DNA repair gene involved in base

excision repair (BER) (Caldecott et al., 2003). The *XRCCI* gene exhibits polymorphic variations, including three common single nucleotide polymorphisms (SNPs) that result in amino acid substitutions in exon 7 (Arg194Trp), exon 9 (Arg280His) and exon 10 (Arg399Gln). These non-conservative amino acid alterations may influence DNA repair capability by altering the protein-protein interactions between *XRCCI* and other BER proteins. The Arg399Gln variant was found to be associated with several phenotypic alterations, including higher levels of sister chromatid exchange (Abdel-Rahman et al., 2000), aflatoxin B1-DNA adducts, glycophorin A mutations (Lunn et al., 1999) and polyphenol DNA adducts (Eric et al., 2000).

In this study carried out for the first time on Kashmiri population we tried to find out any possible association of the three variant alleles of *XRCCI* Arg399Gln gene with the development of CRC. The distribution of genotype frequency observed in our study in controls was 34.2%, 42.5% and 23.3% of GG, GA, AA while as in cases 28.3%, 66.7% and 5% respectively. Our findings suggest that the Arg399Gln polymorphism is associated with the risk of development of CRC. Our findings suggest that the Arg399Gln polymorphism is associated with the risk of development of CRC. No risk could be attributed to subjects with the 399Gln homozygote in the Kashmiri population. Several polymorphic studies have observed a positive association of the *XRCCI* 399 AA allele with various malignant diseases, including cancer of the head and neck (Erich et al., 1999) breast (Duell et al., 2001) lung (Divine et al., 2001) and colon and rectum (Abdel-Rahman et al., 2000). These conclusions corresponded well with the finding that subjects with the 399Gln allele have higher numbers of chromosomal breaks per cell than those with other genotypes (Wang et al., 2003), and that these subjects are more susceptible to chemically induced genetic damage (Li et al., 2003). However Camilla et al. (2006) found 399AA allele to be associated with a reduced risk of high-risk colorectal adenomas. Claire et al. (2002) found that the presence of the *XRCCI* 399Gln allele is protective against the development acute myeloblastic leukemia (Seedhouse et al., 2002; Skjelbred et al., 2006). These studies are in accordance with our study. We found that the presence of the *XRCCI* 399Gln allele is protective against the development of CRC. Allele frequency of Arg and Gln observed in our study in cases is 0.62 and 0.38 respectively, whereas in controls it is 0.55 and 0.45 respectively. A number of studies have reported that the *XRCCI* 399Gln allele has no or an inverse relationship with the risk of neoplasm development (Olshan et al., 2002; Shen et al., 2003; Smith et al., 2003; Hong et al., 2005).

The possible mechanism of the discrepancy that Gln is protective allele and Arg/Gln heterozygous allele is risk allele in the development of CRC could be that the altered protein has modified repair efficiency and increased level of DNA damage that results in apoptosis which ultimately leads to reduced risk of exposure induced cancer (Seedhouse et al., 2002; Jeon et al., 2005) and the possibility of *XRCCI*-independent single-strand break repair (Caldecott et al., 2003). The exact mechanism by

which the *XRCC1* 399Gln allele can have apparently opposite effects on the risk of neoplasm development is not known and should be further studied in detail.

In conclusion our study reports protective role of Gln allele compared to Arg/Gln allele in the risk of development of CRC in Kashmiri population however extensive studies are needed to ascertain this finding.

References

- Abdel-Rahman SZ, El-Zein RA (2000). The 399Gln polymorphism in the DNA repair gene *XRCC1* modulates the genotoxic response induced in human lymphocytes by the tobacco-specific nitrosamine NNK. *Cancer Lett*, **159**, 63-71.
- Abdel-Rahman SZ, Soliman AS, Bondy ML, et al (2000). Inheritance of the 194Trp and the 399Gln variant alleles of the DNA repair gene *XRCC1* are associated with increased risk of early-onset colorectal carcinoma in Egypt. *Cancer Lett*, **159**, 79-86.
- Caldecott KW, Aoufouchi S, Johnson P, Shall S (1996). *XRCC1* polypeptide interacts with DNA polymerase and possibly poly (ADP-ribose) polymerase, and DNA ligase III is a novel molecular 'nicksensor' in vitro. *Nucleic Acids Res*, **24**, 4387-94.
- Caldecott KW, McKeown CK, Tucker JD, Ljungquist S, Thompson LH (1994). An interaction between the mammalian DNA repair protein *XRCC1* and DNA ligase III. *Mol Cell Biol*, **14**, 68-76.
- Caldecott KW (2003). *XRCC1* and DNA strand break repair. *DNA Repair (Amst)*, **2**, 955-69.
- Cancer fact and figures for African-Americans 2011-2012.
- Chih-Ching Y, Fung-Chang S, Reiping T, Chung RCC, Ling-Ling H (2005). Polymorphisms of the *XRCC1*, *XRCC3*, & *XPB* genes, and colorectal cancer risk: a case-control study in Taiwan. *BMC Cancer*, **5**, 12.
- Divine KK, Gilliland FD, Crowell RE, et al (2001). The *XRCC1* 399 glutamine allele is a risk factor for adenocarcinoma of the lung. *Mutat Res*, **461**, 273-8.
- Duell EJ, Millikan RC, Pittman GS, et al (2001). Polymorphisms in the DNA repair gene *XRCC1* and breast cancer. *Cancer Epidemiol Biomarkers Prev*, **10**, 217-22.
- Eric JD, John KW, Tsun-Jen C, et al (2000). Polymorphisms in the DNA repair genes *XRCC1* and *ERCC2* and biomarkers of DNA damage in human blood mononuclear cells. *Carcinogenesis*, **21**, 965-71.
- Erich MS, Edward JC, Lei L, et al (1999). Polymorphisms of DNA repair gene *XRCC1* in squamous cell carcinoma of the head and neck. *Carcinogenesis*, **20**, 2125-9.
- Hisamuddin IM, Yang VW (2006). Molecular Genetics of Colorectal Cancer: An Overview. *Curr Colorectal Cancer Rep*, **2**, 53-9.
- Hong YC, Lee KH, Kim WC, et al (2005). Polymorphisms of *XRCC1* gene, alcohol consumption and colorectal cancer. *Int J Cancer*, **116**, 428-32.
- Jeon YT, Kim JW, Park NH, et al (2005). DNA repair gene *XRCC1* Arg399Gln polymorphism is associated with increased risk of uterine leiomyoma. *Human Reproduction*, **20**, 1586-9.
- Khuroo MS, Zargar SA, Mahajan R, Banday MA (1992). High incidence of oesophageal and gastric cancer in Kashmir in a population with special personal and dietary habits. *Gut*, **33**, 11-5.
- Krupa R, Blasiak J (2004). An association of polymorphism of DNA repair genes *XRCC1* and *XRCC3* with colorectal cancer. *J Exp Clin Cancer Res*, **23**, 285-94.
- Kubota Y, Nash RA, Klungland A, et al (1996). Reconstitution of DNA base excision-repair with purified human proteins: interaction between DNA polymerase and the *XRCC1* protein. *EMBO J*, **15**, 6662-70.
- Lei J, Xiao F, Yi B, Jue-Yu Z, Xiao-Yan S (2013). "Association between the *XRCC1* Polymorphisms and Glioma Risk: A Meta-Analysis of Case-Control Studies" *PLoS ONE*, **8**, 55597
- Li Y, Marion MJ, Rundle A, Brandt-Rauf PW (2003). A common polymorphism in *XRCC1* as a biomarker of susceptibility for chemically induced genetic damage. *Biomarkers*, **8**, 408-14.
- Liu B, Wei J, Zou Z, et al (2007). Polymorphism of *XRCC1* predicts overall survival of gastric cancer patients receiving oxaliplatin-based chemotherapy in Chinese population. *European J Human Genetics*, **15**, 1049-53.
- Lunn RM, Langlois RG, Hsieh LL, Thompson CL, Bell DA (1999). *XRCC1* polymorphisms: effects on aflatoxin B1-DNA adducts and glycophorin A variant frequency. *Cancer Res*, **59**, 2557-61.
- Mariana CS, David VC, Kimberly DS (2007). DNA repair single-nucleotide polymorphisms in colorectal cancer and their role as modifiers of the effect of cigarette smoking and alcohol in the Singapore Chinese Health Study. *Epidemiol Biomarkers Prev*, **16**, 2363-72.
- Masson M, Niedergang C, Schreiber V, et al (1998). *XRCC1* is specifically associated with poly(ADP-ribose) polymerase and negatively regulates its activity following DNA damage. *Mol Cell Biol*, **18**, 3563-71.
- Moullan N, Cox DG, Angèle S, et al (2003). Polymorphisms in the DNA Repair Gene *XRCC1*, Breast Cancer Risk, and Response to Radiotherapy. *Cancer Epidemiol Biomarkers Prev*, **12**, 1168-74.
- Olshan AF, Watson MA, Weissler MC, Bell DA (2002). *XRCC1* polymorphisms and head and neck cancer. *Cancer Lett*, **178**, 181-6.
- Rybicki BA, Conti DV, Moreira A, et al (2004). DNA Repair Gene *XRCC1* and *XPB* Polymorphisms and Risk of Prostate Cancer. *Cancer Epidemiol Biomarkers Prev*, **13**, 23-9.
- Seedhouse C, Bainton R, Lewis M, et al (2002). The genotype distribution of the *XRCC1* gene indicates a role for base excision repair in the development of therapy-related acute myeloblastic leukemia. *Blood*, **10**, 3761-6.
- Shen M, Hung RJ, Brennan P, et al (2003). Polymorphisms of the DNA repair genes *XRCC1*, *XRCC3*, *XPB*, interaction with environmental exposures, and bladder cancer risk in a case-control study in northern Italy. *Cancer Epidemiol Biomarkers Prev*, **12**, 1234-40.
- Skjelbred CF, Saebo M, Wallin H, et al (2006). Polymorphisms of the *XRCC1*, *XRCC3* and *XPB* genes and risk of colorectal adenoma and carcinoma, in a Norwegian cohort: a case control study. *BMC Cancer*, **6**, 67.
- Smith TR, Miller MS, Lohman K, et al (2003). Polymorphisms of *XRCC1* and *XRCC3* genes and susceptibility to breast cancer. *Cancer Lett*, **190**, 183-90.
- Tebbs RS, Flannery ML, Meneses JJ, et al (1999). Requirement for the *XRCC1* DNA base excision repair gene during early mouse development. *Dev Biol*, **208**, 513-29.
- Wang Y, Spitz MR, Zhu Y, et al (2003). From genotype to phenotype: correlating *XRCC1* polymorphisms with mutagen sensitivity. *DNA Repair (Amst)*, **2**, 901-8.
- Wang Y, Yang H, Li H, et al (2009). Association between X-ray repair cross complementing group 1 codon 399 and 194 polymorphisms and lung cancer risk: A meta-analysis. *Cancer Lett*, **285**, 134-40.
- Wei-Hong D, Zhen-Yu Z, Jun-Gui L, et al (2012). "XRCC1 Arg399Gln Gene Polymorphism and Hepatocellular Carcinoma Risk in the Chinese Han Population: A Meta-analysis" *Asian Pac J Cancer Prev*, **13**, 3601-4.