

## RESEARCH ARTICLE

Editorial Process: Submission:01/06/2025 Acceptance:06/04/2025

# Antioxidant property of Silver Nanoparticles Loaded with Alcoholic Extraction of Lycium Shawii

Ali Noory Fajer<sup>1\*</sup>, Meaad Nasser Hussein<sup>1</sup>, Yusra Sebri Abdulsahab<sup>2</sup>

## Abstract

**Background:** Antioxidants play a crucial role in mitigating oxidative stress associated with cancer progression. Lycium shawii is noted for its high content of antioxidants and anti-inflammatory compounds. Silver nanoparticles (AgNPs) stand out as the best among the numerous synthesized and characterized nanoparticles that have been made thus far just to give an example. **Objective:** This study is carried out to synthesize, characterize and later investigate the in vivo antioxidant activity of AgNPs loaded with the alcoholic extract of Lycium shawii. **Methods:** Thirty rats were randomly distributed among five groups. Each group contained six rats. Group 1 (Control) was given 1 mL of normal saline orally daily through a stomach tube. Group 2 was injected with 1 mL of AgNPs at a dose of 100 mg/kg for one day only. Group 3 was treated with 1 mL/day of Acetamiprid at a dose of 25 mg/kg. Group 4 was given AgNPs (100 mg/kg) for three weeks and then treated with Acetamiprid (25 mg/kg) for an additional three weeks. Group 5 was treated with Acetamiprid (25 mg/kg) for three weeks and then AgNPs (100 mg/kg). All treatments were done daily. Serum levels of T-AOC and TOS as well as AST, ALT, urea, and creatinine were measured in all groups. **Results:** A statistically significant rise in TOS, ALT, AST, urea, and creatinine, along with a statistically significant reduction in T-AOC were observed ( $p \leq 0.05$ ) in Group 3 as compared to Groups 1, 2, 4, and 5. **Conclusion:** *In vivo* results showed that AgNPs, especially in combination with Lycium shawii extracts, seemed to exhibit high antioxidant activity, and was able to provide protection against oxidative stress. Retrieval of these results could contribute to a good indication about the potential use of AgNPs in cancer treatment, and as a drug in pandemic of oxidative stress-mediated disease. The molecular basis of its antioxidant effects and the long-term safety and efficacy of the use of AgNPs in clinical practice are yet to be determined.

**Keywords:** Silver nanoparticles- lycium Shawii- cancer- antioxidant activity

*Asian Pac J Cancer Prev*, 26 (6), 2197-2204

## Introduction

Antioxidants play a crucial role in combating oxidative stress associated with cancer development [1]. Cancer cells often exhibit elevated levels of reactive oxygen species (ROS), which contribute to oxidative stress. This oxidative stress can lead to DNA damage and mutations, thereby promoting tumorigenesis. Antioxidants can mitigate these effects by neutralizing ROS, thereby reducing the risk of cancer progression [2-5]. Medicinal plants and dietary supplements are recognized as valuable sources of food antioxidants. Antioxidants derived from natural food sources are generally regarded as safer than synthetic supplements. The interaction of dietary antioxidants with other nutrients can enhance their bioavailability and efficacy, making them more effective compared to isolated supplements [6]. Lycium shawii is rich in antioxidants, anti-inflammatory compounds, and antimicrobial or cytotoxic agents, making it a valuable resource for drug development and dietary supplements

[7]. The active chemical compounds in plant extracts serve as the primary basis for their pharmacological activities. Nanotechnology is a relatively new field in the modern world; however, materials with structural nanometer dimensions are not novel and have significant applications [8]. Nanotechnology offers innovative tools for developing more efficient and secure nanomedicines, presenting various potential benefits in drug formulation and delivery. It involves the engineering and industrialization of materials at the molecular and atomic scales [9, 10]. Currently, different procedures are used for the production of metallic nanoparticles, including biological, chemical, and physical methods. Mechanical milling and high-energy mechanical milling are effective physical methods for synthesizing nanoparticles (NPs). The chemical method typically involves the use of chemical substances for the preparation of NPs, which renders them unsuitable for various applications due to the inclusion of toxic compounds [11]. Among the various nanoparticles synthesized and characterized to date, silver nanoparticles

<sup>1</sup>Department of Chemistry, College of Education, University of Al-Qadisiyah, Iraq. <sup>2</sup>Clinical Pharmacy Branch, College of Pharmacy, Misan University, Iraq. \*For Correspondence: ali.fajer@qu.edu.iq

(AgNPs) hold a prominent position, primarily due to their remarkable ability to act as antibacterial agents, even in solid form. Despite the long-standing recognition of silver's significance, its potential has largely been underutilized, with applications historically limited to coins and traditional medicine. Currently, approximately 320 tons of silver nanoparticles are produced annually, finding applications in food safety, biosensing, and nanomedical imaging [12]. Given the widespread use of silver nanoparticles, risk assessment of these nanoparticles is crucial, particularly in large-scale applications. Numerous studies have demonstrated the effectiveness of nanoparticles and their biological and cellular effects. However, an increase in reactive oxygen species (ROS) can lead to the breakdown of the antioxidant defense system, resulting in damage to DNA, lipids, and proteins [13]. Antioxidant systems consist of enzymatic and non-enzymatic components. Non-enzymatic antioxidants include water-soluble and membrane-bound molecules, such as ascorbic acid and glutathione. Antioxidant enzymes, including superoxide dismutase (SOD), catalase, and glutathione peroxidase (GPX), play critical roles in defending against oxidative stress. Reduced glutathione, a sulfur-containing tripeptide, can either scavenge free radicals or act as a cofactor for GPX, which oxidizes glutathione-S-transferase (GST), thereby reducing the activity of SOD, the first line of defense against ROS. Previous research has shown that ROS acts as a biomarker for increased production and induction of oxidative stress, which can be altered by the activity of silver nanoparticles [14-16].

Acetamiprid, classified as a neonicotinoid insecticide, has the chemical formula  $C_{10}H_{11}ClN_4$ . It is an organic compound used to manage and control sucking insects that threaten a diverse range of crops, including leafy vegetables, fruiting vegetables, and cole crops. Acetamiprid is manufactured by Aventis CropScience and is commercially available under brand names such as Assail and Chipco [17]. The compound exerts its effects by binding to specific receptors within the synaptic cleft of nerve cells, thereby interfering with signal propagation in the insect's nervous system. Acetamiprid shares structural similarities with other neonicotinoids, such as imidacloprid, nitenpyram, and thiacloprid. It features a 6-chloro-3-pyridine methyl group, similar to these compounds, but differs in the substituent present on an acyclic or cyclic moiety, which may include nitroguanidine, nitromethylene, or cyanoamidine. Additionally, acetamiprid, structurally similar to nicotine, exhibits physiological effects comparable to nicotine in the human body. Nicotine, a naturally occurring insecticide, serves as the basis for synthesizing several synthetic insecticides [18, 19]. Acetamiprid is used in agriculture in various forms, such as wettable powders and water-soluble packs. It is essential to follow prescribed application instructions and safety measures to minimize potential risks to human health and the environment [20]. Furthermore, in this study we aimed to synthesize and characterize silver nanoparticles loaded with alcoholic extracts of *Lycium shawii* and evaluate their antioxidant effects *in vivo*.

## Materials and Methods

### *Preparation of Extract from Lycium shawii*

The *Lycium shawii* plant was collected from the outskirts of Diwaniyah. The leaves were thoroughly washed several times with plain water, followed by rinsing with distilled water to remove impurities. After washing, the leaves were shade-dried and ground into a fine powder using an electric mixer. The powder was then stored for future use. Subsequently, 20 grams of the crushed leaves were heated in 200 milliliters of filtered water at 45 °C for 30 minutes in a shaking water bath. After heating, the solution was wrapped in cellophane paper and left to stand for 24 hours before being filtered through filter paper [21].

### *Synthesis of Silver Nanoparticles*

A mixture of 30 mL of 1 mM silver nitrate solution in water and 70 mL of the extract was prepared at room temperature with a pH of 8.5. The mixture was heated in a water bath and agitated at 45°C for 80 minutes to ensure thorough mixing. The appearance of a dark brown color indicated the formation of silver nanoparticles.

### *Experimental animals*

The animals were acclimatized for two weeks prior to the experiments. Rats were divided into five groups, each consisting of six male rats (n=6). Group 1 (Control) received 1 mL of normal saline orally via a stomach tube daily. Group 2 received 1 mL of silver nanoparticles (AgNPs) at a dosage of 100 mg/kg on the day of administration. Group 3 was administered 1 mL/day of Acetamiprid at a dosage of 25 mg/kg. Group 4 received AgNPs (100 mg/kg) for three weeks, followed by Acetamiprid (25 mg/kg) for an additional three weeks. Group 5 was given Acetamiprid (25 mg/kg) for three weeks, followed by silver nanoparticles (100 mg/kg). All doses were administered daily, and serum levels of total antioxidant capacity (T-AOC) [22], total oxidant status (TOS) [23], AST, ALT [24], urea, and creatinine were assessed in all groups [25].

### *Statistical Analysis*

The experiments were carried out three times, and the data shown comes from three separate trials. Statistical analysis was performed using Graph Pad Prism version 8.4.3, with results reported as mean  $\pm$  standard deviation (Mean  $\pm$  SD). A paired Student's t-test and one-way ANOVA were used to evaluate statistical significance, considering p-values less than 0.05 as statistically significant.

## Results

### *UV-Vis Spectrum, SEM, and XRD analysis*

Figure 1 illustrates the UV-Vis spectra results of silver nanoparticles in a 0.1 mM silver nitrate solution. Figure 2 presents the SEM micrographs of the synthesized silver nanoparticles. The Figure illustrates the irregular morphology and size of the synthesized nanoparticles, which were found to fall within the accepted nanoparticle size range of 1 to 100 nm.

X-ray diffraction (XRD) analysis was conducted to determine the size distribution and characteristics of the crystal structure. The results of the XRD analysis are presented in Figure 3.

#### Biochemical parameters

Table 1 shows the effects of green-synthesized silver nanoparticles on oxidative stress in male rats. Compared to the effects of N-[(6-chloro-3-pyridyl) methyl]-N'-cyano-N-methyl-acetamide, a chemical compound, the nature of its interaction with biological systems and the extent to which it contributes to oxidative stress remain unknown.

Table 2 shows the role of green-synthesized silver nanoparticles on liver enzymes in male rats. The results indicate a significant increase ( $p \leq 0.05$ ) in ALT and AST enzyme levels in Group 3 (G3) compared to Groups 2

(G2), 4 (G4), and 5 (G5).

#### Kidney Function

As shown in Table 3, the levels of urea and creatinine increased in Group 3 (G3) compared to Groups 2 (G2), 4 (G4), and 5 (G5).

#### Histological Analysis of Liver and Kidney Tissues

After examining the histological slides of sections taken from the liver and kidney, the results of our study were consistent with the physiological findings, revealing pathological histological changes in the liver and kidney tissues.

#### Liver Tissue Analysis

Figure 4 presents various cross-sectional images of liver tissue. Image (A) depicts the control group,




Figure 1. The Result of UV-Vis Spectra of Silver Nanoparticles in 0.1 mM Silver Nitrate Solution Concentration.

Table 1. The Effect of Green-Synthesized Silver Nanoparticles on Oxidative Stress Markers.

Oxidative stress			
T-AOC ( $\mu\text{mol/L}$ )		TOS ( $\mu\text{mol/L}$ )	
Mean $\pm$ SD	P-value	Mean $\pm$ SD	P-value
0.79 $\pm$ 0.06	G1 & G2 = 0.67	1.75 $\pm$ 0.44	G1 & G2 = 851
	G1 & G3 = 0.000*		G1 & G3 = 0.000*
	G1 & G4 = 0.000*		G1 & G4 = 0.000*
	G1 & G5 = 0.000*		G1 & G5 = 0.000
0.81 $\pm$ 0.05	G2 & G3 = 0.000*	1.8 $\pm$ 0.33	G2 & G3 = 0.000*
	G2 & G4 = 0.000*		G2 & G4 = 0.000*
	G2 & G5 = 0.000*		G2 & G5 = 0.000*
0.11 $\pm$ 0.033	G3 & G4 = 0.61	4.54 $\pm$ 0.55	G3 & G4 = 0.025*
	G3 & G5 = 0.013*		G3 & G5 = 0.003*
0.14 $\pm$ 0.32	G4 & G5 = 0.04*	3.88 $\pm$ 0.46	G4 & G5 = 0.353
0.244 $\pm$ 0.129		3.62 $\pm$ 0.31	

\*Sing to difference significant at  $P \leq 0.05$




Figure 2. SEM Analysis of AgNPS Synthesised by Lycium Shawii.L. Plant Extraction

highlighting the central vein (yellow arrow), blood pockets (blue arrow), and hexagonal hepatocytes arranged in strands (red arrow). Image (B) shows the group treated with the extract, where similar features in the liver tissue are observed, including the central vein (yellow arrow), blood pockets (blue arrow), and hexagonal hepatocytes arranged in strands (red arrow). In images (C) and (D), which represent the group treated with acetaminophen, blood congestion in the central vein (yellow arrow) is noted, along with necrosis (indicated by other arrows) and the accumulation of inflammatory cells (black arrow). Image (E) illustrates the group treated with acetaminophen followed by the extract, revealing natural tissue characteristics, including the central vein (yellow arrow), blood pockets

(blue arrow), and hexagonal hepatocytes arranged in strands (red arrow). Finally, image (F) depicts the group treated with the extract followed by acetaminophen, where a slight expansion of the central vein (yellow arrow) and mild decay (blue arrow) are observed, while the hepatocytes appear normal and arranged in strands (red arrow).

#### Kidney Tissue Analysis

Figure 5 presents various cross-sections of kidney tissue from both control and treated group rats. Image (A) illustrates the kidney tissue of the control group, highlighting nearby urinary tubules (green arrow), distant urinary tubules (black arrow), glomeruli (blue arrow), and




Figure 3. The XRD Analysis of AgNPS Synthesized by Lycium Shawii.L. plant Extraction



Table 2. The Effect of Green-Synthesized Silver Nanoparticles on Liver Enzymes Level

Group	Liver enzyme			
	ALT (U/L)		AST (U/L)	
	Mean $\pm$ SD	P-value	Mean $\pm$ SD	P-value
G1	30.4 $\pm$ 1.67	G1 & G2 = 0.986 G1 & G3 = 0.000* G1 & G4 = 0.000* G1 & G5 = 0.000*	63.56 $\pm$ 3.79	G1 & G2 = 0.725 G1 & G3 = 0.000* G1 & G4 = 0.000* G1 & G5 = 0.000*
G2	30.38 $\pm$ 1.84	G2 & G3 = 0.000* G2 & G4 = 0.000* G2 & G5 = 0.000*	62.26 $\pm$ 4.6	G2 & G3 = 0.000* G2 & G4 = 0.000* G2 & G5 = 0.000*
G3	59.76 $\pm$ 1.17	G3 & G4 = 0.000* G3 & G5 = 0.000*	107.31 $\pm$ 6.05	G3 & G4 = 0.129 G3& G5 =0.001*
G4	47.92 $\pm$ 1.92	G4 & G5 = 0.001*	101.56 $\pm$ 8.9	G4&G5 = 0.021*
G5	43.34 $\pm$ 2.1		92.98 $\pm$ 3.7	

\*Sing to difference significant at  $P \leq 0.05$




Figure 4. Histological Slides of the Sections taken from the Liver. (A): control group, (B): group treated with extract, (C and D): group treated with acetaminophen, (E): group received acetaminophen followed by extract, (F): group received extract followed by acetaminophen.

Bowman's capsule (yellow arrow). Image (B) depicts the kidney tissue of the group treated with the extract, showing similar structures: nearby urinary tubules (green arrow), distant urinary tubules (black arrow), glomeruli (blue

Table 3. The Effect of Green-Synthesized Silver Nanoparticles on Kidney Function in Male Rats

Group	Kidney function			
	Urea (mg/dl)		Creatinine (mg/dl)	
	Mean $\pm$ SD	P-value	Mean $\pm$ SD	P-value
G1	6.63 $\pm$ 0.41	G1 & G2 = 0.642 G1 & G3 = 0.000* G1 & G4 = 0.000* G1 & G5 = 0.001*	1.46 $\pm$ 0.4	G1 & G2 = 0.786 G1 & G3 = 0.000* G1 & G4 = 0.000* G1 & G5 = 0.000*
G2	6.30 $\pm$ 0.40	G2 & G3 = 0.000* G2 & G4 = 0.000* G2 & G5 = 0.000*	1.36 $\pm$ 0.35	G2 & G3 = 0.000* G2 & G4 = 0.000* G2 & G5 = 0.000*
G3	16.32 $\pm$ 2.06	G3 & G4 = 0.000* G3 & G5 = 0.000*	4.68 $\pm$ 0.75	G3 & G4 = 0.04 * G3 & G5 = 0.01 *
G4	10.68 $\pm$ 0.96	G4 & G5 = 0.56	3.88 $\pm$ 0.55	G4 & G5 = 0.094
G5	9.26 $\pm$ 0.76		3.24 $\pm$ 0.72	

\*Sing to difference significant at  $P \leq 0.05$




Figure 5. Histological Slides of the Sections Taken from the Kidney. (A): control group, (B): group treated with extract, (C and D): group treated with acetaminophen, (E): group received acetaminophen followed by extract, (F): group received extract followed by acetaminophen.

arrow), and Bowman's capsule (yellow arrow). Images (C) and (D) display cross-sections of kidney tissue from the group treated with acetaminophen, where atrophy in the glomeruli (black arrow), blood congestion (red arrow), and bleeding (yellow arrow) are observed. Image (E) shows a cross-section of kidney tissue from the group treated with both acetaminophen and the extract, revealing natural tissue with nearby urinary epithelium (green arrow), distant urinary tubules (black arrow), glomeruli (blue arrow), and Bowman's capsule (yellow arrow). Finally, Image (F) presents a cross-section of kidney tissue from the group treated with the extract followed by acetaminophen, where a slight expansion of Bowman's capsule (yellow arrow), mild atrophy in the glomeruli (blue arrow), and expansion of distant urinary tubules (black arrow) are noted.

## Discussion

The homogenized extract demonstrated superior nanoparticle production in the synthesis of silver nanoparticles from the *Lycium shawii* L. plant extract. The reaction environment confirmed the presence of silver nanoparticles. Due to surface plasmon resonance, the color of the reaction medium gradually changed to dark brown. The surface plasmon band is influenced by factors such as the dielectric environment, morphology, size, composition, and shape of the nanoparticles. The degree of absorption increased with the synthesis of more silver nanoparticles, as indicated by previous research [26]. The visible range of UV spectra for silver nanoparticles (AgNPs) typically falls between 410 and 500 nm, while the absorption in this study ranged from 420 to 450 nm. The scanning electron microscope (SEM) has enhanced our understanding of the morphological characteristics of the *Lycium shawii* L. plant, from which silver nanoparticles (AgNPs) were synthesized. Figure 2 presents the SEM micrographs of the synthesized silver nanoparticles [27]. The Figure illustrates the irregular morphology and size of the synthesized nanoparticles,

which were found to fall within the accepted nanoparticle size range of 1 to 100 nm.

Different chemical substances may exert varying impacts on oxidative stress, depending on their mechanisms of action and concentrations. In the absence of specific data or studies on this particular compound, we can only provide general insights into how chemical compounds might be associated with oxidative stress [28]. Substances can act as either pro-oxidants, which contribute to oxidative stress by generating reactive oxygen species (ROS), or antioxidants, which mitigate oxidative stress by scavenging ROS. The characteristics of N-[(6-chloro-3-pyridyl) methyl] warrant further investigation. The role of N'-cyano-N-methyl-acetamidine in this context requires experimental exploration, as there is a possibility that this substance may negatively affect enzymes and proteins, thereby leading to oxidative stress [29, 30]. Studies have demonstrated that the antioxidant properties of silver nanoparticles (AgNPs) inhibit free radicals and reduce oxidative stress. The ability of AgNPs to donate electrons and neutralize reactive oxygen species (ROS) is well-documented [31]. AgNPs contain natural antioxidants, such as vitamins C and E, and their combination with these antioxidants may offer enhanced protection against oxidative damage. AgNPs have been utilized in biomedical applications, particularly in drug delivery systems, where they can transport and release antioxidants at targeted sites within the body. This approach improves the bioavailability and efficacy of antioxidants. While AgNPs exhibit antioxidant properties, it is crucial to recognize that they can also demonstrate cytotoxic effects, particularly at elevated concentrations. The balance between their antioxidant benefits and potential toxicity requires careful consideration in various applications. In some cases, AgNPs are coated with antioxidant compounds to enhance their stability and mitigate potential toxicity. This combination may provide a safer and more effective platform for diverse applications. Ongoing research is exploring the use of AgNPs in the development of new



antioxidant-based therapies and materials, including their incorporation into wound dressings, food packaging, and cosmetic products to extend shelf life and improve product quality. In summary, the relationship between AgNPs and antioxidants is complex. While AgNPs can exhibit antioxidant properties, they are also used alongside antioxidants in various applications, potentially amplifying the overall antioxidant effect. However, their use must be carefully regulated and assessed to ensure safety and efficacy in specific contexts [32].

The elevation in the ALT and AST Levels may be attributed to cell death mechanisms associated with acetamiprid consumption in rats, which generates reactive metabolites and free radicals due to oxidative stress and degradation processes. These factors contribute to increased levels of ALT and AST. Typically, these enzymes are detoxified through conjugation with glutathione (GSH). However, in cases of overdose, the availability of GSH becomes limited, leading to liver damage, including mitochondrial impairment, oxidative stress, and nuclear DNA fragmentation [33]. Conversely, the decrease in AST and ALT enzyme levels in groups treated with silver nanoparticles may be attributed to the role of AgNPs in mitigating damage caused by free radicals or protecting tissues from adverse changes. Additionally, these nanoparticles may enhance the activity of enzymes responsible for detoxification [34].

The enhanced level of urea and creatinine are due to the alteration in the function of the kidney may be due to the action of pesticide. Metabolic deterioration of cells by the generation of toxic metabolites, metabolic byproducts, and pathological molecules leading to tissue damage are a consequence of these perturbations [35]. High levels of urea and creatinine can be harmful to tissue health and affect kidney functions. The elevation of urea and creatinine levels could do to deuterating effects of acetamiprid on oxidative stress and renal tissue. On the other hand, it could be due to a decreased glomerular filtration rate (because of glomerular arterioles' resistance, for example, or a low blood volume, both of which can diminish creatinine excretion and produce its retention in the body) [36]. The discovered increase and decrease in urea and creatinine levels in G4 and G5 could have been as a result of nanosilver antimicrobial effects which is well known as a strong antioxidant as well. Nanosilver seemingly shield cell membrane against toxicities of free radicals and in turn to improve the kidney function and maintain normal GRF [37].

In conclusion, the results of this study suggest that silver nanoparticles loaded with alcoholic extracts of *Lycium Shawii* have significant potential as antioxidant agents. The in vivo analysis showed that AgNPs, especially in the presence of bioactive compounds of *Lycium Shawii*, exhibited strong antioxidant activity and was able to modulate oxidative stress markers, providing protection from oxidative damage. Exposure to Acetamiprid resulted in a significant increase in oxidative stress as indicated by increased levels of oxidative stress markers and biomarkers (TOS, ALT, AST, urea, and creatinine), which was attenuated by treatment with AgNPs, indicating a protective effect of AgNPs against oxidative damage.

These results suggest that AgNPs may have therapeutic potential as an intervention for oxidative stress-related disorders such as cancer. However, further studies are required to understand the molecular mechanisms underlying the observed antioxidant effect and to evaluate the long-term safety and efficacy of AgNPs in clinical settings. These findings provide a solid foundation for further investigation into the role of silver nanoparticles in drug development and as dietary supplements.

## Author Contribution Statement

Meaad Nasser Hussein and Yusra Sebri abdulsaheb: Methodology, Investigation, Data curation, Original draft preparation. Ali NooryFajer: Supervision, Conceptualization, Writing- Reviewing and Editing.

## Acknowledgements

### *Availability of data and materials*

The data and materials that support the findings of this study are available from the corresponding author, upon reasonable request.

### *Ethical Approval*

This research protocol was evaluated and approved by Researches Ethics Committee of AL-Qadisiyah University, Iraq.

### *Conflict of interest*

The authors declare that they have no conflict of interests.

## References

1. Fuchs-Tarlovsky V. Role of antioxidants in cancer therapy. *Nutrition*. 2013;29(1):15-21. <https://doi.org/10.1016/j.nut.2012.02.014>.
2. Luo M, Zhou L, Huang Z, Li B, Nice EC, Xu J, et al. Antioxidant therapy in cancer: Rationale and progress. *Antioxidants*. 2022;11(6):1128. <https://doi.org/10.3390/antiox11061128>.
3. Fadhil J, Nada N, Wasan G. Total antioxidant capacity and homocysteine levels in obese women with polycystic ovary syndrome. *Int J Pharm Pharm Res*. 2017;8:78-86.
4. Adday Ali R, Reda A, Abbas HJ, Addai Ali H, Ali Naser N. Association of vitamin d and antioxidant minerals with interferon- $\lambda$ 1 (interleukin-29) level in type 2 diabetic patients. *Egypt J Vet Sci*. 2025;56(5):1075-80.
5. Kadhum DA, Mashkour MS, Ali HA, Parsian H. Role of interleukin-17 in the pathogenesis and progression of coronary artery disease. *Egyptian Journal of Medical Microbiology*. 2024;33(4):83-89. <https://doi.org/10.21608/ejmm.2024.308246.1295>.
6. Didier AJ, Stiene J, Fang L, Watkins D, Dworkin LD, Creeden JF. Antioxidant and anti-tumor effects of dietary vitamins a, c, and e. *Antioxidants*. 2023;12(3):632. <https://doi.org/10.3390/antiox12030632>.
7. Rogers B, Adams J, Pennathur S. *Nanotechnology: understanding small systems*. Crc Press; 2007 Nov 29.
8. McNeil SE. Nanotechnology for the biologist. *J Leukoc Biol*. 2005;78(3):585-94. <https://doi.org/10.1189/jlb.0205074>.
9. Bawa R, Audette GF, Rubinstein I, editors. *Handbook of*

- clinical nanomedicine: nanoparticles, imaging, therapy, and clinical applications. CRC Press; 2016 Feb 22.
10. Hadi ZA, Odda AH, Jawad AF, Al-Tu'ma FJ. Design and development of fe3o4@ prussian blue nanocomposite: Potential application in the detoxification of bilirubin. *Asian Pac J Cancer Prev.* 2023;24(8):2809. <https://doi.org/10.31557/APJCP.2023.24.8.2809>.
11. Harstad S, Hunagund S, Boekelheide Z, Hussein ZA, El-Gendy AA, Hadimani RL. Gd-Based Magnetic Nanoparticles for Biomedical Applications. In *Magnetic Nanostructured Materials*. Elsevier; 2018. p. 137-55.
12. Singh J, Dutta T, Kim KH, Rawat M, Samddar P, Kumar P. 'Green' synthesis of metals and their oxide nanoparticles: Applications for environmental remediation. *J Nanobiotechnology.* 2018;16(1):84. <https://doi.org/10.1186/s12951-018-0408-4>.
13. Shipelin V, Gmshinski I, Khotimchenko S, editors. Risk assessment of silver nanoparticles. IOP Conference Series: Materials Science and Engineering; 2015: IOP Publishing.
14. Birben E, Sahiner UM, Sackesen C, Erzurum S, Kalayci O. Oxidative stress and antioxidant defense. *World Allergy Organ J.* 2012;5(1):9-19. <https://doi.org/10.1097/WOX.0b013e3182439613>.
15. J Al-Tu F, A Ibraheem A, H Al-Kayatt T. Adiponectin and lipid profile levels in type-2 diabetic obese patients in Karbala Province. *karbala journal of pharmaceutical sciences.* 2011;2(2):157-66.
16. Numan HA, Almohanna TG, Almayali MQ, Ali RA, Taha HKA, Ali HA. The effect of clusterin level as a potential marker in women with polycystic ovary syndrome. *Edelweiss Appl Sci Technol.* 2024;8(5):616-23. <https://doi.org/10.55214/25768484.v8i5.1725>.
17. Yao X-h, Min H, Lü Z-h, Yuan H-p. Influence of acetamiprid on soil enzymatic activities and respiration. *Eur J Soil Biol.* 2006;42(2):120-6. <https://doi.org/10.1016/j.ejsobi.2005.12.001>.
18. Kimura-Kuroda J, Komuta Y, Kuroda Y, Hayashi M, Kawano H. Nicotine-like effects of the neonicotinoid insecticides acetamiprid and imidacloprid on cerebellar neurons from neonatal rats. *PLoS One.* 2012;7(2):e32432. <https://doi.org/10.1371/journal.pone.0032432>.
19. Al-Lami HB, Al-Tu'ma FJ, Al-Safi WG. Association between anti-müllerian hormone and other biomarkers with ovarian function in polycystic ovarian syndrome of iraqi women. *J contemp med sci.* 2020;6(4). <https://doi.org/10.22317/jcms.v6i4.824>.
20. Imamura T, Yanagawa Y, Nishikawa K, Matsumoto N, Sakamoto T. Two cases of acute poisoning with acetamiprid in humans. *Clin Toxicol (Phila).* 2010;48(8):851-3. <https://doi.org/10.3109/15563650.2010.517207>.
21. Ahmed TA, Al Naemi H. Biological activities of lycium shawii leaves extract. *Int J Pharm Biol Arch.* 2010;3:697-700.
22. Erdoğan Ü. Ultrasonic assisted propolis extraction: Characterization by atr-ftir and determination of its total antioxidant capacity and radical scavenging ability. *Int J Second Metab.* 2023;10(2):231-9. <https://doi.org/10.21448/ijsm.1167773>.
23. Diken HO, Demirtas B, Kaya HK. Effects of kefir on paraoxanase activity (pon1), total antioxidant status (tas), total oxidant status (tos), and serum lipid profiles in smokers and non-smokers. *Food Sci Technol.* 2022;42:e92721.
24. Verdian A. Apta-nanosensors for detection and quantitative determination of acetamiprid—a pesticide residue in food and environment. *Talanta.* 2018;176:456-64. <https://doi.org/10.1016/j.talanta.2017.08.070>.
25. Kozitsina A, Dedeneva S, Shalygina ZV, Okhokhonin A, Chizhov D, Matern A, et al. Determination of urea and creatinine by chronoamperometry. *J anal chem.* 2014;69:758-62. <https://doi.org/10.1134/S1061934814080048>.
26. Alwhibi MS, Soliman DA, Awad MA, Alangery AB, Al Dehaish H, Alwasel YA. Green synthesis of silver nanoparticles: Characterization and its potential biomedical applications. *Green Process Synth.* 2021;10(1):412-20. <https://doi.org/10.1515/gps-2021-0039>.
27. Al-Gburi SA. Green synthesis of silver nanoparticles. 2018.
28. Sakr M, Adly MS, Gar Alalm M, Mahanna H. Effective removal of acetamiprid and eosin y by adsorption on pristine and modified mil-101 (fe). *Environ Sci Pollut Res.* 2024;1-25.
29. Annabi E, Ben Salem I, Abid-Essefi S. Acetamiprid, a neonicotinoid insecticide, induced cytotoxicity and genotoxicity in pc12 cells. *Toxicol Mech Methods.* 2019;29(8):580-6. <https://doi.org/10.1080/15376516.2019.1624907>.
30. Hussein KM, Ali HA. Association between interleukin-39 (il-39) with hormonal and metabolic changes in women with polycystic ovarian syndrome. *Anaesth Pain Intensi.* 2024;28(6):1035-42. <https://doi.org/10.35975/apic.v28i6.2612>.
31. Docea AO, Calina D, Buga AM, Zlatian O, Paoliello M, Mogosanu GD, et al. The effect of silver nanoparticles on antioxidant/pro-oxidant balance in a murine model. *Int J Mol Sci.* 2020;21(4):1233. <https://doi.org/10.3390/ijms21041233>.
32. Arozal W, Monayo ER, Barinda AJ, Perkasa DP, Soetikno V, Nafrialdi N, et al. Protective effects of silver nanoparticles in isoproterenol-induced myocardial infarction in rats. *Front Med.* 2022;9:867497. <https://doi.org/10.3389/fmed.2022.867497>.
33. Pravinson M, Pirasath S, Gheteeswaran S, Uthayakumaran S. Acute poisoning with acetamiprid, a type of neonicotinoid insecticide causing severe lactic acidosis: A case report. *SAGE Open Med Case Rep.* 2021;9:2050313X211059296.
34. Wen L, Li M, Lin X, Li Y, Song H, Chen H. Agnps aggravated hepatic steatosis, inflammation, oxidative stress, and epigenetic changes in mice with nafld induced by hfd. *Front Bioeng Biotechnol.* 2022;10:912178.
35. Turan F, Ergenler A. Investigation of the genotoxic effect of acetamiprid in cyprinus carpio using the micronucleus analysis and the comet assay. *Turk J Marit Marine Sci.* 2022;8(2):80-9. <https://doi.org/10.52998/trjmms.1037906>.
36. Zhang X, Huang Y, Chen WJ, Wu S, Lei Q, Zhou Z, et al. Environmental occurrence, toxicity concerns, and biodegradation of neonicotinoid insecticides. *Environ Res.* 2023;218:114953. <https://doi.org/10.1016/j.envres.2022.114953>.
37. Alshubaily FA, Jambi EJ, Khojah SM, Balgoon MJ, Al-Zahrani MH, Alkhatabi NA. Studying the effect of silver nanoparticles synthesized by ulva fasciata aqueous extract against liver toxicity induced by ccl4 in rats. *Journal of Nature and Science of Medicine.* 2020;3(3):182-8. [https://doi.org/10.4103/JNSM.JNSM\\_2\\_20](https://doi.org/10.4103/JNSM.JNSM_2_20).



This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License.